

Jury Member Report – Doctor of Philosophy thesis.

Name of Candidate: Valentina Ekimova

PhD Program: Petroleum Engineering

Title of Thesis: Experimental modeling of gas hydrates interaction with a salt solution in permafrost

Supervisor: Dr. Evgeny Chuvilin

Name of the Reviewer: Jeffrey Priest

I confirm the absence of any conflict of interest (Alternatively, Reviewer can formulate a possible conflict)	Date: 18-09-2022
--	-------------------------

The purpose of this report is to obtain an independent review from the members of PhD defense Jury before the thesis defense. The members of PhD defense Jury are asked to submit signed copy of the report at least 30 days prior the thesis defense. The Reviewers are asked to bring a copy of the completed report to the thesis defense and to discuss the contents of each report with each other before the thesis defense.

If the reviewers have any queries about the thesis which they wish to raise in advance, please contact the Chair of the Jury.

Reviewer's Report

Reviewers report should contain the following items:

- Brief evaluation of the thesis quality and overall structure of the dissertation.
- The relevance of the topic of dissertation work to its actual content
- The relevance of the methods used in the dissertation
- The scientific significance of the results obtained and their compliance with the international level and current state of the art
- The relevance of the obtained results to applications (if applicable)
- The quality of publications

The summary of issues to be addressed before/during the thesis defense

This thesis deals with influence of salt solution on the dissociation of gas hydrate. The research deals primarily conducting laboratory experiments where various sand specimens are formed with methane gas hydrate and ice which are subsequently inserted into salt solutions and track the evolution of temperature and salt ions through the sample as a function of pressure, temperature, and salt concentration.

The thesis is structured through 8 chapters. The thesis is appropriately formatted in terms of thesis structure. The first chapter defines the problem statement, the research goals and the outline of the thesis. This is followed by two chapters that highlight the current state of knowledge of gas hydrates in permafrost and how salt migration occurs and is modelled in frozen sediments. These chapters are very in-depth, although there is much discussion on aspects that are not relevant to the actual research conducted. It would have been beneficial to have appropriate subsections to identify discussion of new and different ideas. At present these two chapters very 'expansive' which makes it somewhat difficult to follow what is important, relevant, and how it differs within the text. Chapter 5 discusses the experimental program, sediment characteristics, and the various methods for determining key observation measurements. I found this chapter to be somewhat repetitive. The test methods were described but were somewhat light of detail. It would have been useful to have a table of all tests that were to be carried out and what the different tests were for. Although tests were described they were not clear. This is compounded in subsequent chapters where results are introduced but changes in hydrate saturations, temperatures, and pressures etc are not clear. In this chapter also terms are introduced that are not discussed etc. Also, some specific tests are highlighted that do not follow the methods described. Chapter 6 and 7 presents the results of testing and hypothesis for observations. As highlighted details were somewhat vague of specific tests with tables of tests that only appear to cover some of the test's results presented. Better clarity in detailing all tests carried out and providing more clear tables of all tests would have helped in understanding results better. Chapter 8 discusses conceptual models, which are generally clear although figure numbering was wrong. Finally, a conclusion chapter is provided, where some conclusions were more statements of what was carried out rather than clear conclusions s are offered.

The research carried out addresses some key aspects of developing oil and gas resources in the Arctic, namely how destabilization of hydrate might occur through salt solution migration, either arising from drilling fluids or naturally through sea water. The research addresses the identified issues and provides some results, with some discussion of the impact of the results gained.

The PhD student has done a great deal of high-quality work as evidenced by the 6 journal publications (5 geoscience and 1 Energy & Fuels) and 4 conference proceedings. These are appropriate journals and conferences for the work carried out.

Overall, the research program addresses the stated objectives and of high quality. However, the thesis does require some work to really make the research stand out clearly. I found the referencing style to be haphazard and did not follow one 'standard' but mixed and matched. There were several figures that were not named correctly in the text. More clear and detailed discussion of test methodology would go a long way to understanding the complexity and magnitude of testing conducted, along with complete list of tests and test conditions.

I have appended an annotated thesis. This contains many questions etc that I would have in a defense, but also highlights some key things that need to be addressed. I feel these should be done prior to defense and believe that the student can make these changes in the month available before the scheduled defense.

Provisional Recommendation

I recommend that the candidate should defend the thesis by means of a formal thesis defense

I recommend that the candidate should defend the thesis by means of a formal thesis defense only after appropriate changes would be introduced in candidate's thesis according to the recommendations of the present report

The thesis is not acceptable and I recommend that the candidate be exempt from the formal thesis defense



Skolkovo Institute of Science and Technology

Skolkovo Institute of Science and Technology

EXPERIMENTAL MODELING OF GAS HYDRATES INTERACTION WITH A SALT
SOLUTION IN PERMAFROST

Doctoral Thesis

by

VALENTINA EKIMOVA

DOCTORAL PROGRAM IN PETROLEUM ENGINEERING

Supervisor

Leading Research Scientist Dr. Evgeny Chuvilin

Moscow 2022

© Ekimova Valentina 2022

I hereby declare that the work presented in this thesis was carried out by myself at Skolkovo Institute of Science and Technology, Moscow, except where due acknowledgement is made, and has not been submitted for any other degree.

Candidate (Valentina Ekimova)

Supervisor (Leading Research Scientist Dr.
Evgeny Chuvilin)

Abstract

Today, one of the most promising area for oil and gas production is the Arctic region. Though, the hydrocarbon production in this region is frequently associated with many complications, among them is the existence of sensitive permafrost component – gas hydrates. Gas hydrates destabilization may occur when thermobaric conditions of their stability change or as a result of inhibition by various chemical compounds, in particular by salts. Consequently, intensive methane emission can happen.

To date, issues related to the influence of geochemical conditions on methane hydrate stability (particularly migration of salt solutions) in the sediments pore space at negative temperatures have turned out to be poorly understood. This research is dedicated to experimental study of the mechanism and regularities of the interaction of gas hydrates with salt solutions in the pore space of hydrate-saturated sediments under various thermobaric conditions. The object of the study was sandy sediments, as well as the ones obtained during drilling in the area of permafrost distribution, where the existence of gas hydrates is assumed by a number of indirect signs.

During the experiments, it was found that the migration of salts in frozen sediments, containing gas hydrates stimulates the hydrate decomposition in pore space of sediments, and can accelerates ice thawing. As a result, a dissociation front of porous gas hydrates was observed.

It has been found that salt migration proceeds more intensively in frozen hydrate-saturated sediment than in frozen non-hydrate-saturated ones. At the same time, a significant decrease in the temperature of a frozen hydrate-containing sample was recorded during the migration and accumulation of salts, while a less significant decrease in temperature was observed in frozen sediment. It is shown that the most important factor determining the intensity of salt ions migration in frozen sandy sediment, containing gas hydrates, is gas pressure, in contrast to frozen hydrate-free sediments, where the influence of pressure was not so significant.

It was revealed that during salt migration in frozen hydrate-saturated sediments, the intensity of methane hydrate dissociation rises with an increase in ambient temperature, salt solution concentration, and with a decrease in the amount of clay particles. Moreover, the activity of hydrate dissociation decreases in the series of salt solution composition: $\text{MgCl}_2\text{-NaCl-CaCl}_2\text{-KCl-Na}_2\text{SO}_4$. On the basis of experimental data, the salt ion flux was calculated, and the critical concentration of complete dissociation of pore gas hydrate was obtained.

A scheme of phase transformations and temperature changes in the pore space of frozen hydrate-saturated sediment during their interaction with salt solutions at gas pressures above and below equilibrium is proposed. A schematic model of the contact of natural and technogenic salt solutions with frozen hydrate-saturated strata in continental conditions and on the shelf is also given.

Publications

Journal publications:

1. Chuvilin E., **Ekimova V.**, Davletshina D., Krivokhat E., Shilenkov V., Bukhanov B. Pressure factor Influence on the Salt Migration in frozen hydrate-saturated sediments. *Experimental Modeling. Energy & Fuels*. 2022.
2. Chuvilin E., **Ekimova V.**, Davletshina D., Krivokhat E., Shilenkov V., Bukhanov B. Migration of Salt Ions in Frozen Hydrate-Saturated Sediments: Temperature and Chemistry Constraints. *Geosciences*. 2022 Jul; 12 (7), 276. <https://doi.org/10.3390/geosciences12070276>, WoS/Scopus indexed journal. Impact Factor: 2.076.
3. Chuvilin E., **Ekimova V.**, Davletshina D., Krivokhat E., Shilenkov V., Bukhanov B. Temperature Variation during Salt Migration in Frozen Hydrate-Bearing Sediments: Experimental Modeling. *Geosciences*. 2022 Jun; 12 (7), 261. <https://doi.org/10.3390/geosciences12070261>, WoS/Scopus indexed journal. Impact Factor: 2.076.
4. Chuvilin E, **Ekimova V**, Davletshina D, Sokolova N, Bukhanov B. Evidence of Gas Emissions from Permafrost in the Russian Arctic. *Geosciences*. 2020 Sept: 10 (10), 383. <https://doi.org/10.3390/geosciences10100383>, WoS/Scopus indexed journal. Impact Factor: 2.076.
5. Chuvilin E, Davletshina D, **Ekimova V**, Bukhanov B, Shakhova N, Semiletov I. Role of Warming in Destabilization of Intrapermafrost Gas Hydrates in the Arctic Shelf: Experimental Modeling. *Geosciences*. 2019 Sept; 9: 407. <https://doi.org/10.3390/geosciences9100407>, WoS/Scopus indexed journal. Impact Factor: 2.076.

6. Chuvilin E, **Ekimova V**, Bukhanov B, Grebenkin S, Shakhova N, Semiletov I. Role of salt migration in destabilization of intra permafrost hydrates in the Arctic Shelf: Experimental modeling. *Geosciences*. 2019 Apr: 9(4), 188. <https://doi.org/10.3390/geosciences9040188>, WoS/Scopus indexed journal. Impact Factor: 2.076.

Conference proceedings:

1. **Ekimova V.**, Chuvilin E., Davletshina D., Bukhanov B., Krivokhat E., Shilenkov V. Salt transfer in frozen hydrate-containing sediments during their interaction with salt solutions. In the Proceeding of IV Conference of Geocryologists of Russia "Monitoring in permafrost", Moscow, Russia, June 14 - 17, 2022, pp. (published in Russian).

2. **Ekimova V**, Chuvilin E, Bukhanov B, Grebenkin S, Shakhova N, Semiletov I. Evaluation of the role of salt migration in the destabilization of intrapermafrost gas-hydrate formations on the Arctic shelf on the basis of the experimental modeling. In Proceedings of the IX International Scientific and Practical Conference "Marine Research and Education (MARESEDU-2020)", Moscow, Russia, 2020, pp. 59-62 (published in Russian).

3. **Ekimova V**, Chuvilin E, Bukhanov B, Grebenkin S, Shakhova N, Semiletov I. Experimental modeling of dissociation of intrapermafrost gas hydrates caused by salt transfer. In Proceeding of the Conference "Global Issues in the Arctic and Antarctic", Archangelsk, Russia, 2020, pp. 101-104 (published in Russian).

4. Chuvilin E., Bukhanov B., **Ekimova V**. Experimental modeling of interaction between salt solutions and frozen sediments containing gas hydrates // The 8th International Conference on Gas Hydrates. Beijing, China, 2014.

Acknowledgements

This work would be impossible without my Research Supervisor Leading Research Scientist of Skoltech Dr. Evgeny Chuvilin. I am grateful for guidelines, and all valuable recommendations.

I would like to extend my sincere thanks to my Ph.D. committee members – Prof. Mikhail Spasennykh, and Dr. Vladimir Istomin for their suggestions and insightful comments during my study and research.

I am deeply grateful to my colleagues, who were ready to assist at every stage of the research work and helped in conducting the experiments: Senior Research Scientist Boris Bukhanov, Engineers Natalia Sokolova, Dinara Davletshina, Sergei Grebenkin, Daria Sergeeva, Ph.D. student Vladimir Shilenkov, Ms student Ekaterina Krivokhat, Ms student Elena Pankratova from Skoltech University, Gleb Berloff from Warwick University.

I want to give my deepest appreciation to the Skolkovo Institute of Science and Technology for amazing 4 years of advanced research, innovations, enjoyable lectures, and extracurricular activity.

I would like to thank all Jury committee members – for their time, recommendations and comments for the work perfection.

Finally, a special appreciations to my family and friends for their belief in me, unwavering support, and inspiration in conducting research work.

The research was supported by the Russian Science Foundation (grants No. 22-17-00112, 21-77-10074).

Table of Contents

Abstract.....	3
Publications	5
Acknowledgements	7
Table of Contents	8
List of Figures.....	10
List of Tables	17
Chapter 1. Introduction.....	18
1.1. Relevance. Statement of the problem	18
1.2. Goal.....	20
1.3. Objectives	20
1.4. Novelty.....	21
1.5. Outline of the Thesis	22
Chapter 2. Arctic permafrost and problems of hydrocarbon fields development in the Arctic	23
2.1. Permafrost-geological structure of the Russian Arctic region	23
2.2. Peculiarities of hydrocarbon deposits development in areas of permafrost distribution	30
Chapter 3. Conditions for gas hydrate existence and formation in permafrost	35
3.1. Characteristics of gas hydrate as a substance.....	35
3.2. Conditions for the existence of permafrost associated gas hydrate and possible causes of their destabilization	45
Chapter 4. The study of salt migration processes in frozen sediments.....	58
4.1. Experimental research of salt migration processes in ice-containing sediments	61
4.2. Existing physico – chemical models of salt transfer processes in sediment media	69
Chapter 5. Method of experimental study of salt ions migration in hydrate-saturated sediments	76
5.1. Experimental setup and preparation technology of hydrate saturated sediments	77
5.2. Sediment characteristics used in experimental simulation.....	81
5.3. Technique for studying the interaction of frozen hydrate-containing sediments with saline solutions in non-equilibrium and equilibrium conditions	83
5.4. Method of the temperature field change investigation during the contact of frozen hydrate saturated sediment with salt solution.....	88
5.5. Method for assessing the salt content in sediment samples	91
5.6. Processing of the experimental data	97
Chapter 6. Mechanism and dynamics of salt transfer processes in hydrate-bearing sediments.....	99
6.1. Influence of hydrate presence on the salt migration in frozen sediments	100

6.2. The interaction of frozen hydrate saturated frozen sediments with salt solutions in time	103
6.3. Temperature changes as a result of salt migration in frozen hydrate-saturated sediments	107
5.4. Mechanisms of salt migration in frozen hydrate containing sediments	117
Chapter 7. Influence of the different factors on the process of salt migration in frozen hydrate saturated sediments	125
7.1. Salt and mass transfer in sediment saturated with gas hydrate, depending on pressure	125
7.2. Temperature influence on the process of hydrate-bearing sediment interaction with saline solutions	132
7.3. Influence of salt solution concentration on the intensity of salt migration and hydrate dissociation in hydrate saturated sediments	137
7.4. The process of interaction of hydrate-containing sediments with a salt solution of various chemical composition.....	147
7.5. Sediment dispersion influences the process of salt migration in hydrate saturated sediments	153
Chapter 8. Concept models of hydrate interaction with natural and technical salt solution in hydrate saturated collectors in permafrost.....	167
Chapter 9. Conclusion	171
Bibliography	176

List of Figures

Figure 1. Map of the permafrost distribution in the Arctic zone of the Russian Federation [71].	27
Figure 2. Simulation of the zero isotherm propagation on the Arctic shelf [80].	29
Figure 3. Structure of gas hydrates [139].	36
Figure 4. Different crystalline arrangement of hydrates: a – structure I, b – structure II, c – structure H and d – host cages of water. Dense line is the hydrate unit-cell [145].	37
Figure 5. Gas hydrate burning (photo by Dinara Davletshina).	38
Figure 6. Stability of CH ₄ hydrate curve (build in HydraFlash): GH – gas hydrate; liq – liquid water.	39
Figure 7. Shift of gas hydrate equilibrium by different methods of methane extraction [144].	40
Figure 8. Scheme of bulk hydrate self-preservation at the negative temperature (picture obtained and analyzed through extensive cryo-FE-SEM imaging [168].	41
Figure 9. Different hydrate inhibitors and scheme of their work principle [145].	42
Figure 10. Hydrate decomposition in water by NaCl ions migration in time by mathematical modeling (red – methane, green – Na ⁺ , yellow – Cl ⁻) [201].	44
Figure 11. Existence of methane hydrate in natural conditions (modified from [17,223]).	46
Figure 12. Hydrate distribution in sediments [12].	48
Figure 13. Schematic map of registered gas emanations in the Russian Arctic [236].	54
Figure 14. Scheme of a chemical reaction during the melting of ice due to the action of NaCl ions [298].	59
Figure 15. Phase diagram of NaCl aqueous solution (modified after [305]).	60
Figure 16. Equipment (pressure cell) for saturation of sediment by hydrate.	78
Figure 17. The appearance of frozen sand saturated with pore hydrate after extraction from the pressure cell.	80
Figure 18. Distribution of water content, porosity, and hydrate coefficient by the length of frozen hydrate saturated sediment samples.	80

Figure 19. Scheme of the installation for the interaction of frozen hydrate-saturated and non-hydrate-saturated sediments with salt solutions under pressure.	86
Figure 20. Scheme of Kriolab Tbf system for temperature field change investigation (a) and picture of the frozen hydrate saturated sand sample with temperature sensors (b). 1 - frozen hydrate-saturated sediment; 2 - solution container; 3 - NaCl solution; 4 - temperature sensors; 5 - thermistor streamer; 6 - branch box; 7 - fan; 8 - USB cable; 9 - PC with Kriolab Tbf software; 10 - cooling chamber. Modified after [309].	89
Figure 21. The water activity of non-saline sand at the various water content.	92
Figure 22. Distribution of pore moisture activity from sand salinity (a) and salt concentration in pore moisture (b).	93
Figure 23. The water activity from NaCl concentration in water solution (HydraFlash).	94
Figure 24. Comparison of the measurement results of the Na ⁺ concentration by a flame photometer and a conductometer.	95
Figure 25. Pore moisture activity versus Na ⁺ ion concentration.	95
Figure 26. Effect of gas pressure on the accumulation of salt ions (Na ⁺) in frozen sand (fine sand -1) samples (W=12%) containing (solid line) and not containing (dashed line) gas hydrates 4 hours after interaction with 0.2 N NaCl solution (t=-6 °C).	101
Figure 27. Accumulation of Na ⁺ ions (A) and changes in the hydrate coefficient (B) in samples of frozen hydrate-containing sand-1 (K _h ⁱⁿ = 0.6, W=12%) at different times after interaction with NaCl solution 0.1 N (P=0.1 MPa, t=-6 °C).	104
Figure 28. Experimental evaluation of the decomposition front of porous gas hydrates and the thawing front in frozen hydrate-containing sand samples (sand-1, W = 12%) when interacting with 0.1 N NaCl salt solution at a temperature -6°C and pressure 0.1MPa.	105
Figure 29. The appearance of frozen sand samples (fine sand 4, W=12%) before (A) and after interaction (B) with 0.1 NaCl solution at -6°C.	106
Figure 30. Average Na ⁺ flux change in time during interaction of frozen hydrate saturated sediment (fine sand-1) with NaCl solution of 0.1N at -6 °C and 0.1 MPa.	107

Figure 31. Dynamics of temperature change in time in the near-contact area (2.5 cm from the contact) in frozen sand samples (fine sand-2) interacting with 0.4 N saline solution at atmospheric pressure: 1 – non-hydrate-containing frozen sample (dotted line); 2 - hydrate-containing (solid line); 3 - initial temperature (t_{in}) -6.5 °C (dashed line).	109
Figure 32. Temperature distribution along the length of a frozen sample (fine sand-2), not containing (1) and containing gas hydrate (2) after 0.6 h of interaction with 0.4 N. NaCl solution at atmospheric pressure. The dotted line - 3 shows the initial temperature $t_{in} = -6.5^{\circ}\text{C}$	110
Figure 33. Dynamics of temperature change in time in a frozen hydrate-containing sample ($S_h^{in}=38\%$, $S_i^{in} =15\%$) on the distance 2.5; 3.5; 5.5 and 7.0 cm from the contact zone with 1 N NaCl solution ($t_{in} = -6.5^{\circ}\text{C}$, $P=0.1\text{ MPa}$).	111
Figure 34. Change in the temperature distribution along the length of the frozen hydrate-saturated sample fine sand-2 ($S_h^{in}=41\%$, $S_i^{in} =12\%$) at the stage of cooling in the first hour of interaction with 1 N NaCl solution ($t_{in}=-6.5^{\circ}\text{C}$, $P=0.1\text{MPa}$).	113
Figure 35. Dynamics of temperature changes in time of frozen hydrate-containing samples (fine sand-2) in the near-contact area (2.5 cm from the contact) interacting with NaCl solution of various concentrations (0.2-1N) ($t_{in} = -6.5^{\circ}\text{C}$, $P=0.1\text{ MPa}$).	114
Figure 36. Influence of the contact solution concentration on the maximum temperature drop ($\Delta t=t_{in}-t_{min}$) in frozen hydrate saturated fine sand-2 (C_{sol} from 0.1 to 1 N, $t_{in} = -6.5^{\circ}\text{C}$, $P=0.1\text{ MPa}$).	115
Figure 37. Influence of contact solution concentration on the rate of temperature decrease (Vt) of frozen hydrate-saturated fine sand-2 (C_{sol} from 0.1 to 1 N, $t_{in} = -6.5^{\circ}\text{C}$, $P=0.1\text{ MPa}$).	116
Figure 38. Influence of concentration on the time taken for the temperature of the sample to return to the initial value (T_r) after the cooling period caused by the interaction of hydrate-containing frozen sand with saline solution ($t_{in} = -6.5^{\circ}\text{C}$, $P=0.1\text{ MPa}$).	116
Figure 39. Scheme of phase-transformation of frozen hydrate saturated sediments during salt migration: (a) metastable conditions (experimental pressure $P_{exp} < \text{equilibrium pressure } P_{eq}$); (b) stable conditions ($P_{exp} \geq P_{eq}$).	119
Figure 40. Scheme of temperature changes during the interaction of a frozen hydrate-containing sample with saline solution.	122

Figure 41. Scheme of temperature changes in a frozen hydrate-containing sample in the process of salt transfer, phase transformations, and changes in ice and hydrate saturation along the height of the investigated sample.	123
Figure 42. Modeling P-T conditions of methane hydrate stability in the HydraFlash program and thermobaric conditions for conducting experiments on the migration of salt ions in frozen hydrate-saturated sediments (T1-T6).....	126
Figure 43. The effect of pressure on the accumulation of salt ions (Na^+) (a) and the decrease in the hydrate coefficient (K_h) (b) in samples of frozen hydrate saturated fine sand-2 ($K_{hin} \sim 0.3$, $W=12\%$) at non-equilibrium and equilibrium pressure 4 hours after the start of interaction with a frozen NaCl solution 0.1 N ($t= -6^\circ\text{C}$).	127
Figure 44. Influence of gas pressure on the flux density of Na^+ ions through the cross section of a frozen hydrate-containing sample 4 hours after the start of interaction with a frozen salt solution 0.1 N NaCl ($t= -6^\circ\text{C}$).....	128
Figure 45. Dependence of the critical concentration (C_{cr}) causing decomposition of pore hydrate in a frozen sand sample on gas pressure ($t= -6^\circ\text{C}$, $C_{sol}=0.1$ N).	129
Figure 46. Altitude distribution of a frozen hydrate-containing sample: (A) salt concentrations and hydration coefficient; (B) – critical pressure of dissociation of methane hydrate (concentration of contacting NaCl solution 0.1 N, $t= -6^\circ\text{C}$).	130
Figure 47. Effect of temperature on the accumulation of salt ions (Na^+) in frozen hydrate-containing samples (fine sand-2) after 2 hours of interaction with 0.2 N NaCl solution.	133
Figure 48. The effect of temperature (-3, -6, -20°C) on the decrease in the hydration coefficient (K_h) in samples of frozen hydrate-containing fine sand-2 ($K_{hin} \sim 0.4$, $W=12\%$) 2 hours after the start of interaction with frozen 0.2 N NaCl salt solution.	134
Figure 49. Effect of temperature on the average Na^+ flux into a frozen hydrate-containing sample (fine sand-2) for 4 hours of interaction with a frozen salt solution of 0.1N NaCl ($P=0.1\text{MPa}$).....	135
Figure 50. Effect of temperature on the critical concentration (C_{cr}) causing decomposition of pore hydrate in a frozen sand sample (fine sand-2, $P=0.1$ and 4MPa , $C_{sol}=0.1\text{N}$).	136
Figure 51. Thermobaric conditions of hydrate stability in the water-NaCl solution of different concentration simulated in “HydraFlash” software.....	138

Figure 52. Effect of contact solution concentration on the accumulation of salt ions (Na^+) in frozen hydrate-containing samples (fine sand-2) after 4 hours of interaction with NaCl solution ($t=-6^\circ\text{C}$, $P=0.1\text{ MPa}$).....	140
Figure 53. Influence of NaCl contact solution concentration on the distribution of hydrate coefficient (K_h) along the length of frozen hydrate-containing samples (fine sand-2) ($K_h^{\text{in}} \sim 0.4$, $W=12\%$) after 4 hours of interaction ($t=-6^\circ\text{C}$, $P=0.1\text{ MPa}$).	141
Figure 54. Influence of the contact solution concentration on the accumulation of salt ions (Na^+) in frozen hydrate-containing samples (fine sand-2) interacting for 4 hours under a pressure of 4.0 MPa with a NaCl solution ($t=-6^\circ\text{C}$).	142
Figure 55. Effect of contact solution concentration on distribution of hydrate coefficient (K_h) along the height of frozen hydrate-containing sample (fine sand-2, $K_h^{\text{in}} \sim 0.4$, $W=12\%$) after 4 hours of the interaction with frozen NaCl salt solution ($t=-6^\circ\text{C}$, $P=4.0\text{ MPa}$).	143
Figure 56. Influence of contact solution concentration on the average Na^+ flux in frozen hydrate-saturated sample (fine sand-2) after 4 hours of the interaction ($t= -6^\circ\text{C}$, $P=0.1\text{ MPa}$ and 4.0 MPa).	144
Figure 57. Influence of a chemical composition of salt solution (Na_2SO_4 , KCl , CaCl_2 , NaCl , MgCl_2) with a concentration of 5% on the conditions of methane hydrate stability in the range of negative temperatures from 0 to -20°C (calculations were made in the “HydraFlash” software).....	148
Figure 58. Accumulation of Na^+ ions in samples of frozen hydrate saturated sand (fine sand-2, $W=12\%$) after 4 hours of contact with 0.1N NaCl solution of different composition ($t= -6^\circ\text{C}$, $P=0.1\text{MPa}$).....	149
Figure 59. The influence of the chemical composition of the contact salt solution on the decrease in the hydrate coefficient (K_h) in frozen hydrate-containing samples (fine sand-2, $K_h^{\text{in}} \sim 0.5$, $W=12\%$) after 4 hours of interaction with the frozen salt solution ($t=-6.5^\circ\text{C}$, $C_{\text{sol}}=0.1\text{N}$, $P=0.1\text{MPa}$).....	151
Figure 60. The influence of the chemical composition of the salt contact solution on the salt flux into frozen hydrate-containing sample (fine sand-2) after 4 hours of interaction with a frozen salt solution ($t=-6^\circ\text{C}$, $C_{\text{sol}}=0.1\text{N}$, $P=0.1\text{MPa}$).	152

Figure 61. Influence of kaolinite content in fine-grained sand (fine sand-2) on the accumulation of Na ⁺ ions in frozen hydrate-saturated samples 4 hours after interaction with 0.1 N NaCl solution at -6 °C.....	155
Figure 62. Effect of kaolinite content in sand (fine sand-2) on hydrate dissociation in a frozen hydrate-saturated sample ($K_h^{in} \sim 0.5$, $W=12\%$) 4 hours after the start of interaction with a frozen salt solution of 0.1 N NaCl at -6 °C.	155
Figure 63. Influence of kaolinite content in sand (fine sand-2) on moisture accumulation in a frozen hydrate-saturated sample ($W=12\%$) after 4 hours of interaction with salt solution of 0.1 N NaCl at -6 °C.....	156
Figure 64. Effect of kaolinite content in sand (fine sand-2) on the average water flux through a frozen hydrate-bearing sample ($W=12\%$) after 4 h of interaction with a frozen salt solution of 0.1 N NaCl at -6 °C.	157
Figure 65. Influence of the mineralogical composition of clay in quartz sand (fine sand-2 with 5% additives) on the accumulation of Na ⁺ ions in frozen hydrate-saturated samples after 4 hours of interaction with 0.1 N NaCl solution at $t=-6^{\circ}C$: A – kaolinite; B – montmorillonite.....	158
Figure 66. Influence of the mineralogical composition of clay in quartz sand (fine sand-2) on the decrease in the hydration coefficient (K_h) in frozen hydrate-containing samples ($K_h^{in} \sim 0.4$, $W=12\%$) 4 hours after the start of interaction with frozen salt solution 0.1 N NaCl at $t= -6^{\circ}C$: A – kaolinite; B, montmorillonite.	158
Figure 67. Influence of the mineralogical composition of clay in quartz sand (fine sand-1) on the accumulation of moisture in a frozen hydrate-saturated sample ($W=12\%$) after 4 hours of interaction with a frozen salt solution of 0.1 N NaCl at -6 °C: A - kaolinite; B - montmorillonite.	159
Figure 68. Influence of the content of clay particles of various compositions (kaolinite and montmorillonite) in quartz sand (fine sand-2) on the average flux density of Na ⁺ ions into a frozen hydrate-containing sample over 4 hours of interaction with a frozen salt solution of 0.1N NaCl ($p=0.1MPa$, $t= -6^{\circ}C$): A – kaolinite; B – montmorillonite.....	160
Figure 69. Influence of the content of clay particles of different mineral composition in quartz sand (fine sand-2) on the critical concentration (C_{cr}) causing the decomposition of pore hydrate in a frozen sand sample ($t= -6^{\circ}C$, $p=0.1MPa$, $C_{sol}=0.1N$): A – kaolinite; B – montmorillonite.	160

Figure 70. Influence of the size of sand particles on the accumulation of salt ions (Na^+) and the decrease in the hydrate coefficient (K_h) in frozen hydrate-containing samples ($K_h^{\text{in}} \sim 0.4$, $W=12\%$, $t=-6^\circ\text{C}$) 4 hours after the start of interaction with frozen salt solution 0.1 N NaCl: A - fine sand-2 (0.1-0.2 mm $\sim 80\%$); B – medium sand (0.2-0.5 mm $\sim 60\%$)..... 161

Figure 71. Influence of silt content on the accumulation of salt ions in frozen hydrate-containing samples 4 hours after interaction with 0.1 N NaCl solution at $t=-6^\circ\text{C}$: A –fine sand-2 (silt $\sim 1\%$); B – fine sand-3 (silt $\sim 5\%$); C – sandy loam (silt $\sim 18\%$). 163

Figure 72. Influence of silt content on the hydrate coefficient (K_h) of frozen hydrate-containing samples ($K_h^{\text{in}} \sim 0.4-0.5$, $W=12\%$) 4 hours after interaction with 0.1 N NaCl solution at $t=-6^\circ\text{C}$: A – fine sand-2 (silt $\sim 1\%$); B – fine sand-3 (silt $\sim 5\%$); C – sandy loam (silt $\sim 18\%$). 163

Figure 73. Influence of silt content on moisture accumulation in a frozen hydrate-saturated sample ($W=12\%$) after 4 hours of interaction with a frozen salt solution of 0.1 N NaCl at -6°C : A – fine sand-2 (silt $\sim 1\%$); B – fine sand-3 (silt $\sim 5\%$); C – sandy loam (silt $\sim 18\%$)..... 164

Figure 74. Influence of fineness and amount of clay particles (kaolinite) on the average migration flow of Na^+ in frozen hydrate-saturated samples interacting with NaCl solution 0.1 N for 4 hours at $t=-6^\circ\text{C}$: A – fine sand-3; B – fine sand-2 with kaolinite (from 0 to 10%); C – medium sand; D – sandy loam..... 166

Figure 75. Proposed model of interaction between saline horizons and frozen deposits containing gas hydrates: a) initial state; b) start of salt migration from cryopegs due to ambient temperature increase; c) dissociation of metastable hydrates occurs; b) dissociation of stable hydrates occurs. 169

Figure 76. The proposed conceptual model on the interaction between frozen hydrate-containing sediments and sea water: a) preliminary state – sea regression; b) after sea transgression – start of interaction between frozen sediments and sea water; c) migration of salt into permafrost sediments (development of salinity and thawing fronts); d) propagation salt front into sediments, saturated with intrapermafrost gas hydrate followed by hydrate decomposition and active methane emission. 170

List of Tables

Table 1. Depth of permafrost and zero temperatures distribution on the north of West Siberia gas fields [8].	29
Table 2. Complications, which can be caused during drilling in the permafrost (modified after [3,49]).	33
Table 3. Registered cases of gas emanations in the Russian Arctic [236].	51
Table 4. Particle size distribution, mineral composition, and salinity of investigated sediments.	81
Table 5. Main initial characteristics of frozen hydrate-containing sand samples.	100
Table 6. Average Na^+ flux in frozen sand (fine sand-1) under the equilibrium (2.5 MPa) and non-equilibrium (0.1 MPa) pressure (4 hours of interaction with 0.2N NaCl solution at -6°C).	102
Table 7. Frozen non-hydrate saturated (sample 1) and hydrate saturated (sample 2) fine sand -2 initial characteristics.	108
Table 8. Initial characteristics of the frozen hydrate-saturated fine sand-2 samples before the contact with NaCl solution of different concentrations.	113
Table 9. Scheme of phase-transformation in frozen hydrate-saturated sediments during increase of salt solution concentration at stable conditions (experimental pressure $P_{exp} \geq P_{eq}$) and metastable conditions ($P_{exp} < P_{eq}$).	145
Table 10. Dependence of the critical concentration (C_{cr}) causing the decomposition of pore hydrate in a frozen fine sand-2 sample on the chemical composition of the contacting solution ($t=-6^\circ\text{C}$, $P=0.1\text{MPa}$, $C_{sol}=0.1\text{N}$).	152
Table 11. Properties of hydrate-saturated frozen sand with different amount of kaolinite.	154
Table 12. Properties of hydrate-saturated frozen sand with different composition of clay.	157
Table 13. Properties of hydrate-saturated frozen sand with different amount of silt.	162
Table 14. Dependence of the critical NaCl concentration on the amount of silt in frozen hydrate saturated sandy sediment at the temperature -6°C .	165

Chapter 1. Introduction

1.1. Relevance. Statement of the problem

Today, the leading area for oil and gas production remains the Arctic region. Nevertheless, the exploitation of hydrocarbon deposits here is often accompanied by many complexities, mainly related to the spread of permafrost [1–8]. Frozen sediments are a complex multi-component and multi-phase system containing natural salt solutions (cryopegs), underground ice, as well as methane hydrates, which **very vulnerable** to different local man-made impacts (mechanical, thermobaric, chemical, etc.) [1,3,9–11].

Gas hydrates (clathrates) are crystalline compounds that look like ice or **snow**, which are formed from water and a low molecular weight gas at high pressure and low temperatures [12–14]. In ~~the~~ nature, hydrates occur in **the sea** and ocean sediments (at the depths of the northern seas from 250-300 meters, and from a depth of 600 meters in the southern seas), along with permafrost areas with satisfactory thermobaric and geochemical conditions in the so-called hydrate stability zone [15–21]. ~~There are requirements for the methane hydrate existence in the frozen strata above the modern stability zone (metastable gas hydrates) due to the effect of self-preservation [15,22–29].~~ **These shallow, metastable gas hydrates** can be primarily destabilized by changes in environmental conditions (pressure decrease, temperature increase, and salt migration) [30–32].

An important characteristic of methane hydrates is their possibility to store a significant gas amount - up to 160 m³ of methane in 1 m³ of hydrate [33]. This allows to consider natural gas hydrate accumulations, on the one hand, as a promising unconventional source of hydrocarbons, and, on the other hand, as a serious geological hazard in the development of traditional oil and gas fields, which, as a rule, are located below the zone of possible hydrate formation [10,12,19,27,33–38].

Gas hydrate formations in the Arctic can be destabilized as a result of permafrost degradation (due to the rise of temperature) [32], and throughout salt migration into the frozen hydrate-containing sediment [31,39,40]. The reasons for the destabilization of gas hydrate accumulations as a result of the impact of salts can be separated into natural and technogenic. Among the natural causes, there are the migration of salt ions into frozen hydrate-saturated sediment from sea water on the Arctic shelf, and from cryopegs or saline sediment horizons in continental conditions. Technogenic causes include the flow of exposed cryopegs and saline sediment horizons from sea water as a result of drilling exploratory and production wells when various salt solutions are used in the drilling process, as well as during the burial of industrial solutions in frozen hydrate-containing sediment [41]. These negative phenomena lead to an increase in the risk of emergencies and negative consequences, leading to an increase in financial costs during the operation of deposits in the Arctic [4,5,10,42].

To date, the issues of salt migration during the interaction of salt solutions with intrapermafrost methane hydrates have been poorly studied, especially from the standpoint of the kinetics and mechanism of the process. Many studies are dedicated to experimental and mathematical research on the interaction between frozen sediments and salt solutions, without taking into account the hydrate component [43–52]; mathematical and experimental modeling of the process of hydrate inhibition in sediments by saline solutions [53–56]. And only recently there have been works on mathematical modeling of the process of dissociation of gas hydrates on the Arctic shelf as a result of the migration of salt ions into hydrate-saturated frozen sediments [57].

In this regard, setting up experimental modeling to study the mechanism and dependences of dissociation of porous hydrates in frozen sediment as a result of salt migration at various pressures (both equilibrium and below equilibrium) is of certain interest. The result of such research can help to evaluate the role of salt migration in the destabilization of intrapermafrost hydrate and methane emission in the Arctic.

1.2. Goal

Experimental study of the mechanism and regularities of the interaction of gas hydrates with salt solutions in the pore space of hydrate-saturated sediments under various thermobaric conditions.

1.3. Objectives

1. Development of an experimental method for the interaction of gas hydrate with salt solutions in the pore space of frozen sediments under pressure equal to or above atmospheric and negative temperatures;
2. Development of an experimental method for the temperature field investigation during salt transfer in frozen hydrate saturated sediments;
3. Experimental study of the mechanism and dynamics of the interaction of gas hydrates with salt solutions in the pore space of frozen sediments;
4. Experimental study of the influence of various factors (temperature, pressure, salt solution concentration and composition, and sediment composition) on the processes of hydrate dissociation during the chemical components penetration into hydrate-bearing frozen sediments;
5. Experimental investigation of the temperature change dependencies in the processes of mass transfer and dissociation of hydrates during salt migration in frozen sediments, containing methane hydrates;
6. Creation of schematic model of the interaction of gas hydrates saturated frozen strata with natural salt and technical solutions in hydrate-bearing reservoirs in permafrost.

1.4. Novelty

- A method for determining the content of salt ions (Na^+) in sand was developed based on the measurement of pore moisture activity;
- An experimental method for the interaction of frozen hydrate-bearing frozen sediment with salt solutions under pressure > 0.1 MPa has been developed;
- An experimental method for study of the temperature change in frozen hydrate-saturated sediment interacting with salt solution was developed;
- For the first time, the effect of salt migration on the dissociation of hydrates in frozen hydrate-saturated sediments was experimentally shown, and compared with the process of salt transfer in frozen non-hydrate-saturated sediment;
- Temperature variation during salt migration in frozen hydrate-bearing sediments was investigated;
- The influence of various factors (temperature, pressure, concentration and composition of salts, clay content and dispersity) on the dissociation of hydrates in the pore space of frozen sediment during their contact with salt solutions was revealed;
- The critical concentration of salts for the complete decomposition of the hydrate (C_{cr}) was determined depending on various factors;
- The change in the average salt flux in frozen hydrate-saturated sediments was calculated depending on various factors;
- Conceptual scheme-model of destabilization of intrapermafrost gas hydrates in areas of permafrost distribution in continental conditions and on the Arctic shelf is proposed.

1.5. Outline of the Thesis

Chapter 2 explores the current understanding of Arctic permafrost from the geological point of view, and possible natural and technogenic geohazards in the area of hydrocarbon production in the continental conditions and on the shelf.

Chapter 3 provides a literature review for the current state of the knowledge connected with gas and gas hydrates as a component of permafrost, and the dependences of their destabilization, with special emphasis on the salt influence investigation on hydrate dissociation.

Chapter 4 introduces existing experimental and mathematical research on hydrate-salt solution and frozen sediment-salt solution interaction.

Chapter 5 illustrates the method for artificial hydrate saturation of frozen sediments, their interaction with the salt solution under stable and metastable conditions, and temperature field investigation during the interaction process. In addition, a new method for salt concentration determination in sandy sediments is proposed. Finally, the sediments characteristic and algorithm for the calculation of the main characteristics are given.

Chapter 6 describes the experimental data on the dynamics of hydrate saturated frozen sediment interaction with salt solution in time. A comparison of salt migration in frozen hydrate-containing and frozen non-hydrate-containing frozen sediments is given. In addition, the temperature change in the process of salt transfer in hydrate saturated frozen sediments is analyzed. Based on the experimental data the description of the mechanism of phase transitions in frozen sediments, containing gas hydrates in stable and metastable form, is given.

Chapter 7 demonstrates, via experimental modelling, the influence of different factors on the interaction of hydrate-saturated frozen sediments with salt solution, such as pressure (above and below equilibrium), temperature (from -3 to -20 °C), salt solution concentration (from 0.1 to 0.4N) and composition (NaCl, CaCl₂, NaSO₄, MgCl₂, KCl). In addition the influence of clay content of different

mineralogical composition (kaolinite and montmorillonite) and dispersity on the intensity of salt migration and hydrate dissociation in frozen sediments is provided.

Chapter 8 cover concept schematic models of the process of hydrate destabilization by natural and technogenic solutions on the Arctic self and in continental conditions.

Chapter 2. Arctic permafrost and problems of hydrocarbon fields development in the Arctic

2.1. Permafrost-geological structure of the Russian Arctic region

~~As it is well known,~~ a large number of hydrocarbon deposits in Russia are concentrated in the Arctic, where more than 83% of natural gas and 12% of oil are produced [7,8,58,59]. However, the largest part of the Arctic territory of Russia is occupied by permafrost, reaching thicknesses from hundreds of meters to 1 km [60,61]. In this regard, in order to plan safe economic development, including the development of energy resources, it is first of all necessary to understand the geological structure and main parameters of frozen sediments (temperature conditions, ice content, sediment composition, frozen sediment thickness, etc.).

Basic information on the permafrost distribution patterns and the main components of permafrost is described in the works of Kudryavtsev V.A., Dostovalov B.N., Romanovsky N.N. Kondratieva K.A., Melamed V.G. (1978) and Ershov E.D. (2002). Frozen sediments refers to sediments whose ground particles are cemented by ice and are in conditions of negative temperatures for more than 1 year. There are also sediments without ice content, ~~but located at~~ a negative temperature: frosty (not containing any water at all) and cooled sediments. The peculiarity of the cooled sediments is that they are saline or saturated with cryohaline waters (cryopegs) with a

concentration of 1 to 300 g/l. At the same time, salts from cryopegs can migrate into the frozen stratum, and thus changing its thickness [60,62].

Chuvilin E.M. with co-authors in 2005 gave a detailed description of the main structural elements of frozen sediment, such as the organomineral skeleton and components that fill the sediment pore space: ice, unfrozen water and gas [63]. Each of those components is important for the investigation of the salt migration.

Ice, mainly hexagonal in structure, is the main cementing component of frozen sediment. In addition, ice is the main transport medium for chemicals in permafrost [48]. The main characteristic of the ice content in the sediment is the coefficient of ice content - the ratio of the mass (volume) of ice contained in pore space to the entire mass (volume) of the sediment. Frozen sediments are subdivided into icy ($L = 0-0.2$), medium icy ($L = 0.2-0.4$), and heavily icy ($L > 0.4$) [58,60].

Unfrozen water is a part of bound water, which has not been frozen. The content of unfrozen water decreases with reducing negative temperature. In this case, ice and unfrozen water are in constant dynamic equilibrium. Also, the amount of unfrozen moisture increases with an increase in the salinity of sediments and with a decrease in gas pressure [60,64–67].

Compared to ice and unfrozen water, the gas component of frozen sediment has been studied much worse, but some systematization of its types is given in the work [63]. So, according to the genesis, gas in permafrost is divided into biochemical, deep (catagen), and coal gas. The emitting gas might migrate to the conductive zone and then gather under the impermeable layer in permafrost creating an area of high pressure [63,68].

According to the phase state, three main classes (forms) of gas are distinguished - sorbed, including adsorbed and absorbed forms, free and dissolved [63]. The sorbed class includes mineral surface gas present in frosty sediment; a gas that exists as a gas hydrate in frozen (stable and metastable form, discussed in detail in Chapter 3) or cooled sediment (stable form); coal gas that can exist in both

frozen and cooled sediment. Dissolved gas is subdivided into free and bound water gas. Free gas - pinched and movable type [63]. Compositionally heterogeneous cryogenic strata with different **capacitive and filtration** properties can accumulate gas, which can be converted into a solid gas hydrate form, or partially accumulate in lenticular bodies and remain sealed both in the gas phase and in the form of hydrates.

In general, permafrost strata along the vertical section are divided into three layers [60]:

1. A seasonal thawing and freezing layer with thickness < 5 meters, characterized by temperature changes;
2. The strata, where the annual temperature fluctuation happens within thickness < 30 meters - with high ice content and permanent temperatures from 0 to -4... -5 °C in the bottom part of the strata (up to minus 4-5 ° C);
3. Frozen sediments with persistent sub-zero temperatures. The upper part of this strata is characterized by the lowest temperatures (up to -4...-5°C). In terms of thickness, this sequence makes up the main and most part of the permafrost section.

Frozen strata can be merging with a layer of seasonal thawing and non-merging, separated from it by a thawed layer of sediment [60-62]. Depending on the distribution of permafrost, there are: continuous (more than 90% of the area, up to 1400 m thick), discontinuous (50% -90%) and isolated patches (10% -50%, up to 10-20 m thick) , between which it is possible to draw conditional boundaries [60,69].

In terms of composition, permafrost strata can be composed of loose sediments of various genesis from Paleogene to modern **Quarter time** of formation, as well as sediments and rocks of sedimentary, intrusive, effusive and metamorphic origin from Proterozoic to Mesozoic time [8].

Permafrost is constantly evolving depending on long-term climate fluctuations or changes in annual and sediment temperatures. Its thickness can increase (aggrandize) or decrease (degrade)

[60,61,70]. In this regard, there are also modern permafrost and relict. Modern strata are developed from the surface under a layer of seasonal thawing, and relict ones occur at a certain depth exceeding this layer [8].

Previously, it was found that in Russia, the thickness of permafrost reaches an average of 600-700 m, and the temperature varies within 0 ... -2.5 °C. At the same time, the maximum depth of permafrost (1400 m) was recorded during drilling in the northwestern part of Yakutia [60].

Figure 1 shows a map of the spreading of the permafrost zone on land in the Russian Arctic, indicating the type of distribution [71].

Scale 1:12 000 000




Figure 1. Map of the permafrost distribution in the Arctic zone of the Russian Federation [71].

The continuous distribution of permafrost is characteristic of the northern regions of Russia. However, under large natural water reservoirs (rivers, lakes) and in tectonic fissured zones of increased groundwater circulation, there are areas composed of thawed sediments for the entire thickness of the permafrost or for part of it, and their number increases from north to south. Such areas of frozen sediments are called taliks (through or non-through), which can serve as zones for unloading a gas deposit [8,72,73].

Frozen sediments are widely developed not only on land, but also on the shelf of the Arctic seas [74–81]. For the first time about permafrost existence on the Arctic shelf, A.I. Fartyshev wrote in 1993, based on his own field observations and calculated data, as well as on the results of drilling on the shelf of the Beaufort Sea [82,83]. However, to date, there are only separate field data on the thickness, distribution, and features of sediments in the permafrost zone of the Arctic shelf of Russia [84,85], as well as mathematical modeling data and individual geophysical surveys.

Mathematical modeling of the permafrost zone on the Arctic shelf of Russia was carried out by various researchers L.A. Zhigarev, Ya.V. Neizvestnov, V.A. Solovyov, A.I. Fartyshev, N.N. Romanovsky, G.S. Tipenko, V.E. Tumsky, A.V. Gavrilov, D.N. Nikolsky, V.V. Malakhova, M.N. Grigoriev, P.P. Overduin, C. Ruppel, A. Vasiliev and others. Figure 2 shows some of the latest data on modeling the propagation of the zero isotherm on the Arctic shelf, which corresponds to the distribution of the permafrost zone there [80]. The depth of underwater permafrost, which was identified earlier (in 2001) on the map of the International Permafrost Association, is indicated by a blue line. In the shaded area, assumed present-day seafloor temperatures result in permafrost formation more than 50 m above the calculated depth [80].




Figure 2. Simulation of the zero isotherm propagation on the Arctic shelf [80].

A distinctive feature of the Arctic shelf is the wide distribution of the subaqueous permafrost zone of complex structure, represented by cooled saline and permafrost sediments [86].

The permafrost zone of the Arctic sea basin was formed as a result of active thermodynamic, chemical and hydrodynamic interaction of the atmosphere, hydrosphere and lithosphere. At the same time, the formation of the submarine permafrost occurred due to freezing from above at the stages of sea regressions; and thawing - mainly from below under the influence of heat flow from the bowels of the earth at the stages of transgressions [74,84,87].

As Yakushev V.S. (2014) noted, three main oil and gas provinces are located wholly or partially in the territory of the permafrost zone on the territory of Russia: Timan-Pechora, West Siberian and East Siberian (Table 1) [8].

Table 1. Depth of permafrost and zero temperatures distribution on the north of West Siberia gas fields [8].

Fild name	Depth of permafrost, m	Depth of zero temperatures, m
Bovanenkovo	120-170	260-310
Kharasaveyskoe	160-180	250
South-Soleninskoye	230-260	420
Yamburg	265-336	400-450
South-Pestsovoye	390-460	460-500
Zapolyarnoye	420-463	460-510
Urengoi	400-480	420-500

And depending on the combination of the above-described characteristics of frozen sediments, three categories of section complexity are defined: with especially difficult geocryological conditions, complex and uncomplicated. So, for example, the Nizhnepechorsko-Korotikhsky, Shapkino-Adzinsky, Yamalo-Gydansky, Tazovsky North-Siberian, Yenisei-Putoransky, Anabarsky, and Priverkhoyano-Central-Yakutsky districts are especially complex. In work [8] these areas are considered in detail, and the main characteristics of geocryological conditions are given for them.

Thus, the major part of energy resources of Russia are located in and under the permafrost. In connection with such a wide distribution of permafrost, significant material has been accumulated in Russia on the properties of frozen sediment, the main parameters, patterns of distribution, and complications associated with the economic development of territories occupied by permafrost. The largest oil and gas provinces are located in the region of the highest values of the geocryological risk index: West Siberian, East Siberian and Arctic shelf [88]. Let us consider the complications arising from the economic development of the Arctic on land and on the shelf.

2.2. Peculiarities of hydrocarbon deposits development in areas of permafrost distribution

A number of authors note that at almost all stages of the "life" of hydrocarbon fields in the areas of permafrost, numerous complications can occur associated with negative technogenic effects of various nature (thermal, mechanical, chemical, etc.) on the components of permafrost, primarily such

as natural salt solutions (cryopegs), underground ice, gas hydrates [1–3,6,89,90]. In most cases, such complications are little controlled due to inertia, and therefore dangerous. But so far in Russia there is not a single state program for the comprehensive study of permafrost.

Drilling a well is the first and main step necessary to start developing a hydrocarbon field [3]. At the same time, the process of drilling various wells in the permafrost distribution zone differs significantly from analogues located in areas with a mild climate and positive sediment temperatures. The specificity is due to the complex interaction and influence of mining and geological, mining and technical, permafrost and climatic factors [58,91].

As mentioned earlier, the main cementing component of frozen sediment is ice, therefore, the complicated conditions for the construction of wells in permafrost are based mainly on the temperature factor, which determines, on the one hand, the efficiency of the process of destruction and transportation of frozen sediment to the surface, and on the other hand, the stability of the future wells. And above all, it should be mentioned the destruction and melting of ice occurs due to its thermal, erosive and physico-chemical interaction with the drill bit and the circulating flushing fluid, even if it is pre-cooled [92,93].

Drilling fluid is the first process fluid that interacts with newly opened frozen sediment and, accordingly, has a primary effect on permafrost [92–95]. At the moment, the issue of choosing the appropriate composition of the drilling agent, drilling mode to prevent the degradation of permafrost and secondary ice formation has been worked out in sufficient detail. **In the main document regulating well drilling [96] the main recommendations on this issue are spelled out.** It is recommended to use high-viscosity polymer-clay and biopolymer solutions with controlled solids content, bottomhole purge with air or foams. However, most works devoted to drilling fluids do not take into account the hydrate component of frozen sediment, and even more so, this is not mentioned in regulatory documents.

At the same time, it is allowed to use aqueous salt solutions (NaCl, KCl, CaCl₂, MgCl₂) as an additive to drilling fluids, and in some cases as a full-fledged drilling fluid when drilling in permafrost. [95]. Previously, it was found that saline solutions cause thawing of permafrost, as was shown experimentally ~~in the works~~ [49,97], and therefore potentially can also cause destabilization of gas hydrate formations.

The temperature effect of drilling fluid on frozen sediment is difficult to describe. On the one hand, a permafrost system initially in equilibrium is exposed to a drilling fluid with a temperature elevated relative to the formation (this can happen if the solution was not sufficiently cooled initially, or was heated as a result of the mechanical rotation of the drill bit, or the temperature of the overlying warm sediments) [93]. In this case, the equilibrium will be shifted, a temperature gradient will arise, and the pore ice will begin to melt ~~under the influence of a cleaning agent~~, turning into liquid water. As a result, the sediment will go into a softened state. However, the thermal effect can be partially absorbed by the endothermic process of the ice-liquid water phase transition and, accordingly, somewhat smoothed out. On the other hand, if the treatment agent is at a lower temperature than the host sediment, then the opposite effect may occur. Unfrozen moisture under the influence of a decrease in temperature will turn into ice with the release of heat. During the transformation of unfrozen water into ice, the salts contained in it can also be squeezed out increasing salt concentration in the remaining pore moisture, therefore increasing its freezing temperature [1,93,98]. A decrease in the temperature of unfrozen water leads to a change in the physical and mechanical properties of sediments: their viscosity and plasticity increase, and a tendency to creep appears, up to a complete loss of sediment cohesion [99].

Thus, field experience shows that during the well drilling in the area of permafrost distribution on land, phase transitions are observed in frozen sediments with a change in the strength characteristics of the rock as a result of both temperature and chemical effects of technogenic nature

[92]. Opening of gas-saturated lenses, opening of cryopegs and saline frozen horizons can also occur, followed by the flow of salts down the section [100,101]. As a result, a number of negative consequences can occur, one of which is hydrate dissociation, accompanied by powerful methane emissions, subsidence of sediment masses around wells, the formation of a deformation field, absorption of drilling fluid, cavern formation, griffon formation, and many others [4,5,10,42,102].

Another important issue is the exploration of deposits on the Arctic shelf. However, in this area, the issues of possible complications have been studied less. Nevertheless, it can be noted that all of the above-mentioned complications typical of land permafrost are possible. In addition to them, salt ions migrate from sea water to sediments on the Arctic shelf, causing degradation of the permafrost zone [74,103] and relic methane hydrate dissociation [31].

The generalization on the complications that arise when drilling wells in the areas of permafrost is carried out on the basis of several publications and is given in Table 2 [1,3,104]. It should be noted that the types of impact of the drilling process on permafrost and its consequences are divided conditionally. Consequences of this or that influence are appearing in nature in a complex.

Table 2. Complications, which can be caused during drilling in the permafrost (modified after [3,49]).

Type of drilling influence	Typical processes	Consequences	Associated phenomena
Thermal	Temperature change, thawing	Change in sediments temperature, sediment thawing (taliks formation), column subduction	Axial deformations of the metal lining, the cause of which is the loss of longitudinal stability of the columns due to stresses arising from the consolidation of frozen sediment thawed in the borehole zone
Mechanical	Visco-plastic deformation; ground discontinuity	Deformation due to porosity change, collapse of the sediments, funnels formation in subsidence sediments	Bending of the above-ground well structure. Threat of deformation fracture of casing pipes and open flowing; inability to maintain the wellhead and perform repair work

Chemical	Dissolution, substitution, redox	Transformation of mineral and chemical composition (with formation of the new one), cavern formation	Collapse of the wellbore walls, formation of cavity widening in the wellbore, mud contamination, sticking, clogging of the wellbore with particles of thawed sediment, decrease in ROP, complications during well cementing, drilling fluid loss
Physical and chemical	Texture transformation, mass transfer, secondary ice formation	Transformation of structure up to sediment damage, change in chemical composition and moisture content of sediment	Loss of cohesion of permafrost sediments with a well walls and poor-quality cementing ring due to repacking of grains, formation of ice plugs in borehole's channel, deformation of the drill string, freezing of the drill pipe

These negative phenomena lead to an increase in the risk of emergencies and negative consequences, leading to an increase in financial costs during the operation of deposits in the Arctic [1,105,106].

In addition to all the above complications of the development and production of hydrocarbon deposits, the environmental component of the potential permafrost thawing and hydrate dissociation is also relevant, which is especially important to take into account for the very vulnerable environmental conditions of the Arctic. As noted above, in addition to changes in the bearing capacity of sediment and the emergence of a number of associated hazardous geocryological processes, there is possibility of gas pockets opening and gas hydrate dissociation. Thus, a number of authors note numerous gas manifestations when drilling wells in areas where the permafrost zone is distributed [36,42,99,107–112]. Some generalization of the recorded cases of gas shows is given in Table 3.

Thus, active methane emission can occur, the transformation of the greenhouse effect, a possible change in climatic conditions, the formation of ozone holes and the so-called "methane catastrophe" [113–118].

Summarizing all of the above, ensuring well stability is one of the most important problems, especially in the permafrost area [36,98,119]. At the same time, a vast area of the main oil and gas provinces in the Arctic is located in an area of very complex and complex geocryological conditions. The existing domestic and foreign experience in the construction of wells in permafrost and their

subsequent operation have shown that insufficient knowledge of permafrost conditions, unreasonable choice of technical, technological and design solutions has a causal relationship with the occurrence of accidents and complications. Such serious problem as wellbore instability can be caused by fluid flow, heat transfer, pressure redistribution, and also hydrate decomposition. The quantitative change in the natural temperature field of the surrounding sediment depends mainly on the temperature of the cleaning agent, the sediment thermal characteristics, the radius of the well, the drilling method, and also on the amount of heat released due to the mechanical destruction (cutting) of sediments.

Chapter 3. Conditions for gas hydrate existence and formation in permafrost

Scientists from various fields of science have been studying gas hydrates for more than 200 years, since the accidental discovery of sulfur oxide (SO₂) hydrate by chemist Joseph Priestley in 1777-1778, and the first mention of the term "hydrate" by Davy in 1811 [15,120]. At the moment, the main issues of consideration of gas hydrates can be divided into two large groups: 1) gas hydrates as one of the promising non-traditional sources of hydrocarbons, the energy capacity of which is comparable to the potential of traditional energy resources [12,121–128]; 2) gas hydrates as a factor complicating the development of hydrocarbon deposits and transforming climatic conditions [10,11,107,129–132].

3.1. Characteristics of gas hydrate as a substance

At present, the structure and general characteristics of gas hydrate as a chemical compound are known in sufficient detail thanks to the work of a number of scientists, such as Makogon, Trofimuk, Cherskiy, Vasiliev, Chuvilin, Istomin, Yakushev, Sloan, Collett, Kvenvolden, Moridis, Uchida and many others. The clathrate structure of gas hydrate was first suggested by Nikitin in 1935, and a few

years later this assumption was confirmed experimentally [133]. A clathrate (from the Latin *clathratus* - “to place in a cage”), or an inclusion compound, is a compound of the “host-guest” structure. In the case of hydrates, the clathrate structure is described as follows. Water molecules form a polyhedral structure with cavities that represent the “host” structure. These cavities are filled with gas molecules (“guests”) (Figure 3). There is no chemical bond between gas and water molecules, only a weak van der Waals interaction [14,33,115,134–138].




Figure 3. Structure of gas hydrates [139].

Gas hydrates have a **fluctuating** composition (i.e., non-stoichiometric chemical compound). The following chemical formula defines hydrates composition: $M \times nH_2O$ (M – gas, forming hydrate, n - the amount of water molecules per one gas molecule, also named the hydrate number [12,140]).

Typically, the hydrate number (n) value determined accordingly to the temperature, pressure, and gas composition, and located in the range from 5.75 to 17. For instance, the hydrate number equals 5.9 in the case of methane hydrates. The hydrate number is of particular importance for all calculations related to the study of gas hydrates [115,141–143].

Figure 4 shows different hydrate structures: (a) cubic structure I, (b) cubic structure II, (c) hexagonal structure H [144].




Figure 4. Different crystalline arrangement of hydrates: a – structure I, b – structure II, c – structure H and d – host cages of water. Dense line is the hydrate unit-cell [145].

Structure I is most widespread in nature and it is presented by 2 small dodecahedral voids, 6 large tetradecahedral voids formed by 46 water molecules. Such structure is able to include gases of the molecular size less than 5.2 Å, particularly CH₄, CO₂, C₂H₆ and some other gases [126,146].

Structure II is able to include gas molecules of the size 5.9 - 6.9 angstroms (for example C₃H₈ or C₄H₁₀) and consists of 8 large hexadecahedral and 16 small dodecahedral voids composed of 136 H₂O molecules [120].

Less common structure is H structure, which is formed only by gas mixtures (like CH₃C₆H₁₁ mixed with H₂S) [15].

Such natural gases as CO₂, C₂H₆, C₃H₈, C₄H₈, O₂, N₂, H₂S, and more often CH₄, are able to form hydrates both separately and in mixtures. Some other gases are able to go into a hydrate form only at the very high pressures (such gases as H₂, He, Ne) or unable to make a hydrate at all (for example C₅H₁₂ et al.) [147]. The composition of natural gas hydrates may include hydrocarbons (methane, ethane, propane, butane), as well as gases such as CO₂ and SO₂. As shown by the data of chemical analysis of gas from intrapermafrost gas shows, the most common gas that forms natural gas hydrates is methane. However, pure methane is extremely rare. Almost always, nitrogen, carbon dioxide,

hydrocarbon homologues of methane, hydrogen sulfide are present in small amounts in the composition of the gas [12,33,120].

In appearance, gas hydrates resemble loose ice or snow (Figure 5), which was one of the reasons for the relatively late start of the study of gas hydrates.



Figure 5. Gas hydrate burning (photo by Dinara Davletshina).

~~Petroleum specialists, scientists and others have definitely encountered this substance for a long time, but could not explain the cause of many phenomena, such as, for example, the sudden emission of methane, the release of a sediment mass, the formation of plugs that resemble ice in a well, etc. [2,148].~~ Active exploration of gas hydrates only began in the 1940s and 1960s with the work of Makogon Yu. F., Strizhov I.N., Trofimuk A. A., Mokhnatkin M.P., Chersky N. V., Vasiliev V. G., Trebin F. A., who suggested the hydrate existence in nature in the permafrost area and proved it in the laboratory conditions [140,149].

The external similarity of gas hydrate and ice, due to the framework of water molecules with van der Waals bonds, determines the closeness of the values of their physical properties (density, heat capacity, acoustic properties, Poisson's ratio) [15,140]. At the same time, a number of properties are noted for which there are significant differences: thermal (ice thermal conductivity (λ) equals 2.2 W / m K, and methane hydrate $\lambda=0.5$ W / m K) and dielectric (dielectric constant of ice - 94, methane hydrate - 58). Also, during hydrate formation, the specific volume can increase by 26%, while when water freezes, the volume increases by 9% [15,138].

Under normal conditions, one volume of methane hydrate, as mentioned above, contains 160 volumes of gas, which emphasizes its potential use as an energy source, on the one hand, and the danger of dissociation of gas hydrate formations in the sediment, on the other. [33].

Figure 6 shows the phase diagram of the existing conditions of methane hydrate, one of the most common types of gas hydrates in nature. When the thermobaric conditions change, gas hydrates dissociate into water and gas [150].




Figure 6. Stability of CH_4 hydrate curve (build in HydraFlash): GH – gas hydrate; liq – liquid water.

There are many works devoted to the study of the p-T shift of the conditions for the hydrate existence and dissociation and the influence of various factors on this process (inhibition by various

substances, type of gas hydrate, etc.) [33,135,151]. Works on the study of the dissociation of gas hydrates are mainly devoted to their inhibition in the development of hydrocarbon deposits (to prevent hydrate formation) [152–156], or for the purpose of methane production from hydrate-bearing sediment [55,138,157–161]. Within the framework of this work, it is interesting to review the hydrate dissociation at temperatures below 0 °C.

As written in a number of works [127,136,150,162,163] the dissociation of methane hydrate can occur as a result of a change in thermobaric conditions, when the pressure drops below the stable one or the temperature rises and the conditions go out of equilibrium. Also, dissociation can occur when the gas hydrate stability curve shifts as a result of various factors (mainly due to the presence of inhibitors in the system) [14,164]. Figure 7 schematically shows the shift in hydrate stability as a function of various factors.





Figure 7. Shift of gas hydrate equilibrium by different methods of methane extraction [144].

Hydrate dissociation is an endothermic process (i.e., it occurs with the absorption of heat) [14,165,166], and the following chemical equation describes that process [136]:



where n - hydrate number, and ΔH - hydrate formation (or dissociation) enthalpy (equals around 520-540 kJ/mol by [14,165]).

Most authors distinguish two stages in the process of hydrate decomposition due to pressure reduction at negative temperatures: active hydrate dissociation and then slowing of the hydrate dissociation ~~up to almost complete fading~~. Such a slowdown in the hydrate dissociation can be explained by the self-preservation effect of gas hydrate (Figure 8) [23,25,27]. When the pressure drops below equilibrium and the gas hydrate begins to dissociate, the water formed as a result of the decomposition of the hydrate at a negative temperature turns into a fairly thick and strong ice shell around the crystals or aggregates of gas hydrates, under which there is an excess internal gas pressure that prevents further dissociation of the hydrate [22,167]. Such an ice cover ensures the long-term preservation of gas hydrates even at atmospheric pressure without significant changes in their gas content under conditions of negative temperatures [135].




Figure 8. Scheme of bulk hydrate self-preservation at the negative temperature (picture obtained and analyzed through extensive cryo-FE-SEM imaging [168]).

Works devoted to the study of the gas hydrate dissociation as a result of exposure to various chemicals were carried out from the standpoint of hydrate inhibition for the extraction of methane (for

example [169]) or to prevent hydrate formation in wells and pipelines (for example [154,155,170]). A hydrate inhibitor is a chemical compound, which shift the hydrate phase equilibrium to the side of lower temperatures and higher pressures. Consequently, the hydrate formation process is slowed down in the reservoir [125,136].

There are organic and inorganic (brines, sea water) inhibitors of gas hydrates (Figure 9) [15,145,154].




Figure 9. Different hydrate inhibitors and scheme of their work principle [145].

Within the framework of this work, of particular interest are studies devoted to the effect of inorganic inhibitors, particularly salts, on the dissociation of gas hydrate. There are several works devoted to the investigation of hydrate dissociation, mainly from the standpoint of their potential development by the method of inhibition with saline solutions.

First of all, there are a number of works on the thermodynamic study of the effect of salts on the shift in the conditions for the gas hydrate existence, which is directly related to their potential destabilization [12,14,15,171–191]. Part of the works devoted to the study of salt water influence on

the shift of hydrate stability [192]. It was found, both experimentally and as a result of thermodynamic calculations, that an increase in the concentration of chloride salts shifts the equilibrium conditions towards higher pressures and lower temperatures [136,140,184].

In addition [53,54,56,161,169,172,180,181,183,193–198] conducted a study on the effect of salt solutions of electrolytes of various concentrations and temperatures on the hydrate component. The rate of hydrate dissociation has been shown to depend on brine temperature, brine concentration, hydrate/brine interaction surface area, and pressure. Because of both experimental and mathematical modeling, it was found that the rise in temperature, the salt concentration of the solution, and reduction in pressure contribute to the intensification of the hydrate dissociation process. An increase in the contact area also contributes to a more intensive decomposition of methane hydrate. Zheng with co-authors also noted a decrease in temperature in the course of experimental modeling of the NaCl influence on the hydrate decomposition, and the higher the concentration of the injected brine, the greater its decrease. However, the paper does not provide an explanation for this temperature effect [180].

In works [164,172,197,199–201] the mechanism of gas hydrate inhibition under the influence of NaCl salts at positive temperatures is described. The author believes that the mechanism by which the salt solution stimulates the dissociation of hydrates is caused by the interaction between water molecules involved in the structure of the hydrate complex and NaCl ions in an aqueous solution. The salt is ionized in an aqueous solution and interacts with hydrogen-water molecules on a hydrate cell, and its Coulomb bond is much stronger than the hydrogen bond or the Van der Waals force. Consequently, a stronger bond between salt and water ions will lead to the destruction of the initially stable hydrate structure and methane releases (Figure 10).




Figure 10. Hydrate decomposition in water by NaCl ions migration in time by mathematical modeling (red – methane, green – Na⁺, yellow – Cl⁻) [201].

Scientists [164,172,175,176,179,184,196,200,202–204] conducted an experimental study of the influence of the salts composition of (NaCl, CaCl₂, MgCl₂, KCl) on the processes of hydrate formation and dissociation in water free volume. An increase in the intensity of dissociation of gas hydrates in the series without saline solution - KCl solution - CaCl₂ solution - NaCl solution - MgCl₂ solution was revealed, however, with a change in concentration, the ratio may change. Additionally, Saw with co-authors (2014) determined the enthalpy of hydrate dissociation depending on the salt composition. So, more heat will be absorbed in the presence of sodium chloride salts in the system, and less heat in the case of calcium chloride salts, which means that secondary ice formation is more likely to be more intense in the system with NaCl [164].

However, it should be noted that the issues of inorganic salts influence on the kinetics of hydrate dissociation are less studied [205].

The conditions for the hydrate formation and dissociation in the sediments pore space is a more complex process. A number of scientists believe that a porous medium contributes to a more intensive process of hydrate formation compared to a free one because more nucleation sites are provided, the energy of the nucleation process is lower, the gas-liquid contact increases and the barrier to hydrate

nucleation decreases [185,206–210]. But at the same time, there is also an opinion that the pore space can act as an inhibitor of hydrate formation (by reducing the activity of water and improving the dissolution of methane), and at the moment this view is the most common [211–214]. Wu with co-authors suggests that both formation and dissociation of hydrate more actively occurs in a porous medium than in the free volume [215].

Also, a number of authors describe the effect of the pore space: there is an optimal size of the pore space at which hydrate accumulation occurs faster and the hydrate is subsequently more stable. [210,216–218]. In addition, there is the concept of non-clathrate water - moisture that does not go into a hydrated state and remains in a liquid state in equilibrium with the rest of the system components (ice, hydrate) [64,219]. Four models for hydrate formation is defined in the sediments pore space: supporting, cementing, floating, and coating [12].

Let us consider the conditions of hydrate formation and existence in nature in the permafrost zone.

3.2. Conditions for the existence of permafrost associated gas hydrate and possible causes of their destabilization

Methane hydrates are widely distributed throughout the world. To date, more than 200 gas hydrate deposits have been registered. The volume of methane contained in the hydrated form in sediment is estimated over a wide range: $3.1 \cdot 10^{13}$ – $3.4 \cdot 10^{16}$ on land and $2 \cdot 10^{14}$ – $7.6 \cdot 10^{18}$ m³ in marine conditions [2,15,33,138,220,221].

In nature, gas hydrates exist in the layer of the lithosphere and hydrosphere, where there is suitable thermobaric and geochemical regime, in so-called hydrate stability zone (HSZ) [33,121,222]. Such conditions for methane hydrate exist in the permafrost zone (on land) (Figure 11B) and on the

sea shelf from a depth of 600 m (in the southern seas) and 300 m (in the northern seas) (Figure 11A) [15,33,121,150].




Figure 11. Existence of methane hydrate in natural permafrost conditions (modified from [17,223]).

Estimates of the depth of the upper and lower HSZ boundaries for onshore permafrost conditions have been obtained from field data and through mathematical modeling and vary widely, mainly depending on the thickness of the permafrost. The upper boundary of HSZ for methane hydrates is located at the depth of about 150 m in continental permafrost regions where surface temperatures are maintained at about 0 °C [224]. The lower limit of the existence of methane hydrates can reach a depth of 1000 meters (for example, on the Siberian platform) (Figure 11B).

At the same time, the thickness and depth of HSZ on the Arctic shelf is a much more complex issue due to insufficient field data. Romanovsky with co-authors obtained some data on the thickness, history of formation and evolution of the permafrost zone and HSZ of the Arctic shelf based on a

retrospective analysis, and Malakhova and co-authors made a publication based on mathematical modeling [57,225]. The HSZ upper boundary is traced at around 100-145 m beneath the sea bottom, deepening towards the coast, which is determined by an additional baric factor due to the weight of the water layer on the outer shelf. The thickness of HSZ increases with decreasing sea depth, which is explained by lower temperatures of sediments in the shallow part of the shelf and the greater permafrost thickness. The current value of the thickness of the stability zone, according to the simulation results, is approximately 700 m for the outer shelf and 1130 m for the inner shelf [84].

However, gas hydrate in the permafrost zone can also exist above the gas hydrate stability zone in the region of relatively low pressures (below equilibrium) at temperatures below 0 °C for a long time, in the so-called hydrate metastability zone (HMZ). It represents the permafrost interval above the HSZ, where relic hydrates exist due to the self-preservation effect [23]. The hydrate metastability zone, in contrast to the zones of stability, is not a thermodynamic, but a geological zone. Relic gas hydrates is not formed in MHZ, but exist in the conserved form under the condition of ice coexistence for a long geological time [23].

There are several scenarios for the history of the formation of metastable (relict) hydrate in the permafrost zone [226]:

- 1) With rapid freezing of closed taliks saturated with methane, due to the creation of excess pressure;
- 2) During periods of glaciation due to the pressure created by ice on the underlying gas-saturated sediments;
- 3) Due to the increased pressure exerted by the water column on gas-saturated sediments during transgression.

Due to the creation of increased pressure in gas-saturated sediment, the gas hydrate is formed in the gas hydrate stability zone. However, later, during the evolution of the permafrost zone, as a

result of changes in thermobaric conditions, part of the relict gas hydrate formations passed into the area of pressures below equilibrium, while maintaining negative temperatures of the host sediment [84].

V.A. Istomin and V. Yakushev (1992) defined three main type of natural hydrate formation depending on their location in permafrost: supraperafrost, intrapermafrost and subpermafrost [15,30,227]. Subpermafrost gas hydrates are currently the most studied due to the many cases of real registered natural core extraction of gas hydrate saturated sediments (Mallik, Canada [162]; Prudhoe Bay and mount Elbert, Alaska, USA [13,126,228]; Qilianshan, Tibet, China [35,229–232] etc.).

Also, gas hydrate formations can be distributed differently in sediment under natural conditions (Figure 12) [12].




Figure 12. Hydrate distribution in sediments [12].

As mentioned above, the specificity of the hydrocarbon fields development is that many large hydrocarbon fields (Bovanenkovskoye, Yamburgskoye, Zapolyarnoye, Medvezhye, Urengoyskoye, etc.) are located in the permafrost zone. Both exploration and production wells can pass through the gas hydrate stability interval. Consequently, during the development of deposits, destabilization of gas hydrate formations may occur [10,107,233].

Currently, there are only a few areas of permafrost where the natural hydrates existence has been proven by direct data (extraction of hydrate-saturated deposits as a result of drilling). However,

there are many prospects where the presence of hydrate-bearing formations is suspected due to active methane emission, well logging and stability conditions [17].

Purposeful study of gas manifestations in the Arctic regions of Russia began relatively recently (about 60 years ago), which was associated with the discovery of the largest hydrocarbons fields in the North of Western Siberia and their development: Messoyakha (1967), Urengoyskoye (1966), Yamburgskoye (1969), Bovanenkovskoye (1971), etc [8,33,127,234]. When arranging the field infrastructure and laying transport routes, prospectors and builders began to encounter numerous gas emissions from the upper horizons of the permafrost zone, which often accompanied drilling operations. At the same time, both spontaneous short-term gas emissions and long-term gas emissions from engineering and geological wells drilled in monolithic permafrost occurred [42,111,235,236]. At the same time, they began to pay attention to natural gas emissions from the bottom of the Arctic seas and lakes [116,118,237,238].

An analysis of gas manifestations described by researchers shows that they can be conditionally divided according to the place of manifestation - on land and on the Arctic shelf, and according to the conditions of occurrence - into natural (natural) and man-made, associated with anthropogenic impact on the permafrost. Natural gas vents on land are often associated with various natural sources of water (lakes, rivers) and wetlands [236]. In the frame of this work, the gas that is generated in the seasonally thawed layer is not considered. Recently, another natural source of methane emission is gas emission craters, recorded in recent decades during the development of the Arctic zone of Western Siberia [72,106,236,239].

The collection and analysis of available literature data on gas shows in the Russian Arctic made it possible to compile a summary table (Table 3), reflecting the detection area, characteristic features and a possible source of gas, as well as a schematic map of gas shows in the Arctic region of Russia (Figure 13) [236].

Table 3. Registered cases of gas emanations in the Russian Arctic [236].

		No in map	Location	Signatures	Possible gas sources	Reference	
Onshore	Natural	Lakes	1	Yamal Peninsula, Lake Neyto	Gas plumes on water surface, mud flows, holes in ice	Deep gas from Neyto field	[240]
			2	Yamal Peninsula, Crater Lake	Craters and parapets on lake bottom, mud flows	Possibly, deep gas-water fluids	[238,240]
			3	Yamal Peninsula, group of lakes south of Bovanenkovo field	Blue water, holes in ice, mud flows	Active emission, possibly, deep gas	[241]
			4	Gydan Peninsula, a lake on the left bank of Yuribey River	A crater encircled by a parapet	Prolonged ascending flow, possibly, deep gas	[241]
			5	Yamal Peninsula, 54 km northeast of Arctic field, Otkrytie Lake	Large craters (up to 40 m in diameter on lake bottom)	Deep gas, possibly, from Cenomanian gas reservoirs	[110]
		6	Craters	Yamal Peninsula, 30 km south of Bovanenkovo field	A large crater (up to 40 m in diameter and ~70 m deep), encircled with a well-pronounced parapet; ground dispersed to a distance of 120 m	Possibly, intrapermafrost gas	[242–247]
		7		Gydan Peninsula, 100 km northwest of Antipayuta Village	A crater, 10-13 m in diameter and ~15 m deep; no parapet	Possibly, intrapermafrost gas	[248]
		8		Yamal Peninsula, floodplain of Erkuta-Yakha River	A crater, 10-12 m in diameter and ~20 m deep, with a preserved fragment of a 2-3 m high mound and ground dispersed to a distance of 100 m	Biogenic gas in a talik and deep gas	[249]
		9		Yamal Peninsula, 33 km northwest of Seyakha, Myudriyakha River	Fire gas explosion that produced a 50 m deep 50×70 m crater in a river, with dispersed blocks of permafrost and ice-rich soil, up to 150 m ³	Possibly, deep gas	[250,251]
		10		Gydan Peninsula, north of Deryabino field, bank of Mongoche River	Fire gas explosion that produced a 20 m deep crater, with dispersed large fragments of sediment	Possibly, deep gas	[85,241]
	Man-caused	Drilling in permafrost	11	North Yakutia, Anabar—Khatanga interfluve	Gas shows at depths 70-120 m	Intrapermafrost gas	[252]
			12	Yamal Peninsula, Yuribey River	Gas shows at depths 10-50 m	Intrapermafrost gas	[111,253]
			13	Taz Peninsula, Zapolyarny field	Gas shows at depths 50-120 m	Intrapermafrost gas, possibly relict gas hydrates	[33]
			14	Taz Peninsula, Yamburg field	Gas shows at depths 45-55 m	Intrapermafrost gas, possibly relict gas hydrates	[33,254]

Offshore		15	Ascending flows and seeps	Yamal Peninsula, Kharasavey field	Gas shows at depths 10-210 m	Intrapermafrost gas, possibly relict gas hydrates	[255,256]
		16		Yamal Peninsula, South-Tambey field	Gas shows at depths 40-60 m	Intrapermafrost gas, possibly relict gas hydrates	[257]
		17		Gydan Peninsula, Pelyatka field	Gas shows at depths 20-30 m	Intrapermafrost gas	[33]
		18		Gydan Peninsula, Salman (Utrenneye) field	Gas shows at depths 50-150 m	Intrapermafrost gas, possibly relict gas hydrates	[23]
		19		Yamal Peninsula, Bovanenkovo field	Gas shows at depths 20-130 m	Intrapermafrost gas, possibly relict gas hydrates	[68,256,258,259]
	Natural	20	Laptev Sea, Yana Delta	Ascending flows and seeps	Ebullition	Deep gas	[33,260]
		21	East Siberian Sea, Bennett Island		Gas plumes, up to 1000 km long	Possibly, deep gas	[260,261]
		22	Laptev Sea, between Semyonovsky Island and Lena Delta		Gas seeps	Possibly, deep gas	[262]
		23	Laptev Sea, Ivashkina Lagoon		About twenty gas seeps; high concentration of methane in air	Deep gas	[263–265]
		24	Laptev Sea, Kotelny Island		Gas seeps	Possibly, deep gas	[266]
		25	Laptev Sea shelf, New Siberian Islands		A cluster of gas seeps at 50-90 m sea depths	Possibly, deep gas	[267]

Man made	26	Laptev Sea (between 76.5–77.5° N; 121–132° E)	More than 700 gas seeps, up to 1.3 km in diameter	Possibly, deep gas	[265,268]
	27	Chukchi Sea, Herald Canyon and Wrangel Island	About 90 gas seeps at 50-95 m sea depths	Deep gas	[269]
	28	Kara Sea, Universitetskaya structure	Gas seeps from a depth of 80 m and pingo-like features	Deep gas	[270,271]
	29	Pechora Sea, Kara Gates	10 m high gas fountain from a borehole at subbottom depth 50 m	Deep gas	[111,254]
	30	Pechora Sea, Vaygach Island	10 m high gas fountain from a borehole at subbottom depth 50 m, with a ebullition zone up to 200 m in diameter on the sea surface	Gas from subsea permafrost	[272,273]
	31	Kara Sea, Baydaratskaya Bay	Gas show from subbottom depth 10-50 m	Possibly, deep gas	[251]
	32	Kara Sea, Leningrad field	Gas show from a 200 m deep borehole	Possibly, deep gas	[251]
	33	Laptev Sea, Lena Delta	Gas show from a borehole at 9 m depth	Gas from permafrost	[274]
	34	Laptev Sea, Buor-Khaya Gulf	Gas show from a 13-16 m deep borehole	Gas from permafrost	[274]
	35	Laptev Sea, Mamontov Klyk Cape	Gas show from boreholes up to 80 m deep	Gas from permafrost	[275]




Figure 13. Schematic map of registered gas emanations in the Russian Arctic [236].

Connection of part of the gas manifestation with hydrate formation in permafrost was shown by N.N. Romanovsky et al. (2006), D.Yu. Nikolsky et al. (2012), D. Delisle (2000), N.E. Shakhova et al. (2010), V.I. Bogoyavlensky, O.S. Sizov, I.V. Bogoyavlensky, R.A. Nikonov (2016), Bondur, T.V. Kuznetsova, V.E. Vorobyov, V.V. Zamshin (2014) and other scientists [224,240,250,276–282].

However, there is an alternative point of view. So, for example, O.A. Anisimov et al. (2012, 2014) conducted research on the Arctic shelf based on mathematical modeling and concluded that deep-seated methane that is not associated with hydrates is biogenic [114,283]. C.D.Ruppel and J.D.Kessler in their studies emphasize as well on still unproven connection of gas manifestations and gas hydrate destabilization, that methane out of hydrates at all reaches atmosphere [284].

Hydrate-saturated sediment in the areas of permafrost distribution can be affected by various technogenic and natural solutions, such as highly mineralized solutions left from the development of deposits and / or buried in permafrost massifs, cryopegs discovered during well drilling, wastewater of various origins (industrial, municipal, etc.), drilling fluids (including salt-containing ones) used in the process of drilling wells, as well as the penetration of sea water into hydrate-saturated sediment on the Arctic shelf [60,72,74,100,103,285]. However, at the moment there is a lack of understanding of the patterns of occurrence of the whole complex of complications during well drilling associated with the dissociation of gas hydrate formations [18].

There are a number of works devoted to the experimental study of the interaction of drilling fluids (including salt-containing ones) [3,93,98] and salt solution [43,45,46,97,286] with frozen sediment, excluding the hydrate component; experimental simulation of drilling fluid interaction [94,108,287–289] and salt solution [197,205,290–292] with hydrate-containing non-frozen sediments; experimental and mathematical modeling of the effect of salts on methane hydrate in free volume at negative temperatures [208]; mathematical modeling of the interaction of salt solutions with thawed and frozen sediment [49]; and only a few works are devoted to mathematical modeling of the

interaction of frozen hydrate-saturated sediment with drilling fluids that do not contain salt [129,288,293,294].

Prasad with co-authors (2019) made an investigation of the salt influence on the hydrate stability and their self-preservation. The authors consider salt solutions as a thermodynamic inhibitor of hydrate formations. In this regard, the penetration of salts into the thickness of frozen hydrate-saturated sediment can shift the HSZ deeper to the reservoir, thus reducing the HSZ parameters (length and thickness) [208]. The authors note that in the MHZ permafrost zone, the hydrate exists until the concentration of the pore solution reaches the melting point of ice around the preserved hydrate.

According to the results of experimental studies [208] 3 stages of interaction of salts with sediment saturated with metastable gas hydrates are distinguished: replacement of the initial fresh pore solution of hydrate-containing sand (no dissociation is observed at this stage), when salinity reaches a certain critical value of ice melting, the hydrate begins to dissociate; the third stage, when there is no hydrate in the system and only heat is absorbed from the environment to return the initial temperature state of the system. By analyzing the change in temperature, the authors traced the dissociation of pore gas hydrate by a sharp decrease in temperature. So, the higher the salt concentration, the more significant was the decrease in temperature, that is, the more intensively the pore hydrate dissociated. It should also be noted that the authors consider concentration diffusion as the main mechanism of salt migration [208].

S. Sun with co-authors (2011) [290] considers the p-T shift of the conditions for the hydrate existence in the pore space due to NaCl ions at positive temperatures based on experimental modeling of the interaction of sea water with hydrate-bearing sediment. The authors note that conditions are shifting towards higher pressures and lower temperatures, and the higher the salt concentration, the greater this shift. The authors attribute the intense dissociation of gas hydrates in the pore space due to the migration of salts to a decrease in the activity of pore moisture during the hydration of salt

ions, and also to the fact that weak van der Waals bonds are less preferable than electrostatic ones - therefore, salt ions will also be easier to "pull" water molecules from the crystal lattice of the hydrate, thereby destroying its lattice and facilitating the release of methane [208].

Perlova E.V. et al. (2017) suggest that the destabilization of hydrate-bearing horizons can occur as a result of penetration of drilling fluid, as well as **Prelove a result** of the penetration of salts from highly mineralized horizons (cryopegs) based on the results of the analysis of the geological section and comparison with the highest production rates of methane emissions (exactly where cryopegs, the most intense methane emissions were observed) [119].

Another side of the issue is the use of permafrost as a storage of liquid hydrocarbons or oil and gas production waste. The attractiveness of using the permafrost as a storage facility is explained by the almost complete impermeability of permafrost ~~to fluids~~ [49].

In the 1970s, storage facilities for waste from the development of mineral deposits were put into operation in the permafrost zone in the north of the Magadan Region, at Cape Schmidt, in the area of the Leningradskoye and Bovanenkovskoye deposits [49,295–297]. However, cases of dispersion of chemical elements in the frozen sediment around such storages have been recorded. For example, migration of buried drainage fluids was registered in the frozen sediment in Western Yakutia [49,297]. Dissipation halos were found, which led to the conclusion that active migration of chemicals into frozen sediment (**for more information on these works and migration processes, see Chapter 3**) [295,296]. Intense gas manifestation was recorded during the injection of industrial brines into the aquifers of the permafrost zone of Western Siberia, in the Daldynsky district, up to the release of a gas-water mixture and the flowing of gas with a high content of **heavy methane homologues, which may indicate the gas hydrate nature of the emission**. Later, in this area, two hydrate-bearing intervals were identified in the permafrost strata at depths of 80-100 and 140-190 m, as well as similar signs -

subpermafrost, but negative - temperature (within the permafrost zone) hydrate-bearing horizons in the depth range from 330 to 550 m [296].

Then a field experiment was carried out on the injection of brines into the expected hydrate-containing intervals for 3.5 months. During the experiment, **absorption of the salt solution**, its desalination and cooling by 1.5-2 °C were observed. The volume of released methane amounted to about 50 million m³ [296,297].

Thus, today there is a sufficient amount of information about gas hydrates, the forms and conditions of their existence in nature, as well as the causes of destabilization (pressure decrease, temperature increase, exposure to chemical compounds). However, the effect of salts on gas hydrates is considered mainly from the position of thermodynamics, and rare publications are devoted only to the assumption of a possible destabilization of gas hydrates in frozen sediment as a result of salt migration.

Chapter 4. The study of salt migration processes in frozen sediments

As noted above, one of the reasons for the hydrate dissociation in permafrost zone can be the migration of salts into the strata of frozen sediments containing gas hydrates. At the moment, in literature any special studies on the interaction between frozen hydrate saturated sediments and salt solution **are upsent**. While these studies are vital for assessing the intensity of methane emission during the permafrost degradation.

However, there are data on the study of salt migration in frozen sediments that do not contain gas hydrates, which is undoubtedly of interest for the problem under consideration, given the similarity of the ice and gas hydrate components of sediment. Before proceeding to the consideration

of existing works on the interaction between salt solution and frozen sediments, it should be reviewed the mechanism of ice melting, as one of the most important components of frozen sediment.

Salt ions can cause ice to melt in the pore space of frozen rocks. The melting of ice due to the migration of salt ions is a complex physical and chemical process, the intensity of which is determined by a number of factors. Ice is a crystal lattice composed of dipoles of water molecules. Salt ions, when in contact with ice, pull the dipole of water towards themselves, thus destroying the crystal lattice of ice - i.e. causing it to melt. So, in the case of the action of sodium chloride on ice, a negatively charged chloride ion attracts the water dipole with a positively charged end, and a positively charged sodium ion - negative (Figure 14). Thus, the salt ions are hydrated, causing the crystal structure of the ice to break down (melting), and hence a slurry of salts is formed whose freezing point is lower than the freezing point of pure non-salt water [298].




Figure 14. Scheme of a chemical reaction during the melting of ice due to the action of NaCl ions [298].

When a salt solution interacts with pore ice, salt ions migrate from an area of high salt concentration (brine) to an area of low concentration (films of unfrozen water around ice crystals) due to the so-called concentration diffusion. Salt ions cause the ice to melt, and the resulting moisture lowers the concentration of salts in the solution. An increase in the content of unfrozen moisture in the contact zone leads to the appearance of density inhomogeneities and a more pronounced convective mass transfer in this area [299–301]. In the case when the mass of the solution becomes

large enough compared to the mass of melting ice, the ice regime is considered to be quasi-steady. Ice melts until the salt concentration in the pore solution reaches the equilibrium concentration corresponding to the beginning of the solution freezing. The melting of ice is accompanied by a decrease in temperature, but the temperature levels off when the equilibrium concentration is reached [302–304].

At the same time, the temperature effect depends on the composition of the salts. So, when exposed to NaCl ions, the process of ice melting is endothermic (with heat absorption), and in the case of CaCl₂, it is exothermic (with heat release). Therefore, the effect of sodium chloride is slower than the effect of calcium chloride ions [298].

It should be noted that there is also the concept of the eutectic point for each salt solution. The lowest freezing point and the highest concentration of the solution (brine) corresponding to it is called the eutectic concentration at which solid matter crystallizes. If, as a result of lowering the temperature and increasing the concentration, the eutectic point is reached, then ice and salt crystals exist in the system, and not a liquid salt solution (Figure 15) [303].




Figure 15. Phase diagram of NaCl aqueous solution (modified after [305]).

It is according to the patterns described above that cryopegs exist in ~~the thickness of~~ permafrost. Cryopegs do not freeze, since the crystallization of salts dissolved in them requires deeper cooling, for example, to -21.4 °C for sodium chloride, or -55 °C for potassium chloride [101,302]. Such solutions are not in equilibrium with ice and can convert it into a liquid phase, eliminate the fluid resistance of frozen strata and migrate through cryogenic strata, including into horizons containing gas hydrates [41,49,51]. In this regard, special experimental and mathematical studies were carried out on the interaction of frozen rocks with salt solution.

4.1. Experimental research of salt migration processes in ice-containing sediments

Until a certain point in the development ~~of the history~~ of geocryology, frozen rocks were considered to be an absolute seal. However, in the ~~60s of the last century~~, based on field observations of the haloes of dispersion of chemical elements in frozen sediment in the north of Russia, it was concluded that the permafrost is an active environment where chemical and physicochemical processes take place: **moisture** migration, transport of chemical elements, structure and texture formation (ice release, swelling, cracking, etc.). Such field studies were carried out by **Safronov N.I., Glazovskaya M.A., Targulyan V.O., Shvartsev S.L., Ivanov A.M., Ivanov O.P., Makarov V.N., Pitulko V.M., Chibisov N .P. Shilo N.A., Vinokurov I.P. and some other scientists.** The researchers also noted the particular ecological danger of this phenomenon for an extremely sensitive complex multicomponent system of frozen sediment. In this regard, it became necessary to study and describe the mechanisms and patterns of salt ions migration in frozen sediment and to set up special experimental studies.

Separate experimental studies on the migration of chemical elements were carried out in the ~~60-80s of the last century~~, among which are the works of Murrman, **Hoekstra, Nacano (1971),**

devoted to the study of the migration of sodium ions in frozen bentonite pastes; Nechaev, Kan (1980), Romanov V.P. (1985) on salt migration in frozen sands; Ostroumova V.E. (1989), on the salt migration in frozen sand and loam in a gradient temperature field; as well as in general the work on the study of the migration of ions of various elements by Tyutyunov (1951), Derbenev (1967), Savelyev (1971) and some other scientists. After the 80s, the process of interaction of frozen rocks with salt solutions was studied in sufficient detail thanks to the works of I.A. Tyutyunov, M.A. Derbeneva, V.E. Ostroumova, E.A. Nechaev, E.V. Kan, V.P. Romanov, E.D. Ershov, E.M. Chuvilin, Yu.P. Lebedenko, I.A. Komarov, J.M. Marrion, Xu Xiaozu, B. Hallet, V.N. Makarov, K.V. Biggar, Murrman, Hoekstra, Nacano and other scientists.

In the investigations of E.D. Ershov, E.M. Chuvilin, Yu.P. Lebedenko, N.A. Benediktova, O.G. Smirnova on the basis of an experimental study and previous works, a description of the mechanisms of ion migration in frozen sediment is given. The authors explain the course of such a process as follows. Before the contact, there are two systems - solution and frozen sediment, in each of which the chemical potentials of moisture are in equilibrium, but not equal to each other. The same should be said about concentrations. After the contact, the sediment and the solution begin to interact as a single system in which the state of equilibrium is disturbed. In connection with the desire of the new system to again achieve a state of equilibrium, a moisture potential gradient arises in it, causing a flow of moisture into the frozen sediment, which coincides in direction with the salt flow. Salt ions capture water in solvate films, i.e. reverse osmosis happens. Over time, the flow of solution into the frozen sediment rapidly decreases. The process of migration of the solution into the frozen sample causes the thawing of the sample and the formation of the thawed part of the sample. Due to the desalination of the migrating solution, in the frozen zone partial freezing of residual water occurs and rise in water - ice content, immediately before the thawing front [43,45,48,286,306].

The main media for ions migration in frozen sediment are films of unfrozen water that exist on the mineral particles surface, as well as between boundaries of pore ice crystals. The capillary and film unfrozen water are distinguished [104]. Also, the flow of salt ions through the ice component of frozen rocks is carried out through films of quasi-liquid water. A quasi-liquid layer is a disordered transitional structure of water in a solid and liquid state, the thickness of which can vary from several tens to hundreds of angstroms [307,308]. The physical role of the quasi-liquid layer is to neutralize the effect of instability caused by the breaking of hydrogen bonds at the boundary of the ice structure. In addition, this film serves as an optimal transition medium for restructuring during the mass transfer of ice with surrounding water phases, including saline. The thickness of the layer of the liquid phase of water decreases with decreasing temperature up to its complete disappearance at a temperature of about -45°C . Also [309] notes that the content of the liquid phase is the greater, the higher the dispersion of rocks. So, in clay soils, the content of unfrozen moisture can reach 13%, while in sand its content does not exceed 1%.

~~Ice is an important component of frozen rocks.~~ Depending on the structure of pore ice, gas inclusions, density, size of crystals, presence and composition of impurities, its ion permeability will be determined [48]. Air inclusions in ice (and, accordingly, less dense ice) contribute to a decrease in the flux of salt ions. The main flow of salts is carried out along the contact of ice crystals, respectively, in single-crystal ice, where the flow occurs along structural deformations, dislocations, the migration of ions is lower than with an increase in the number of crystals [97].

As driving forces of mass transfer in frozen rocks Ershov E.D. et al. pointed out the electrokinetic potential gradient (due to the difference in ion concentrations on the surface of the rock skeleton, in the pore and in the external solution), adsorption forces (due to interaction with the particle surface), and electrostatic forces (due to the distribution of electric charge over the surface of negatively charged minerals and positively charged pore ice). A similar statement, but using a

different term, was given in the work of Nechaev, Kan (1980) and Romanov V.P. (1985). They pointed out that the migration of salt ions is influenced by electrosurface properties. The authors attribute the migration of metal ions to the action of electrochemical processes in pore waters. As the driving force of moisture migration, the magnitude of the gradient of the total thermodynamic potential of ground moisture or the Gibbs free energy is taken. The thermodynamic potential is the sum of particular values [310,311].

$$\mu_w = \psi_{\text{KK}} + \psi_p + \psi_o + \psi_{es}$$

ψ_{KK} – frame-capillary thermodynamic potential;

ψ_p – hydrostatic thermodynamic potential;

ψ_o – osmotic thermodynamic potential;

ψ_{es} – electrostatic thermodynamic potential.

When the thermodynamic parameters change, the migration of moisture and chemical elements is possible, since the system comes into non-equilibrium both in terms of the chemical potential and in terms of the general thermodynamic potential. The authors also note that the osmotic potential decreases with increase in salt concentration in pore solution. When the salt concentration drops to a certain minimum value, moisture transfer in the system may stop due to the equality of the total thermodynamic potential of ground moisture to the osmotic potential of water in the external solution [310–312].

One of the most difficult questions to study was the question of the mechanisms of migration of salt ions in frozen rocks. Back in 1965, M.M. Derbeneva wrote that the transfer of salt ions occurs due to moving films of unfrozen water and due to diffusion with an existing difference in concentrations, that is, the presence of a concentration gradient. Later, these two mechanisms were named by E.D. Ershov et al. (1989) convective and concentration diffusion, respectively, and they also singled out a third mechanism, adsorption [286,312].

Concentration diffusion is characteristic of the process of salt migration due to a concentration gradient at high negative temperatures, when the content of unfrozen moisture is relatively high. In this case, the migration of salt ions occurs along the films of unfrozen water from the region with the highest concentration to the region with a low concentration of salt ions [60,97].

Convective transfer occurs at high flux densities of salt ions into the frozen rock and is usually observed at high negative temperatures. At the same time, a feature of this mechanism is the joint moisture and salt transfer, that is, not only the migration of salt ions takes place, but there is a joint movement of water with salt ions. And due to this mechanism, there is an intensification of mass transfer, structure-forming and physico-chemical processes [60,97].

As the temperature decreases, the convective component of the total ion flux decreases, but rather high values of migration fluxes continue to be observed. This is no longer due to the action of the convective mechanism, but to the activation of other ion transport mechanisms, primarily adsorption [48,60,97,313].

The adsorption mechanism of salt transfer occurs in frozen sediments in the range of low negative temperatures (from -20 to -60 °C) due to the exploit of the ice surface forces. In this case, the thickness of unfrozen water films becomes equal to or close to the size of salt ions and migration almost completely stops [46].

It should also be noted the work carried out on the study of deformation processes in frozen rocks in the process of salt migration. The general nature of the deformations is positive. The most important contribution is made by the processes of swelling and swelling due to the migration of the solution in frozen samples. The development of deformations is directly related to the change in the density of the migration flow [45].

Another important aspect of the interaction between salt solution and frozen sediments is the temperature changes that occur during this process. E.D. Ershov et al. (1993) studied the heat fluxes

arising from the interaction of frozen rocks with salt solutions, which made it possible to interpret the phenomena occurring in the process of migration. Thus, the authors argue that before the start of contact with the solution, a stationary temperature field was observed in the sample, i.e., the temperature in the soil did not change, and the heat fluxes were constant and close to zero [45]. After the contact of the soil and the solution, the incoming migration flow increases the concentration of salts in the frozen rock, and, consequently, lowers the temperature of its thawing. As a result, “ice-water” phase transitions occur in the frozen ground. The temperature of the sample and solution at the initial moment of the experiment is constant, but as soon as the salt concentration increases, phase transitions begin, which, in turn, are associated with thermal effects. In case of defrosting, heat is absorbed. In the frozen zone of the sample, near the thawing front, schlieren evolution takes place, which goes along with the release of heat. Thus, in the interacting soil-mortar system, phase transitions occur that occur in two areas: in the thawing zone and in the schlieren zone. This causes a complex picture of heat flows in frozen rocks interacting with salt solutions [45].


As described earlier, the migration of salt ions in frozen rocks occurs due to the difference in chemical potentials. And it, in turn, is determined by the temperature gradient, humidity, dispersion and mineralogical composition of the soil system [306]. Hence, separate studies are devoted to the influence of various factors (external and internal) on the process of salt migration in frozen rocks. The ions migration is predominantly defined by the features of the frozen sediments: moisture content, mineral and chemical composition, dispersion, cryogenic structure, and also external parameters: salt solution concentration, interaction temperature, etc. [45,314,315]. Yu.P. Lebedenko (1989), E.M. Chuvilin (1999), E.D. Ershov et al. (1999, 1998) and others.

For instance, studies demonstrated, that total ions permeability and accumulation can increase with the rise in the ice and water content in the pore space of frozen sediments [46].

The study of the effect of temperature showed that with increasing temperature, the intensity of ion migration increases, while the nature of deformations changes as a result of mass transfer processes. As the temperature decreases, bound water films freeze in the rock, the density of the migration flow decreases, and the accumulation of ions has an extreme character. At temperatures down to -5 ... -10 °C, swelling, swelling, ice release, soaking and sedimentation are observed. And at lower temperatures - the processes of brittle fracture. These regularities are characteristic mainly for finely dispersed soils at high negative temperatures, while the second type of deformations is characteristic of sediment [45,312].

The authors also observed that with the increase in the salt concentration intensifies the salt migration and an increase in the content of unfrozen moisture. In frozen rocks, due to the freezing of part of the water, the negative thermodynamic potential due to the action of the free unspent surface energy of mineral particles is significantly reduced. As the concentration of the external solution further decreases, a gradient of the thermodynamic potential of moisture arises, directed from the solution to the frozen rock [97].

Ershov et al. (1999), when analyzing the migration of ions of various salts in frozen rocks, found that the activity of ions decreases in the series [312,315]:



However, depending on the granulometric composition, the dependence may change. So, for example, in clays, the mobility of Ca^{+2} is much lower, which is due to its adsorption characteristics, that is, the mobility of cations and anions is also determined by their adsorption capacity. Also interesting are the results of an experimental study on ice release depending on the composition of the contacting solution. So in the series KCl, NaCl, MgCl_2 , CaCl_2 , ice release increases [97].

With an increase in dispersion, the accumulation of moisture and ions increases in the sand-clay series. This is explained by the fact that the ion permeability of frozen rocks decreases (that is,

salts penetrate a shorter distance) with increasing dispersion, while more dispersed soil has a greater exchange capacity (sorption), i.e. the ability of the rock surface to bind salt ions, as a result of which their mobility decreases, and, consequently, the intensity of the migration flow decreases [48,315]. At the same time, in a series of clays, the greatest accumulation is observed in kaolinite clay, the smallest in montmorillonite (weak moisture and ionic conductivity due to the high sorption capacity of montmorillonite) [315].

The main role in **coarsely dispersed rocks** is played by concentration diffusion through pore ice, since there is no unfrozen moisture or its content is extremely low (up to 1%). With an increase in dispersion, the amount of unfrozen moisture increases, and along with diffusion, convection begins to play a role in the transfer of ions. However, as noted above, sorption is also observed, which reduces the ion permeability in **finely dispersed rocks** [313].

Ershov et al. also showed different migration abilities of cations and anions using the example of the migration of sodium ions and chlorine ions. During the experiment, chloride ions migrate more from solution to frozen rock than sodium ions, which is due to the greater migration ability of chloride ions [45].

Another study shows, that the increase of **compression pressure** cause the decrease of salt ions migration in sediments with low dispersion (mainly clay-containing sediments). Thus, in clays there is a certain critical pressure at which the migration of salt ions almost completely stops, since the convective component of the migration flow, which provides the highest values of salt transport, is suppressed. While in the sands such an effect is not observed [313].

~~Now there is a point to review the existing mathematical models of the interaction between salt solutions and frozen sediments for calculating the main diffusion characteristics.~~

4.2. Existing physico – chemical models of salt transfer processes in sediment media

To describe the transfer processes in **frozen soils**, several concepts of mass transfer, the concept of suction forces, the concept of crystallization forces, and the osmotic pressures concept have been proposed, ~~which correspond to certain~~ experimental facts. One of the most widely applicable is the approach based on effective transport equations. This model is established on the laws of energy conservation and mass of pore moisture, supplemented by moisture migration under the action of a moisture gradient, and/or a temperature gradient, and/or any other gradients. However, this model also has its own difficulties, mainly associated with the need for experimental data (obtaining diffusion coefficients), but also with the determination of boundary conditions on the phase transition surfaces. Therefore, this model was partially **supplemented G.G.Tsyarkin in 2009** [316].

In general, the salt migration in thawed sediment is defined by the processes of concentration diffusion. And it, in turn, is described by the ~~first~~ Fick's law, which was formulated back in 1855. ~~Therefore, mathematical modeling of the processes of diffusion of salt ions is mainly based on Fick's law, which~~ has the form of the following equation:

$$\Phi = -K \frac{\partial C}{\partial x}$$

where Φ – rate of salt ions diffusion; K – salt diffusion coefficient; C – concentration of dissolved components.

However, E.D.Ershov (1999) pointed out the imperfection of the calculation scheme for diffusion coefficients through Fick's law for frozen rocks, since the coefficient of proportionality between the total ion flux and the concentration gradient is calculated and there is no consideration of other mechanisms of salt transfer besides diffusion: convective and adsorption. To obtain the actual parameters of the migration of salt ions, it is necessary to take into account all the components in the total ion flux, and not just diffusion. Nevertheless, on the basis of this law, numerous schemes for

calculating diffusion coefficients in thawed soils have been proposed [317,318], and several schemes in frozen and freezing soils [49,50,319–321], however, no universal one has been singled out. Moreover, there is no such scheme for frozen hydrate-bearing rocks.

The second important part of the mathematical model for describing transport processes with phase transitions, which are undoubtedly observed during the interaction of salt solutions with frozen rocks, is based on the Stefan problem. The classical Stefan problem assumes that the phase transition is localized in a narrow region, modeled by a front, which is the weak discontinuity surface for the temperature function. The position of the phase transition front relative to the medium changes with time and is in the process of solving the problem [316].

The Stefan problem is based on the law of conservation of energy, and heat transfer in a stationary medium is described by the heat equation. In the one-dimensional case, the melting problem has the form:

$$\frac{\partial T_j}{\partial t} = a_j \frac{\partial^2 T_j}{\partial x^2},$$

where solid phase takes the area:

$$j = 1, \quad X(t) < x < \infty, \quad t > 0,$$

and liquid phase – area:

$$j = 2, \quad 0 < x < X(t), \quad t > 0$$

The initial and boundary conditions can be described in the following form:

$$X(0) = 0; \quad T_1(x, 0) = T_0 < T_f, \quad t > 0; \quad T_2(0, t) = T^o > T_f, \quad x > 0$$

As noted above, the main medium for the transfer of salt ions in frozen rocks are films of unfrozen water, which means that the processes of transfer of salt ions should be determined by the amount of unfrozen **moisture**. So, the higher its content, the more intense the process of salt migration. In non-saline soils at negative temperatures, the connection among the temperature T and unfrozen

water content W_{unf} and is unambiguously designated by the “unfrozen moisture curve”. Though, the quantity of unfrozen water W_{unf} in saline sediments is determined also by pore solution concentration. Moisture migration followed by the increase in moisture content causes temperature increase in frozen sediments. On the contrary, a salt concentration increase in the pore solution causes temperature decrease. The salt concentration in the pore solution C during the frozen sediments thawing changes in accordance with the dependance:

$$C = \frac{C_0 W}{W_{\text{unf}}}$$

where W – total water saturation in the frozen sediment; W_{unf} – unfrozen water content in the frozen sediment; C_0 – initial salt concentration in the pore solution.

Salt diffusion coefficient is one of the main parameters of the salt ion migration process in frozen sediments, which depends on various factors. First of all, it is determined by the sediment dispersity, the geometry and relative position of the ice crystals and water films, and the parameters of the water films in the pore space of sediments.

One of the main parameters of the salt migration in frozen sediments is the diffusion coefficient of salt ions, which depends on many parameters. First of all, it depends on the type of soil, on the thickness of water films in the pores of the soil, on the geometry and relative position of water films and ice crystals. Thus, the greater the number of crystals present in the pore space, the higher the diffusion coefficient will be, since the migration of ions occurs along the boundaries of the crystals.

One of the schemes for calculating the diffusion coefficient was proposed by V.I.Aksenov et al. (2010). The authors propose to calculate the salt diffusion coefficient in frozen soil using the following formula [322]:

$$D = W \cdot D_0$$

where D_0 – diffusion coefficient of salt in free water; for sodium chloride (NaCl) D_0 is approximately $1.1 \cdot 10^{-5}$ – $1.4 \cdot 10^{-5}$ cm^2/c .

Another design scheme was proposed by Ershov E.D., Komarov I.A., Chuvilin E.M. in 1997 [49]. The salt transfer model is presented with the following assumptions:

- convective-diffusion mass transfer of salt ions occurs under isothermal conditions;
- there is no interaction of salt ions with pore solutions with the release (absorption) of heat;
- ion migration occurs in a semi-infinite array evenly along the height of the well;
- one-dimensional process;
- taking into account the pore space characteristics, the fracturing of the host sediments is carried out using the effective porosity parameter n_e , and the **massif** is considered as quasi-homogeneous;
- the concentration of salts in the operating brine is taken constant;
- the influence of pressure and the processes of mechanical interaction of brines with the rock is not taken into account.

The problem of salt transfer is solved in a self-similar approximation, and the problem of mass transfer in a quasi-homogeneous massif and its solution have the form [49]:

$$D \frac{\partial^2 C}{\partial x^2} - v \frac{\partial C}{\partial x} = n_e \frac{\partial C}{\partial \tau}$$

$$C(\infty, \tau) = C(x, 0) = C_f$$

$$C(\infty, \tau) = C_s$$

where C – current salt concentration; D – diffusion coefficient; n_e – porosity; x – current coordinate; v – average volumetric velocity of the filter flow, determined through the flow rate; C_s – ions concentration; C_f – initial concentration of that ions in pore solution of the sediment.

$$\bar{C} = \frac{C - C_\varphi}{C_p - C_\varphi} = 0.5 \left[\operatorname{erfc} \theta + \exp(P_e) \operatorname{erfc}(\theta^2 + P_e)^{\frac{1}{2}} \right]$$

where

$$\theta = \frac{n_e x - v \tau}{2(nD\tau)^{\frac{1}{2}}}, \quad P_e = \frac{vx}{D} - \text{Peclet criterion } \operatorname{erfc}(\cdot) =$$

$1 - \operatorname{erf}(\cdot)$, $\operatorname{erf}(\cdot)$ – Gaussian error function.

The diffusion coefficients of chlorides in frozen rocks obtained using this calculation scheme were $10^{-6} - 10^{-7} \text{ cm}^2/\text{h}$ [49]. While in thawed rock the diffusion coefficient varies from 10^{-6} to $10^{-10} \text{ cm}^2/\text{sec}$, and in ice, the diffusion coefficient can range from 10^{-7} from $10^{-9} \text{ cm}^2/\text{sec}$ [313].

~~Relatively recently, work has appeared on the~~ migration of sea water on the Arctic shelf and the influence of this process on the evolution of permafrost zone and HSZ. ~~As shown in the work of V.V. Malakhova and V.E. Eliseev (2020),~~ salt (for example sodium chloride) remarkably decreases the temperature of hydrate stability comparatively to the temperature of unsalted water. There is also a corresponding change in the pressure of the thermodynamic stability of hydrates [57].

The authors compared the displacement of the depth of the gas hydrate stability zone with an increase in seawater salinity, and also compared two calculation schemes. Thus, the shift in equilibrium temperature is calculated using the expression:

$$T_D = \frac{T_{D,ref} \cdot \log(1 - x_s)}{\log(1 - x_{s,ref})}$$

$$T_D = T_{D,ref} \cdot \log(1 - x_s) / \log(1 - x_{s,ref})$$

where $T_{D,ref}$ – temperature reference value for hydrate decomposition at the corresponding reference salt content value $x_{s,ref}$ and x_s – mole fraction of salt in pore water.

In work [175] a method is proposed for predicting the shift in the equilibrium temperature of a hydrate in a free volume based on information on a decrease in the freezing point of aqueous solutions containing various salt concentrations in relation to the freezing point of fresh water (ΔT_F):

$$T_D = 0.6825 \cdot \Delta T_F$$

In the calculations, the authors also used the dependence of the freezing point of water ($^{\circ}\text{C}$) on the salt concentration S (‰) and pressure in bottom sediment P (MPa) is presented in the work [323]:

$$T_F = -0.073P - 0.064S$$

Salt diffusion coefficient was taken equal to 10^{-9} m^2/sec in accordance with estimates based on drilling data in the Laptev Sea [57,324].

Based on the modeling results, the authors argue that an increase in salinity causes degradation of the gas hydrate stability zone and leads to a displacement of the boundary down the section. However, the authors note that due to the great depth of hydrate occurrence on the Arctic shelf, the influence of salts reaches no more than a few percent. It should be noted that these calculations do not take into account the presence of metastable gas hydrates, which can be affected by salts and which, as noted in Chapter 2, are especially sensitive to changes in external conditions, the authors also point out this fact, but without taking it into account in the calculations. In addition, the diffusion parameters of salt ions are not calculated, but are given according to previous calculations, which means that the power of the ion flow and the time factor are not taken into account.

Another study is dedicated to the hydrate decomposition as a result of the migration of salt drilling fluid into hydrate-bearing rocks (in the thawed state) [325]. According to the proposed calculation scheme, the phase equilibrium pressure and phase equilibrium temperature for hydrate-bearing rocks satisfy the exponential function:

$$P_{eq} = e^{AT_{eq}+B}$$

The authors assume that drilling fluids consisting of sea water and may be comparable to a sodium chloride solution with a mass fraction of 0.03.

$$A = 0.5212w^2 - 0.0609w + 0.1109$$

$$B = -119.76w^2 + 18.638w - 29.306$$

Rate of hydrate dissociation connected with hydrate equilibrium pressure in the reservoirs:

$$\frac{dc_H}{dt} = k_d^0 \cdot e^{-\frac{E}{RT}} \cdot A_{hs} \cdot (P_{eq} - P_g) = k_d^0 \cdot e^{-\frac{E}{RT}} \cdot \sqrt{\frac{\varepsilon^3 (S_l + S_g)^3}{2k}} \cdot (P_{eq} - P_g)$$

The porous medium conservation equations describe changes in mass and temperature per unit volume of the formation. Takes into account the porosity and properties of the porous medium:

$$\text{Gas phase} \quad \frac{\partial}{\partial t} (\rho_g S_g \varepsilon) + \nabla \cdot (\rho_g \vec{v}_g) = -x_g \frac{dc_H}{dt}$$

$$\text{Liquid phase} \quad \frac{\partial}{\partial t} (\rho_l S_l \varepsilon) + \nabla \cdot (\rho_l \vec{v}_l) = -x_l \frac{dc_H}{dt}$$

$$\text{Hydrate phase} \quad \frac{\partial}{\partial t} (\rho_H S_H \varepsilon) = \frac{dc_H}{dt}$$

$$\text{Energy} \quad \rho_{eff} c_{eff} \frac{\partial T}{\partial t} + \nabla \cdot (\rho_g \vec{v}_g c_{pg} \Delta T + \rho_l \vec{v}_l c_{pl} \Delta T) = \nabla \cdot (k_{eff} \nabla T) + \dot{q}$$

$$\text{Salts} \quad \frac{\partial}{\partial t} (\rho_S S_S \varepsilon w) + \nabla \cdot (\rho_S \vec{v}_S) = 0$$

Methane is characterized by changing density and follows the equation of state for an ideal gas.

$\nabla \cdot (\rho_g \vec{v}_g c_{pg} \Delta T + \rho_l \vec{v}_l c_{pl} \Delta T)$ - convective energy transfer initiated by gas migration.

Salt migration can be described by the equation:

$$\vec{v}_S = w \vec{v}_l - D_S \nabla w$$

Cohesion determines the hydrate-bearing sediment strength and it is related to hydrate content.

$$C = C_0 (1 + \psi (S_H - S_0))$$

This calculation scheme was proposed by Wei et al. with the following assumptions [325]:

- The model is limited to 1 m from the wellbore;
- 2D model;
- External boundary conditions are the initial reservoir parameters;
- Internal boundary conditions are downhole pressure and temperature.

Based on this model, the authors calculated the depth of dissociation of gas hydrate formations from the well during the penetration of saline [325].

In general, today there are practically no works on experimental modeling of the interaction of salt and drilling fluids with frozen hydrate-saturated rocks, just as there are no attempts to describe the process of such interaction and obtain diffusion characteristics, including through mathematical modeling. In this regard, setting up experimental modeling to study the mechanism and patterns of hydrate dissociation during interaction of frozen sediments and salt solution is of particular interest to ensure the safe and uninterrupted construction of exploration and production wells in the area of permafrost.

Thus, today there is a lot of information about the processes of mass transfer in frozen sediments during their interaction with salt solutions, which creates good prerequisites for setting up special experimental studies to study the behavior of frozen hydrate-bearing rocks under saline conditions. The specificity of permafrost sediments: the presence of driving forces of moisture migration. At high negative temperatures, the migration flow of moisture has a significant effect on ions migration in finely dispersed sediments, which is typical for ions of light elements.

Chapter 5. Method of experimental study of salt ions migration in hydrate-saturated sediments

Series of experiments were conducted in order to study the mechanisms and dependences of methane hydrate dissociation in porous media of frozen sediments during their interaction with salt solutions at the range of negative temperatures ($-3 \dots -20 \text{ }^{\circ}\text{C}$) in equilibrium (at the pressure $>2 \text{ MPa}$) and non-equilibrium (at a pressure from atmospheric - 0.1 MPa to 2 MPa) conditions. ~~An experimental study on the interaction between salt solution and methane hydrate saturated frozen sediments. The experiments procedure is following:~~

1. Artificial hydrate saturation of frozen sediments samples;
2. Determination of the initial parameters;
3. Bringing frozen samples, containing methane hydrates, and salt solution into the contact;
4. Assessment of the main characteristics change (changes in water, hydrate content and salt ions concentration) during the interaction between salt solution and hydrate-saturated frozen sediments, and determination of the main interaction parameters.

5.1. Experimental setup and preparation technology of hydrate saturated sediments

The technique for obtaining frozen hydrate saturated media included preparation of sediment samples with given water content, locating samples in the pressure cell, hermetic closure of the pressure cell and the creation of a vacuum, increase of the pressure up to the hydrate equilibrium values by the methane injection of CH₄ [326,327].

The sediments are saturated with hydrates using a special technique that allows samples of hydrates evenly distributed in the porous medium of the sediments to be obtained. The initial hydrate saturation of the sediment sample was carried out at a negative temperature of -4...-6°C. In order to increase the hydrate saturation of the sample, a heating-cooling cycle was made with the transition through 0°C. When the gas pressure is higher than the stable pressure, the thawing and freezing of the aqueous compound sample leads to the activation of the hydrate formation process and the increase of the overall hydrate saturation of the sample. The formation of natural gas hydrates happens on the pore ice surface, which inhibits the water transfer process and makes the natural gas hydrates evenly distributed in the sediment samples [327].

For artificial hydrate saturation of sediment media, a special pressure cell was used (Figure 16). Pressure cell volume is 419 cm³ and it was designed for pressures up to 20 MPa.



Figure 16. Equipment (pressure cell) for saturation of sediment by hydrate.

Initially, sediment used for experimental modeling was saturated with a set water content (around 11-12%). Then wet sediment was placed in special perforated plastic containers of cylindrical shape, which height equals to 6-9 cm and diameter - 3 cm. Sediment samples in these plastic containers were then frozen for 1 day at temperatures of $-6 \dots -8 \text{ }^{\circ}\text{C}$ to limit any potential migration of water, which can lead to the non-uniformed distribution of pore hydrate.

Frozen sediments in plastic containers were then placed in the pressure cell. The pressure cell was able to fit from 5 to 7 perforated plastic containers with the studied frozen sediment. A steel cover was hermetically fastened to the body of the cell through a teflon gasket with studs, on which a pressure gauge with a division value of 0.05 MPa was used. After the pressure cell was sealed and depressurized, and cooled hydrate-forming gas (methane, volume fraction 99.98%, $t = -6 \dots -8^{\circ}\text{C}$) was supplied into the system. Then, the pressure cell with samples of frozen sediments is placed in a freezer. First, a cooling cycle ($t = -10 \text{ }^{\circ}\text{C}$) is set, lasting about 3-5 days, then a defrosting cycle ($t = 0 \dots +1 \text{ }^{\circ}\text{C}$), lasting about 1-2 days. Such a change of cycles is carried out 3-4 times. By transmission of the sediment samples through negative temperature by cycles the bigger amount of the residual ice, which was not transferred in the hydrate form before, was transited to form hydrate afterward.

Due to that, sufficient saturation with methane hydrate of the sediment porous media is achieved. Thus, the saturation of the studied frozen sediment samples with hydrate occurs within 2 weeks to a month.

Afterward of the hydrate saturation, hydrate-containing samples were cooled to the temperature -6 ± 1 °C in the pressure cell. Meanwhile, the residual pore moisture, which did not transform into hydrate, freezes out. This approach makes it possible to obtain sediment samples with a high degree of pore filling with hydrates up to 60% or more [23]. ~~Thus, several frozen hydrate-containing samples were obtained, which were in equilibrium conditions in the pressure cell.~~

Then, gas pressure was reduced to the atmospheric (0.1 MPa) in the pressure cell at the negative temperature (6 ± 1 °C). Subsequently, part of the samples for experimental modeling at the atmospheric conditions were kept in a non-equilibrium state for about a day to stabilize the self-preservation process. After that, the frozen hydrate-saturated samples were removed from the pressure cell outside at a negative temperature. Due to self-preservation effect at negative temperatures, the porous hydrate formations in a frozen state at a temperature of -6 ± 1 °C had good preservation for a long time, during all time of the experiment (1-2 days and longer).

Before the contact with salt solution, several characteristics for frozen hydrate-containing samples, including weight moisture (W, %), density (ρ , g/cm³), and specific gas content were obtained.

The appearance of frozen hydrate-saturated sediments before the experiment is shown in Figure 17. Subsequently, artificially saturated hydrate samples were used in experiments on interaction with salt solutions.




Figure 17. The appearance of frozen sand saturated with pore hydrate after extraction from the pressure cell.

All obtained samples had similar characteristics ($W=11-12\%$, $n=35-40\%$, $K_h=40-60\%$) and were used in the experiment as “twin” samples. Figure 18 shows an example of the distribution of water content, porosity, and amount of water in the hydrate form (hydrate coefficient) by intervals of obtained frozen hydrate saturated sediment samples. Water content by the length of the samples varies with a difference of around 1-2%, porosity – around 2-5%, and hydrate coefficient 5-7%.




Figure 18. Distribution of water content, porosity, and hydrate coefficient by the length of frozen hydrate saturated sediment samples.

After measuring the initial parameters, the hydrate-saturated frozen samples were brought into contact with the test saline solution at the end face under different thermobaric conditions.

5.2. Sediment characteristics used in experimental simulation

For the experimental modeling, both model sediments (clean quartz sand and sand with the addition of clay particles of kaolinite or montmorillonite) and natural sediments are taken in the areas of gas manifestations. Parameters of gas emission here indirectly demonstrates their possible connection with natural porous hydrate [113,236]. Sediment characteristics are shown in Table 4.

Table 4. Particle size distribution, mineral composition, and salinity of investigated sediments.

Sample	Place of Sampling	Size of Sediment Particles, %						Mineralogy	Salinity, %
		1 - 0.5	0.5 - 0.25	0.25 - 0.1	0.1 - 0.05	0.05 - 0.001	< 0.001		
Fine sand 1	Moscow region	6	7	80	2	3	2	>98% quartz	0.01
Fine sand 2	Moscow region	0	2	80	17	1	0	>98% quartz	0.01
Medium sand	Moscow region	0	63	25	11	1	0	>98% quartz	0.01
Fine sand 3	South Tambey gas field	0	12	63	22	2	1	>98% quartz	0.11
Sandy loam	Laptev Sea, Buor-Khaya	1	9	52	20	16	2	54% quartz 41% microcline + albite 4% illite	0.4

*The listed mineral phases have percentages > 1%

The determination of the mineral composition was carried out using an X-ray diffractometer (XRD) in the laboratory of sediment science and technical sediment reclamation of the Department of Engineering and Ecological Geology, Faculty of Geology, Lomonosov Moscow State University. The granulometric composition was obtained by the sieve method. While composition and content of water-soluble salts was defined by the method of water extracts on the flame photometer PFP-7 (Jenway) [328,329].

The physical characteristics of the samples were determined according to GOST 5180-84 [330].

Fine sand-1, fine sand-2 and medium sand are represented by homogeneous quartz sand (quartz > 98%) of marine upper Jurassic genesis mJ₃ with the predominance of 0.1-0.2 mm fraction (fine

sand-1 and fine sand-2 around 80%), and 0.2-0.5 mm (medium sand 63%). Fine sands of light yellow color, homogeneous, well sorted. Non-saline (salinity is 0.01%). The specific active surface is 0.6 g/m². The density of the solid component is 2.65 g/cm³.

In addition, two sediments were obtained during drilling in permafrost area, where by indirect signs hydrate formation may exist, and they represented by:

- Fine sand 4 is a marine deposit from the upper Pleistocene (mQ_{III}²⁻³). The sampling site is a permafrost zone close to the South Tambey gas condensate field at a the depth of 32 - 35 m. Sand is light grey, well-sorted, homogeneous in grains, and composed of quartz (>98%). Characterized by predominance of 0.1-0.2 mm (63%), and also 0.1-0.05 mm (22%) fraction. The density of the solid component is 2.68 g/cm³. Salinity is 0.11% presented mainly by sodium chloride composition.
- Sandy loam, a polymineral of alluvial genesis from the Neopleistocene (aQ_{II}). It was extracted during drilling of the permafrost zone of the Arctic shelf in the western part of the Buor-Khaya Bay (Laptev Sea), 17 km northeast of the island Muostakh, where parameters of gas emission indirectly demonstrates their possible connection with natural hydrate existence [30,113,264,331,332]. The depth of sediment sampling is 43-45 m. The sandy loam is light gray, and rounded, with the predominance of 0.1-0.2 mm fraction (52%), and 0.1-0.001 (36%). The mineral composition is dominated by quartz – 54%, microcline+albite – 41%, and illite – 4%. The density of the solid component is 2.72 g/cm³. Salinity is 0.4% represented mainly by sodium chloride composition.

Salinity in natural sediments was decreased to 0.1-0.05% by removing salts through washing with distilled water several times.

In addition to the described sediments, special model mixtures were prepared from fine sand-2 and clay. As clay contaminants, kaolinite and montmorillonite were taken in different percentages (up to 15%):

- Montmorillonite clay (eP_{2ogl}) was sampled near the Dzembel train station, in Turkmenistan. It is presented with montmorillonite (94%), andesite (3%) and biotite (3%).
- Kaolinite clay (eP₂) was taken at the town Novokaolinoviy in the Chelyabinsk region. It is presented by kaolinite (92%), quartz (6%), and muscovite (2%).

Described above model sediments and natural sediments of disturbed composition were used to form frozen hydrate-saturated sediment samples for experimental modeling. The sand was given a moisture content of about 11-12%. Cylindrical sediment samples were formed, about 10 cm long and 3 cm in diameter.

5.3. Technique for studying the interaction of frozen hydrate-containing sediments with saline solutions in non-equilibrium and equilibrium conditions

The study of salt migration processes in frozen hydrate-containing sediment was based on the experimental modeling of the interaction between salt solution and frozen hydrate-bearing sediment samples at the fixed temperature below zero and various pressures. Its procedure is following:

- 1) frozen sediment samples were filled with water up to 12% and consequent preparation with a given water content (12%) and then saturated with methane hydrates in a special pressure cell;
- 2) the saturated hydrate sample is frozen in a pressure chamber and transferred to a metastable state by reducing the pressure to atmospheric pressure at a negative temperature freezing hydrate-

saturated samples in a pressure cell and transferring them to the metastable state by reducing the pressure to atmospheric at a negative temperature;

3) frozen samples, saturated with hydrate, was extracted from the pressure cell, measuring the initial parameters (initial water content, density, initial hydrate coefficient K_h^{in}), and bringing them into the contact with a frozen NaCl solution in the state of salted ice at a constant negative temperature and various pressure conditions (below (<2MPa) or above (>2 MPa) equilibrium);

4) monitoring the process of interaction between salt ice and frozen hydrate-containing samples in time.

The technique for obtaining frozen hydrate-containing media is described in chapter 5.1. As a result of artificial hydrate dissociation of frozen sediments, 5-7 samples-twins were obtained with unified water content, porosity, and hydrate distribution.

Afterwards, the gas pressure drops to 0.1 MPa at temperature $-6\text{ }^{\circ}\text{C}$, transferring the frozen hydrate-bearing sediment samples into metastable conditions. Frozen hydrate-saturated samples can maintain a sufficiently high hydrate content for a long time as a result of self-preservation effect manifestation. Then, the frozen saturated hydrate samples are removed from the pressure cell at temperature below zero. The samples had massive ice-hydrate structure, and the pore hydrate content is evenly distributed by the height of the sample.

Subsequently, the pressure in the pressure cell, which was at a negative temperature, was dropped to 0.1 MPa, transferring frozen hydrate formations in sediment samples to the metastable state. Due to the self-preservation effect, frozen hydrate-saturated samples retained a sufficiently high hydrate content for a long time. Then the frozen hydrate-saturated samples were removed from the pressure cell to the outside at the negative temperature. The samples had massive ice-hydrate texture with pore hydrate contents uniformly distributed over the sample height [23,327].

The initial petrophysical characteristics (moisture content, density, initial hydrate coefficient K_h^{in}) were determined for the obtained frozen hydrate-containing samples before their contact with the salt solution. The samples were characterized by a uniform distribution of the initial characteristics along with the height of the sample, as well as close values between the compared samples. Thus, the initial moisture content of the samples was within 11-12%, porosity 35-40%, density 1.8 g/cm^3 , initial hydrate coefficient $K_h^{in}=0.4-0.6$. Before and after the experiment on the interaction of hydrate-saturated sediment with a saline solution, a sample interacting with a saline solution was photographed.

Experiments on the study of salt transfer in frozen sandy sediments containing porous gas hydrates were carried out at a temperature from -2.5 to $-20 \text{ }^\circ\text{C}$ and pressures from atmospheric to 4 MPa. The contacting solution was a NaCl with a concentration from 0.1 to 0.4N, and also with KCl, Na_2SO_4 , CaCl_2 , MgCl_2 solution with a concentration of 0.1 and 0.2 N.

In the course of experimental modeling on the interaction between salt solution and frozen hydrate saturated samples at atmospheric pressure, the samples were brought into contact with salt solution when the gas pressure in the cell was decreased to 0.1MPa and initial parameters were measured. Usually, around 5-7 samples were brought into the contact at the same time for comparison. Contact of the hydrate saturated frozen sample with the salt solution was made in a sealed container to prevent sublimation processes, and the entire setup was placed in a thermal box to maintain a constant negative temperature (Figure 19). During all experimental modeling under atmospheric pressure, the negative temperature was fixed (from 2.5 to -20°C).

When studying the processes of salt transfer in frozen hydrate saturated sediment under gas pressure, hydrate – bearing frozen samples, after being brought into contact with salt ice, were again placed in a pressure cell. Then the gas pressure of methane increased to values below (from 0.1 to

2.5 MPa) or above equilibrium (from 2.5 to 4 MPa) for a given temperature condition (-4 or -6 °C). Then the installation was also placed in a thermal box (Figure 19).




Figure 19. Scheme of the installation for the interaction of frozen hydrate-saturated and non-hydrate-saturated sediments with salt solutions under pressure.

For comparison, experiments were also carried out with frozen samples that did not contain methane hydrates. At the same time, in experiments with frozen non-hydrate samples under gas pressure, nitrogen was used to prevent hydrate formation processes.

At certain intervals of time from the start of the interaction, one of the hydrate-saturated sample twins was removed from the contact with salt solution at certain intervals of time from the start of the interaction. In experiments under pressure > 0.1 MPa, for this, the gas pressure was first released in the pressure chamber, after which it was opened, while under the atmospheric conditions this additional step was not required. For each sample, the number of parameters was determined that characterize the content of salts, moisture, and gas hydrate (through the hydrate coefficient) along the length separately in 5-8-mm-thick layers of the samples. Two probes were taken from each interval: for water/salt content and hydrate content (with 2 repetitions). In parallel with frozen

hydrate-containing samples, frozen non-hydrate-saturated samples were taken from the contact with salt solutions under the same experimental conditions, and their analysis was made.

Such interval analyses of the samples allows to follow the dynamics of salinization, dissociation of pore gas hydrates, and thawing in frozen sediment samples under various thermobaric conditions, and to study different factors that influence the salt transfer process.

Gas content was assessed by evaluating the gas volume, which was released (with 2-3 times duplicating) during thawing of sediment probes in sodium chloride solution. As a result, the released gas displaces a certain volume of water. A graduated and pre-calibrated glass tube was used to measure changes in water volumes. The latter was placed in a vessel filled with warm water. This tube was filled with water using a rubber bulb, after which the valve was closed and the initial level was recorded in the log. Then a probe of the studied hydrate containing sediment was placed under the expanded part of the tube, and as a result of gas release during the thawing of the hydrate-containing sediment, the water level in the tube changed.

For estimation of hydrate content through hydrate coefficient gained gas content values were used [327,333,334].

Experimental modeling was aimed to study of dynamics of salt migration in frozen hydrate saturated sediments, and the influence of different factors on that process:

- 1) thermobaric conditions – temperature from -2.5 to -20 °C, pressure from 0.1 to 4 MPa
- 2) the concentration of the salt solution – from 0.1 to 0.4 N
- 3) composition of the contact solution – NaCl, KCl, MgCl₂, Na₂SO₄, CaCl₂
- 4) sediment composition – model fine sand of 0.1-0.2 mm fraction, model fine sand 0.2-0.5 mm fraction, natural fine sand from South Tambey gas-condensate field, and natural sandy loam from the Arctic shelf, Buor-Khaya

5.4. Method of the temperature field change investigation during the contact of frozen hydrate saturated sediment with salt solution

Method of study of the temperature field change during interaction of frozen hydrate saturated sediments with salt solution consists of the following steps:

- 1) preparation of the sand sediments with defined water saturation in a special plastic container and freezing it;
- 2) artificial hydrate saturation of frozen sediments in the pressure cell under high pressure and negative temperatures (-4...-6±1 °C);
- 3) the pressure drop in the pressure cell, extraction of the hydrate saturated frozen sediment samples at negative temperature, and atmospheric pressure;
- 4) measuring initial parameters of the frozen hydrate - bearing samples;
- 5) inserting the temperature sensor into hydrate-saturated sand samples and bringing the samples into contact with the salt solution.

Experimental modeling of the temperature field change during interaction of frozen hydrate saturated sediments with a salt solution at the atmospheric pressure and negative temperature (-6.5°C) was made using special temperature registration Kriolab Tbf system designed at the KrioLab LTD (Russia) equipped with the respective built-in software (Figure 20). It consists of a special thermal box for maintaining negative temperatures, thermal sensors (8 pieces, d=2.5 mm), and an automatic recording system, making temperature records with defined time intervals. In the frame of this work time interval of 10 seconds was chosen for a more precise register of the fast temperature drop during hydrate saturated sediments interaction with the salt solution. The temperature sensors were calibrated to a precision of ±0.05 °C alongside a special reference sensor in a liquid thermostat. The temperature in the freezing cell was programmed to a certain value (-6.5 °C) and maintained during whole experiment. Finally, the sensor through the **transmission chain** and branch-box was connected

to the PC, and the software was set up for continuous acquisition. The measurements in a temperature field change were made whilst the frozen hydrate-saturated sediments interacted with the salt solution. The measurements were finished when the recorded temperature returned to its initial value and remained stabilized at that value.




Figure 20. Scheme of Kriolab Tbf system for temperature field change investigation (a) and picture of the frozen hydrate saturated sand sample with temperature sensors (b). 1 - frozen hydrate-saturated sediment; 2 - solution container; 3 - NaCl solution; 4 - temperature sensors; 5 - thermistor streamer; 6 - branch box; 7 - fan; 8 - USB cable; 9 - PC with Kriolab Tbf software; 10 - cooling chamber. Modified after [309].

Model fine sand-2 was used for this investigation as an object of research. The studied sediment was saturated with distilled water up to 12%, then placed in special plastic perforated tubes of 5 cm diameter and around 8 cm length, and frozen. Prepared frozen samples were placed in the pressure cell (up to 2 samples per one experiment), sealed and depressurized, and methane (CH_4 99.9% mole fraction) was added to the pressure of hydrate formation (>3 MPa). The model frozen sand was artificially saturated with hydrates for about 2 - 4 weeks. A more detailed description of the hydrate saturation process is given in chapter 5.1. After hydrate saturation, the frozen hydrate-saturated sand was transferred in metastable conditions by decreasing pressure below equilibrium at the negative

temperature (-6.5°C). Due to the self-preservation effect in frozen sediments hydrate can be preserved in the porous media for a long time without sufficient change in hydrate content.

Then the main initial parameters of hydrate-saturated sand samples were measured (initial water content, density, hydrate coefficient). After that, 4 holes of 2.5mm diameter were drilled in each sample, up to a depth of 3cm in the direction of the center. The distance between temperature sensors is around 1.5 cm. Thus, sensors are located from the contact zone up to the top of the sample deep in the frozen hydrate saturated sediment, and register the temperature of the sample whilst the hydrate is dissociating as a result of salt migration.

After placing the sensors, frozen hydrate-saturated frozen sand samples were brought into contact with NaCl salt solution at atmospheric pressure and a fixed negative temperature (-6.5°C). In parallel with hydrate-containing frozen samples, frozen non-hydrate-saturated samples were brought into contact with salt solutions under the same experimental conditions ($P=0.1\text{MPa}$, $t=-6.5^\circ\text{C}$).

Temperature registration was carried out from the beginning of the experiment on the interaction of a frozen sample with a salt solution to the stabilization of temperature conditions in the sample. The average duration of the experiment was about 5-9 hours.

To compare the results against a baseline, a sample made of non-hydrate-saturated sediment sand was investigated under the same conditions and in the same way ($P=0.1\text{MPa}$, $t=-6.5^\circ\text{C}$, $C_{\text{sol}}=0.1\text{N}$).

In addition to the study of the dynamic of the temperature field change in frozen hydrate saturated and frozen non-hydrate saturated sediment as a result of salt migration, different contact solution concentration was tested (from 0.1 to 1 N).

5.5. Method for assessing the salt content in sediment samples

Over the course of the **experimental modelling**, samples of saline sand were obtained after the interaction of frozen hydrate-saturated sediment with saline solutions. It was necessary to check the salt content to assess dependences of salt migration in frozen hydrate-bearing sediments. Nowadays, there is a problem of a rather high cost of chemical analysis of sediment samples for the content of ions of water-soluble salts, among which one of the most applicable are atomic adsorption and flame photometric methods. In addition, these methods use aqueous extracts of salts, which requires long-term sample preparation, and dilution of concentrations is used, which can lead to inaccurate results and additional errors. In this regard, within the framework of this work, a new method was proposed for determining the concentration of salts in a certain range of moisture and concentrations through the activity of pore moisture.

Thus, the measurement of the content of Na^+ cations in the course of experiments during the interaction of frozen hydrate-containing sand with NaCl salt solution was carried out by the method of water extracts using a flame photometer, a conductometer, as well as by a specially developed method for determining the concentration of salts through the activity of pore moisture (potentiometric method).

The samples on which moisture content was determined were subsequently (before drying, right after the experiment) used for the interval determination of the salt ions content that migrated from the contact saline solution.

One way the analysis of ion concentration was carried out was using the method of water extraction of water-soluble salts using a MARK 603 conductometer. The measurement error of the conductometer was checked on standard solutions of NaCl salts in the concentration range from 0.02% to 0.1%, which corresponds to the range of NaCl concentration in the studied water extracts.

The accuracy of determining the concentration of Na^+ ions using a conductometer in the samples under study was $0.1 \pm 0.05 \text{ mg-EQ/100 g}$.

Another method, a newly proposed one, was based on the measurement of the water potential (activity). Pore moisture activity is the ratio of the vapor pressure of water over a given material to the vapor pressure of pure water at the same temperature. First of all, it was found that the moisture content in a certain moisture range does not affect the activity of pore moisture (Figure 21).




Figure 21. The water activity of non-saline sand at the various water content.

Therefore, any change in the pore moisture activity value in a given moisture range will be determined solely by the salt content in the studied sand samples. To further test this theory, and to identify empirical dependencies, sand samples with a given salinity (% of salts to a dry sample of sand) were prepared and their activity was studied depending on moisture content. Based on the measurement results, graphs of the distribution of pore moisture activity against salinity for several values of moisture (10, 12, 15, and 20%) (Figure 22a), as well as graph of the concentration of salts in the pore moisture against activity was plotted (Figure 22b).




Figure 22. Distribution of pore moisture activity from sand salinity (a) and salt concentration in pore moisture (b).

It should be noted that when recalculating the salt concentration from salinity to concentration per pore moisture, all obtained values of the pore moisture activity gave a linear dependence. The results showed a high correlation between the activity of pore moisture and the concentration of salts in sand samples. The correlation coefficient was 0.9963.

In addition, the dependence of activity on the concentration of salts in the solution (outside the pore space) was tested using the **HydraFlash program**. This software allows ~~allows~~ the determination

of the activity for salt solutions but is unable to determine the concentration of salts through activity. Therefore, this dependence was obtained through data analysis "salt concentration → activity" using the program Excel (Figure 23).




Figure 23. The water activity from NaCl concentration in water solution (HydraFlash).

A linear dependence of moisture activity on the concentration of salts in the solution was also obtained.

To finally confirm the theory and establish the limitations and accuracy of the method, the salt concentration in sand samples was assessed after the contact of frozen hydrate-saturated sediment with saline by two methods through water extracts: conductometric and flame photometric, followed by the comparison of the results with each other. The results showed a high dependence (correlation coefficient 0.9984) between the methods (Figure 24), which makes it possible to compare the proposed potentiometric method with one of the methods of aqueous salt extracts.




Figure 24. Comparison of the measurement results of the Na⁺ concentration by a flame photometer and a conductometer.

To analyze the accuracy and limitations of the potentiometric method, we compared the results of measurements of the activity of pore moisture on sand samples with a given salinity and samples of saline sand after experimental modeling with data from a conductometer, which made it possible to identify an empirical equation for calculating the salt concentration (mg-EQ/100 g) through activity (Figure 25).




Figure 25. Pore moisture activity versus Na⁺ ion concentration.

Thus, the procedure for calculating salt concentration through water activity consists of the following steps:

- 1) Obtaining the parameters of the activity of pore moisture of soil samples on the WP4C device
- 2) Calculation of salt concentration in pore moisture through empirical dependence (C_w , %):

$$C_w = -2209.5 * a + 2209.7$$

where a – water activity, u.f.

- 3) Calculation of the percentage concentration of salt in relation to a dry sample of sand (C_s , %):

$$m_{salt} = C_w * m_{moisture} = C_w * (m_{wet\ sed} - m_{dry\ sed})$$

$$C_s[\%] = \frac{m_{salt}}{m_{dry\ sed}} = \frac{C_w * (m_{wet\ sed} - m_{dry\ sed})}{m_{dry\ sed}}$$

- 4) Conversion to standard concentration units – mg-EQ / 100 g by sodium ion:

$$C_s\left[\frac{mg * EQ}{100g}\right] = \frac{C_s[\%]}{0,023}$$

The proposed method has several advantages:

- 1) No need for special sample preparation. The WP4C can measure pore moisture activity even at low sample weights. Only 1-2 g of sample is sufficient for measurements;
- 2) Relatively high measurement accuracy: +- 0.5 mg-EQ/100 g (+-0.1%);
- 3) Direct measurement method without dilution. Due to the absence of additional manipulations with the sand sample, the measurement error is reduced solely to the instrumental error of the device, which is + -0.0004;
- 4) Cheapness. The instrument for determining the activity is much cheaper than the equipment used in atomic absorption analysis or flame photometry.

The method has a number of limitations:

- 1) Applicable in the concentration range from 0.02 to 5% in terms of pore moisture;
- 2) Method accuracy $\pm 0.1\%$;
- 3) Mainly applicable to sandy sediments.

5.6. Processing of the experimental data

For all samples during processing, initial characteristics (before the start of the experiment) and residual characteristics for the selected intervals (after the experiment) were obtained.

Processing of experimental data related to the study of the kinetics of gas release from frozen hydrate-containing sediment samples when interacting with salt solutions was carried out in the following order:

The moisture content (W) of the samples was determined by the gravimetric method [328]:

$W = ((m_{\text{wet sed}} - m_{\text{dry sed}}) / m_{\text{wet sed}}) \cdot 100\%$, (%.), where $m_{\text{wet sed}}$ – mass of the wet thawed sediment after the experiment, $m_{\text{dry sed}}$ – dry sediment mass.

Density (ρ) determined by the method of geometric measurement:

$\rho = m / V$, (g/cm^3), where m – mass of the frozen hydrate saturated sediment sample, V – volume of the sediment sample.

Calculation of the density of the sediment skeleton (ρ_d) was made according to the formula:

$\rho_d = \rho / (1 + W)$, (g/cm^3), where ρ – sediment density, W – gravimetric water content.

Defining volume water content (W_{vol}):

$W_{\text{vol}} = \rho_d \cdot W$, (%), where ρ_d – density of sediment skeleton, W – gravimetric water content.

Calculation of porosity (n):

$n = (\rho_s - \rho_d) / \rho_s$, (u.f.), where ρ_s – density of solid particles of sediment, ρ_d – density of sediment skeleton.

Then, the initial specific gas content (g) of the hydrate-containing sample and the residual specific gas content, both calculated by the formula:

$$g = (V_2 - V_1) * T / m_{\text{sam}} \text{ (cm}^3\text{/g)},$$

where $(V_2 - V_1)$ – change of the water volume in the glass gas collecting tube (cm^3); T – temperature correction; m_{sam} – mass of the sediment sample (g).

Based on the obtained values of specific gas content, the proportion of gas hydrate inclusions located in the pore space of frozen samples was quantitatively determined - weight hydrate content (H):

$$H = (m_H / m_{\text{sam}}) * 100\%, \text{ (\%)}$$

The degree of filling the pores with ice (S_i) was calculated by the formula:

$$S_i = W_{\text{vol}} / (0.92 \cdot n), \text{ (u.f.)},$$

where W_{vol} – volume water content, n – porosity.

Based on the data obtained, the initial and residual hydrate coefficient (K_h) was calculated for the intervals - the proportion of pore water that passed into hydrate from the total amount of water in the sample.

$$K_h = W_h / W, \text{ (u.f.)},$$

where W_h – the amount of moisture that has passed into the hydrate (% in relation to the mass of the dry sample), W – weight water content of the sample (%).

Volumetric hydrate content (H_v) was defined by the formula:

$$H_v = H \cdot \rho / \rho_H, \text{ (\%)},$$

where ρ – sediment density (g/cm^3), ρ_H – hydrate density (g/cm^3) and methane hydrate density equal 0.9 g/cm^3 .

Knowing the volumetric hydrate content, it is possible to determine the degree of pore filling with hydrate – hydrate saturation:

$$S_h = H_v / n, (\%)$$

Volumetric ice saturation (I_v) was calculated by the formula:

$$I_v = (W - W_h) / \rho_{\text{и}} * 100\%, (\%),$$

where $\rho_{\text{и}}$ – ice density (g/cm^3), equals $0.9 \text{ g}/\text{cm}^3$.

To calculate the parameters of the hydrate content of sediment, the coordination number for methane hydrate was used 5.9 [335].

The change in the flow of Na^+ ions through the cross section of a frozen hydrate-containing sample was also calculated (J , $\text{mol}/\text{m}^2 \cdot \text{sec}$):

$$J = \frac{v}{S * t}$$

where S – sample cross-sectional area, m^2 ; t is the salt migration time, sec; $v = m/23$, m is the mass of sodium ions, g; 23 – atomic mass of sodium, g/mol.

The result of processing the experimental data obtained are graphs of changes in the concentration of salt ions along the height of the sample (C , $\text{mg} * \text{EQ} / 100 \text{ g}$) and the hydrate coefficient (K_h , u.f.). Also, considering the results of temperature measurements, graphs of changes in the temperature field over time were plotted.

Chapter 6. Mechanism and dynamics of salt transfer processes in hydrate-bearing sediments

Experimental modeling of the dynamics of hydrate saturated frozen sediments with the salt solution was carried out on the fine sand-1. The prepared sediment samples had an initial moisture content of 12%. After hydrate saturation, the frozen samples were removed from the pressure cell at

a low negative temperature ($-6 \pm 1^\circ\text{C}$) under the atmospheric pressure. Due to the self-preservation effect, the frozen hydrate saturated samples were well preserved during the whole experiment (2-3 days). The fraction of pore moisture converted to hydrate (K_h) was 0.6.

Typical characteristics of frozen samples containing conserved gas hydrates are presented in Table 5.

Table 5. Main initial characteristics of frozen hydrate-containing sand samples.

Sediment	W,%	Density, g/cm ³	Porosity, u.f.	S _i ,%	K _h
Fine sand-1	12	1.69	0.39	47	0.6

Then the frozen hydrate-containing samples at a negative temperature ($-6 \pm 1^\circ\text{C}$) were brought into contact with a 0.2N NaCl solution previously cooled to the experimental temperature.

At the end of the experiment, for the samples interacting with the salt solution, layer-by-layer determination of the parameters of water content, gas, and salt content was carried out.

6.1. Influence of hydrate presence on the salt migration in frozen sediments

Experimental modeling of the interaction of hydrate saturated frozen sediment with the cooled (frozen) salt solution shows that active diffusion of salt ions into the sand sample occurs, which resulted in salinization of the frozen sample and dissociation of methane hydrate in porous media [31].

To identify the effect of the hydrate component of frozen sediments on the process of the salt migration occurring during the interaction of saline solution with hydrate-containing samples was compared to the salt migration involving samples, where the hydrate was absent, with the salt solution under equilibrium pressure (2.5 MPa) and non-equilibrium pressures (0.1 MPa) at the constant negative temperature ($-6 \pm 1^\circ\text{C}$). It was found that the migration of salt ions into a frozen

hydrate-containing sample occurs more intensively than in a sample without hydrates. This dependence is observed both at non – equilibrium pressure under conditions of hydrate self-preservation (0.1 MPa) and at pressures above equilibrium (2.5 MPa) (Figure 26).

As a result, at atmospheric pressure, Na^+ ions penetrated deep into the sample from the contact zone to 5.3 cm in a frozen hydrate-containing sample, whilst the penetration of salts did not exceed 2.6 cm in the frozen non-hydrate-containing sample after 4 hours from the start of interaction with 0.2 N NaCl solution (Figure 26). At the same time, the total accumulation of salts in a frozen sample that did not contain hydrates was significantly lower than in a sample saturated with hydrate. An experiment at a pressure above equilibrium (2.5 MPa) demonstrates a similar dependence but differs in a lower intensity of salt accumulation (Figure 26). Thus, Na^+ ions penetrated to a depth of 3.6 cm in a frozen hydrate-containing sample, while in a frozen non-hydrate-containing sample only by 2.2 cm after 4 hours from the onset of interaction.




Figure 26. Effect of gas pressure on the accumulation of salt ions (Na^+) in frozen sand (fine sand -1) samples ($W=12\%$) containing (solid line) and not containing (dashed line) gas hydrates 4 hours after interaction with 0.2 N NaCl solution ($t=-6\text{ }^\circ\text{C}$).

Thus, it can be observed, that gas pressure has a strong influence on the salt migration in hydrate saturated frozen sediments, while in frozen non-hydrate saturated sediments pressure barely has an effect on salt accumulation almost does not affect salt accumulation.

For the studied samples, the average Na⁺ flux (J, mol/cm²·sec) was calculated for different gas pressure conditions and fixed temperature (-6 °C) (Table 6). Results of calculation demonstrate that salt flow is less in non-hydrate samples and barely depends on pressure (equals 1.2-1.7·10⁻¹⁰ mol/cm²). While the flow in hydrate saturated samples under atmospheric pressure is more than twice higher and reaches 5·10⁻¹⁰ mol/cm²·sec. However, at the stable pressure (2.5 MPa) average Na⁺ flux is just slightly higher than flow in the frozen non-hydrate bearing sediments and equals 1.9·10⁻¹⁰ mol/cm² at a pressure 2.5 MPa.

Table 6. Average Na⁺ flux in frozen sand (fine sand-1) under the equilibrium (2.5 MPa) and non-equilibrium (0.1 MPa) pressure (4 hours of interaction with 0.2N NaCl solution at -6 °C).

Pressure	Sample	Average Na ⁺ flux J·10 ⁻¹⁰ , mol/cm ² ·sec
0.1 MPa	non-hydrate	1.2
	hydrate	5
2.5 MPa	non-hydrate	1.7
	hydrate	1.9

The specificity of salt migration in frozen hydrate-containing sediments is due to the presence of not only the process of ice melting but also the dissociation of pore gas hydrate with the formation of a liquid phase of water and the release of large volumes of methane. When gas is released, the structure of gas hydrate formations also loosens, and formed moisture becomes gas-saturated. The greater the content of the liquid phase – the more active the salt migration, therefore, the higher the

diffusion coefficient. In addition, the absorption of ions through gas-saturated pore water also promotes higher salt flow to pass through hydrate saturated frozen sediment.

Therefore, salt migration is more active in hydrate - containing sediments at atmospheric pressure. Moreover, if salt migration in frozen non-hydrate-containing sediments does not depend on the pressure, in the case of frozen hydrate containing sediment there is a much higher flow of salts at the non-equilibrium pressure. When pressure is changed from equilibrium to non-equilibrium the average density flow rises more than 2 times.

6.2. The interaction of frozen hydrate saturated frozen sediments with salt solutions in time

Interaction of hydrate saturated sediments with NaCl (concentration 0.1N) was studied during different time period (3.9, 17.5, 25.5 and 28.8 hours after the start of interaction) at a constant negative temperature ($-6\pm 1^\circ\text{C}$) and pressure (0.1 MPa). It was observed that there is a direct dependence between salts accumulation and hydrate dissociation. So, when salts migrate from the zone of contact, hydrate start to dissociate in the same direction – from the zone of contact.

Figure 27 shows the accumulation of Na^+ ions (A) and the amount of pore water in hydrate form (hydrate coefficient) (B) in frozen samples containing preserved (metastable) hydrate at atmospheric pressure after contact with a frozen NaCl solution with a concentration of 0.1N at different times. After 4 hours, the salts penetrated to a depth of 2.3 cm, and the maximum concentration increased from 0.1 to 0.7 mg-EQ/100 g in the contact zone. After 29 hours, the accumulation of salts was recorded at a distance of 5 cm from the contact area with maximum value of concentration 1.6 mg-EQ/100 g.




Figure 27. Accumulation of Na⁺ ions (A) and changes in the hydrate coefficient (B) in samples of frozen hydrate-containing sand-1 ($K_h^{\text{in}} = 0.6$, $W=12\%$) at different times after interaction with NaCl solution 0.1 N ($P=0.1$ MPa, $t=-6$ °C).

Joint analyses of the salt distribution (Figure 27A) and hydrate coefficient (K_h) by the height of the sample (Figure 27B) shows, that the salinity value determines the residual content of methane hydrate in porous media. In the initial state (before the contact with the solution) for sand samples (fine sand-1), about 60% of the pore moisture was in the hydrate form initially. In the process of **unilateral salinization**, there was a decrease in the proportion of pore moisture in the hydrate form (K_h), and the appearance of a front of complete hydrate decomposition (Figure 27B). This way, after 4 hours of interaction hydrate completely decomposed within 1 cm from the contact with salt solution, while after 29 hours no residual hydrate was observed within 4 cm.

In this case, it is possible to distinguish a certain critical content of salt ions in sand samples, which causes complete gas hydrate decomposition in the sediments (C_{cr} , mg-EQ/100 g). Under given conditions (temperature -6°C and concentration of NaCl contact solution 0.1 N) in sediment samples (fine sand 1), the critical value of salt accumulation was about 0.7-0.8 mg-EQ/100 g.

During the process of salt migration in a frozen hydrate-containing sample, a thawing front can also occur in addition to the decomposition front of pore hydrates, when the accumulated salt ions completely transfer the pore ice into water. In Figure 28 the movement of two fronts is shown: hydrate decomposition and thawing boundary.




Figure 28. Experimental evaluation of the decomposition front of porous gas hydrates and the thawing front in frozen hydrate-containing sand samples (sand-1, $W = 12\%$) when interacting with 0.1 N NaCl salt solution at a temperature -6°C and pressure 0.1MPa.

The pore hydrate decomposition front is ahead of the thawing front since thawing requires a higher value of salt accumulation in the frozen sample. This way, after 4 hours of interaction hydrate dissociation reached 1 cm from the contact with salt solution, while the thawing boundary was not registered at all. Only after 17.5 hours thawing front reached 1 cm, while the hydrate dissociation front at that time was already 2 cm from the contact with 0.1 NaCl solution at -6°C and pressure 0.1

MPa. Moreover, with time the difference between fronts became bigger. That is, the critical concentration of pore hydrate dissociation is lower than the critical concentration of thawing and is about 0.7 mg-EQ/100 g and 1.5 mg-EQ/100 g, respectively. Thus, the thawing critical concentration is 2 times higher than hydrate dissociation C_{cr} in the porous media of the studied model sand.

The thaw front in the sample is well fixed by the characteristic color distribution of the sample (Figure 29). Additionally, it was evaluated using a special probe-needle.



Figure 29. The appearance of frozen sand samples (fine sand 4, W=12%) before (A) and after interaction (B) with 0.1 NaCl solution at -6°C .

In addition, the average Na^+ flux was calculated in time based on the data of salt ions concentration in the frozen hydrate saturated sediments. The Na^+ flux in the first 5 hours is $2 \cdot 10^{-10}$ mol/cm²·s, with time the flux decreases, and after 30 hours it decreases to values of about $0.2 \cdot 10^{-10}$ mol/cm²·s, so 10 times less than at the beginning of the interaction.




Figure 30. Average Na⁺ flux change in time during interaction of frozen hydrate saturated sediment (fine sand-1) with NaCl solution of 0.1N at -6 °C and 0.1 MPa.

The presented experimental data on the interaction of frozen hydrate-containing sediments with salt solutions showed that the processes of salt ions migration in frozen hydrate saturated sediment can occur more **intensively** compared to frozen sediments that do not contain hydrates. That is apparently associated with the processes of dissociation of pore hydrate and the appearance of a **liquid phase of** water. Accumulation of salts causes active hydrate dissociation in porous media, so the concentration of ions defines residual hydrate content and it is possible to define the critical concentration of complete hydrate dissociation (C_{cr}). Salt migration contributes to the formation of two fronts: the gas hydrate dissociation front and the thawing front, while the dissociation front is ahead of the thawing front of a frozen hydrate-containing sample.

6.3. Temperature changes as a result of salt migration in frozen hydrate-saturated sediments

As shown earlier, when frozen hydrate-saturated sediments interact with a cooled (frozen) saline solution, active diffusion of salt ions occurs into these samples, resulting in salinization of the frozen

sample. When a certain critical value of salt concentration in a frozen hydrate-containing sample was reached, the hydrate dissociation front was formed.

The process of salinization of a frozen hydrate-containing sample, as well as a frozen non-hydrate-saturated sample, is associated with a temperature decrease since the methane hydrate dissociation and the ice melting under the influence of salts is an endothermic process. Given that the enthalpy of hydrate dissociation is 1.5 times higher than the amount of heat required to melt ice, a more significant decrease in temperature can be expected in a hydrate-containing sample compared to non-hydrate-containing one during their interaction with a salt solution. Experimental data on the dynamics of temperature change over time close to the contact area (2.5 cm from contact with NaCl 0.4N saline solution) in frozen sand samples (fine sand-2), saturated and not saturated with hydrates, confirm this (Figure 31).

In these experiments, samples (fine sand-2) were used, which had an initial moisture content of 12%, and porosity of the order of 38-40%. At the same time, the saturation of the pore space with ice in a non-hydrate-containing frozen sample was 50%, and in a hydrate-containing one with a total degree of pore filling of 53%, the saturation with hydrate was 35% (Table 7).

Table 7. Frozen non-hydrate saturated (sample 1) and hydrate saturated (sample 2) fine sand -2 initial characteristics.

Sample	Water content, %	Density, g/cm ³	Porosity, %	Degree of pore filling with ice, %	Degree of pore filling with hydrate, %
1	12	1.8	40	50	0
2	12	1.85	38	18	35

As follows from Figure 31, a decrease in temperature in the samples (fine sand-2) is recorded almost immediately after being brought into contact with a saline solution. However, the temperature

reaches its minimum value at a distance of 2.5 cm from the contact area in a frozen sample after 0.7 h, and in a hydrate-containing sample only after 1 h. At the same time, the value of temperature decrease in the hydrate-containing sample is 0.6°C and only 0.2 °C in the frozen one. The rate of temperature decrease after bringing a frozen non-hydrate-containing sample into contact with a saline solution was 0.4 °C/h; for hydrate samples, 0.6 °C/h.

After the temperature decrease in the samples, its gradual increase to the initial value of -6.5°C is observed, which is achieved in the hydrate-containing sand sample after 5 hours, and in the frozen one - after 3.7 hours.




Figure 31. Dynamics of temperature change in time in the near-contact area (2.5 cm from the contact) in frozen sand samples (fine sand-2) interacting with 0.4 N saline solution at atmospheric pressure: 1 – non-hydrate-containing frozen sample (dotted line); 2 - hydrate-containing (solid line); 3 - initial temperature (tin) -6.5 °C (dashed line).

The temperature distribution along the length of a frozen sample containing and not containing pore hydrates at a fixed point in time (0.6 h after the start of contact with the salt solution) can be traced in Figure 32. As follows from the obtained data, at the time point of 0.6 hours after the start of the interaction of fine sand-2 with 0.4N NaCl solution at an initial temperature of -6.5°C, a regular

decrease in the temperature deviation from the initial value is observed with distance from the contact area. This deviation in frozen non-hydrate bearing sand is much less than in frozen hydrate-bearing sample. For a sample containing hydrates, a temperature deviation is registered along the entire length of the hydrate saturated frozen sample in contrast to a non-hydrate-containing sample, where temperature decrease was observed only in the area closest to the contact with salt solution (fine sand-2).




Figure 32. Temperature distribution along the length of a frozen sample (fine sand-2), not containing (1) and containing gas hydrate (2) after 0.6 h of interaction with 0.4 N. NaCl solution at atmospheric pressure. The dotted line - 3 shows the initial temperature $t_{in} = -6.5^{\circ}\text{C}$.

Figure 33 demonstrates the temperature dynamics of a frozen hydrate-containing sample (fine sand-2) overtime at different distances from the area of contact with 1N NaCl solution. In the initial state, both the frozen hydrate-containing sample and the solution had a temperature of -6.5°C . After bringing them into contact, a regular decrease in temperature in the sample was observed. This was fixed at different distances from the contact area. The temperature decrease in the sample relative to the initial value (-6.5°C) reached 1.9°C . At the same time, at different distances from the contact area, this decrease shifted in time. So, at a distance of 2.5 cm from the contact (value for 1 sensor),

the minimum temperature (-8.3 °C) was reached 20 minutes after contact with saline solution, and at a distance of 7 cm, this temperature decrease was recorded after 35 minutes. At the same time, there was some delay in the start of sample cooling depending on the distance from the contact area, while the general cooling rate at different intervals remained almost the same (0.08-0.1 degrees per minute). After reaching the minimum temperature value in each interval, a gradual increase in temperature to the initial values (-6.5°C) was observed. Moreover, after 3 hours after the start of the interaction, the difference in temperature between the intervals was practically not observed, although later in the sample a gradual increase in temperature continued.




Figure 33. Dynamics of temperature change in time in a frozen hydrate-containing sample ($S_h^{in}=38\%$, $S_i^{in}=15\%$) on the distance 2.5; 3.5; 5.5 and 7.0 cm from the contact zone with 1 N NaCl solution ($t_{in}=-6.5\text{ }^\circ\text{C}$, $P=0.1\text{ MPa}$).

The temperature distribution along the height of a frozen hydrate-containing sample (fine sand-2) at various times after the interaction of a frozen hydrate-containing sample with NaCl (1 N) salt solution at the cooling stage can be seen on Figure 34. Prior to contact with the saline solution, the temperature of the sample practically did not change and was equal to -6.5°C. After the frozen sample was brought into contact with saline solution, a decrease in temperature was observed within a minute

at a distance of 2.5 cm from the contact with the saline solution ($\Delta t=1\text{ }^{\circ}\text{C}$). At the same time, at a distance of 3.5 cm or more, the temperature remained the same (-6.5°C). After 4 min, a decrease in temperature was recorded to a depth of 5.5 cm from the contact area, and after 8 min, a change in temperature was observed throughout the whole sample. At the same time, at a depth of 2.5 cm, the temperature reduced by 1.8°C , and the temperature gradient along the length of the sample reached $0.4\text{ }^{\circ}\text{C}/\text{cm}$. Subsequently, the temperature in the sample was equalized (fine sand-2) by stabilizing the temperature near the contact and lowering the temperature away from the contact area. As a result, after 30 min, the temperature distribution in the observed region (from 2.5 to 7 cm) was almost linear. The temperature of the sample varied within $-8.2\text{...}-8.4\text{ }^{\circ}\text{C}$. Then, a gradual increase in the temperature of the sample was observed, primarily from the side of the area adjacent to the contact with the salt solution. So, one hour after the interaction of a frozen hydrate-saturated sample (fine sand-2) with a saline solution, the temperature at a depth of 2.5 cm increased by $0.3\text{ }^{\circ}\text{C}$ from the minimum value (at a time point of 30 min), and at a depth of 7 cm it practically did not change (Figure 34).

Thus, when a cooled salt solution interacts with a frozen hydrate-containing sample (fine sand-2), a cold wave arises due to the dissociation of pore gas hydrate caused by salt migration. As a result, a decrease in temperature along the length of the sample (by $1.8\text{ }^{\circ}\text{C}$) can be observed. After completion of the hydrate dissociation process, a gradual increase in temperature takes place, which begins primarily in the interval of the sample adjacent to the area of contact with the salt solution.




Figure 34. Change in the temperature distribution along the length of the frozen hydrate-saturated sample fine sand-2 ($S_h^{in}=41\%$, $S_i^{in}=12\%$) at the stage of cooling in the first hour of interaction with 1 N NaCl solution ($t_{in}=-6.5$ °C, $P=0.1$ MPa).

The intensity of the temperature decrease of a frozen hydrate-saturated sample (fine sand-2) upon contact with a saline solution strongly depends on the concentration of the contact solution. In the course of the experiments, the influence of various NaCl solution concentrations (0.2, 0.4, 0.6, 1 N) on the temperature change of frozen hydrate saturated fine sand-2 dynamics was studied. Table 8 shows the initial characteristics of the studied samples, which were used to assess the effect of contact solutions. The saturation of the pore space of frozen samples with gas hydrate ranged from 35 to 41% and saturation with ice from 11 to 18%.

Table 8. Initial characteristics of the frozen hydrate-saturated fine sand-2 samples before the contact with NaCl solution of different concentrations.

Solution, N	Sample	Water content, %	Density, g/cm ³	Porosity, %	Ice saturation, %	Hydrate saturation, %
0.2	1	12	1.8	40	12	41
0.4	2	12	1.85	38	18	35

0.6	3	12	1.79	39	11	40
1	4	12	1.83	38	15	38

A plot of experimental data illustrating the temperature change of frozen hydrate-saturated sand in time when it comes into the contact with a NaCl solution of various concentrations are shown in Figure 35.




Figure 35. Dynamics of temperature changes in time of frozen hydrate-containing samples (fine sand-2) in the near-contact area (2.5 cm from the contact) interacting with NaCl solution of various concentrations (0.2-1N) ($t_{in} = -6.5^{\circ}\text{C}$, $P = 0.1 \text{ MPa}$).

As follows from the obtained data, the magnitude of the decrease in temperature in the studied samples (fine sand-2) depends on the concentration of the contact solution. With an increase in the concentration of the contact solution from 0.1 to 1 N, the temperature decrease in the near-contact region (2.5 cm) changed from 0.3°C to 1.9°C . In addition, the time to reach the minimum temperature was reduced from 1 hour to 15 minutes (Figure 35).

A more significant decrease in the temperature of a frozen hydrate-containing sample (fine sand-2) with an increase in the concentration of the interacting solution is associated with an increase in the

rate of gas hydrate dissociation in the pore space of sediments. Dependence of the maximum decrease in temperature during the interaction of a frozen hydrate-containing sample (fine sand-2) with a saline solution (Δt) on the concentration of the contact solution ($t_{in}=-6.5^{\circ}\text{C}$, $p=0.1\text{MPa}$) is presented on Figure 36.




Figure 36. Influence of the contact solution concentration on the maximum temperature drop ($\Delta t=t_{in}-t_{min}$) in frozen hydrate saturated fine sand-2 (C_{sol} from 0.1 to 1 N, $t_{in}=-6.5^{\circ}\text{C}$, $P=0.1\text{MPa}$).

At a concentration of NaCl contact solution of 0.1 N, the decrease in temperature was only 0.1 $^{\circ}\text{C}$. In addition, with an increase in the concentration of the solution to 1 N, the decrease in the temperature reached 2 $^{\circ}\text{C}$. Thus, with an increase in the concentration of the contact solution from 0.1 to 1 N, the decrease in the temperature of a frozen hydrate-saturated sand sample increases by a factor of 20.

It should be noted that with an increase in the concentration of the contact solution, there is a significant increase in the rate of temperature decrease ($V_t, ^{\circ}\text{C}/\text{hour}$) (Figure 37). Thus, at a contact solution concentration of 0.1 N, the rate of temperature decrease in a frozen hydrate-containing sample (fine sand-2) caused by salinity was 0.4 $^{\circ}\text{C}/\text{h}$, and at a contact solution concentration of 1 N it increased to 4.7 $^{\circ}\text{C}/\text{h}$, which is more than 10 times higher than in contact with 0.1 N solution.




Figure 37. Influence of contact solution concentration on the rate of temperature decrease (V_t) of frozen hydrate-saturated fine sand-2 (C_{sol} from 0.1 to 1 N, $t_{in} = -6.5^\circ C$, $P = 0.1$ MPa).

The time of the sample temperature recovery to its original value (equals to the surrounding temperature $-6.5^\circ C$) after the initial period of cooling, caused by the interaction of hydrated frozen fine sand-2 with saline solution, also depends on the concentration of the contact solution (Figure 38).




Figure 38. Influence of concentration on the time taken for the temperature of the sample to return to the initial value (T_r) after the cooling period caused by the interaction of hydrate-containing frozen sand with saline solution ($t_{in} = -6.5^\circ C$, $P = 0.1$ MPa).

Thus, it takes about 5 hours for the sample temperature to return to its original state value (-6.5 °C) when a frozen hydrate-saturated sample (fine sand-2) interacts with a saline solution with a concentration of 0.2 N, and about 12 hours are required at a concentration of 1N.

Thus, based on experimental studies, it can be inferred that when samples containing ice and pore hydrate interact with a salt solution, a change in the temperature field in the sample was observed, caused by the processes of pore hydrate dissociation and ice melting. The amount of cooling of the sample due to the dissociation of the pore gas hydrate significantly exceeds the effect of cooling from the melting of the pore ice interacting with the salt solution. At the same time, the concentration of the contact solution has a great influence on the intensity of temperature changes.

5.4. Mechanisms of salt migration in frozen hydrate containing sediments

The experimental data make it possible to represent the dynamics of the interaction of frozen hydrate-containing sediments with saline solution in the following way. When a hydrate-containing sample is brought into contact with a saline solution, salt ions penetrate deep into the sample, which causes melting of pore ice and dissociation of pore hydrate. As a result, a porous hydrate dissociation front appears, which expands from the area of contact with the salt solution. At the same time as the dissociation front is expanding, a thawing front forms at the same time with a more significant accumulation of salt ions.

As the pressure increases, the stability of the gas hydrate component increases because a higher concentration of salts in the pore space is needed to cause the pore hydrates to dissociate in accordance with the phase diagram (Figure 49).

In this regard, with an increase in gas pressure in the saline solution-sample system, the flow of salt ions decreases, as well as the amount of liquid water in the sample, which is formed due to

the dissociation of pore hydrate, decreases. In **frozen rocks** that do not contain gas hydrates, gas pressure in the studied pressure range (up to 4.0 MPa) practically does not have a significant effect on the phase composition of **moisture**, which is not noticeably reflected in the accumulation of salts in frozen sandy sediments in the absence of moisture migration.

Analysis of experimental ~~modeling~~ data on the interaction of frozen hydrate-saturated sediments with NaCl salt solution allows us to describe phase changes in frozen hydrate-containing sand in equilibrium and non-equilibrium (with self-preservation) pressure and negative temperature (-6°C) as follows (Figure 39). At gas pressure values below equilibrium (<2.4 MPa), the pore hydrate in frozen sand exists in a metastable state due to the manifestation of the self-preservation effect (Figure 39a). **Residual gas hydrate accumulations are located under the ice shell, which was formed during the crystallization of water released during the partial dissociation of the pore gas hydrate.** Migration and accumulation of salt ions in frozen hydrate-saturated sediment when interacting with a salt solution occurs primarily along films of **unfrozen water on the surface of sand particles**, as well as along the boundaries of ice crystals. This is accompanied by an increase in the thickness of unfrozen water films due to melting of pore ice. **With the disappearance of the ice film on the surface of the hydrate**, its active dissociation begins with the release of gas and water, which, under conditions of negative temperature and the endothermic effect of dissociation, partially freezes with the formation of ice (Figure 39a).




Figure 39. Scheme of phase-transformation of frozen hydrate saturated sediments during salt migration: (a) metastable conditions (experimental pressure $P_{exp} < P_{eq}$); (b) stable conditions ($P_{exp} \geq P_{eq}$).

At an initial pressure above equilibrium (2.4 MPa and above), the pore methane hydrate in non-saline frozen sand is in stable conditions (Figure 39b). The equilibrium content of the liquid phase of water under these conditions should be less than in the sample containing preserved gas hydrate. Considering that the process of salt ions migration occurs primarily through films of unfrozen water in a hydrate-containing frozen sample, at pressures above equilibrium the migration of ions occurs more slowly than at gas pressure below equilibrium. In addition, it can be assumed that the intergranular permeability of the gas hydrate is much lower than the permeability of ice. Under these conditions, taking into account a significant shift in pressure (ΔP) relative to the equilibrium value, when the salt solution interacts with a frozen hydrate-containing sediment sample, higher critical salt ion concentrations are required to destabilize the pore hydrate, which are achieved over a longer time. When critical concentrations are exceeded, gas hydrate dissociation is observed. However, unlike metastable conditions, this process is less active and the process of formation of pore ice is

slower. The gas formed during the dissociation of the pore hydrate can later concentrate in the void space with an incomplete degree of filling of the pores, and also be in the form of separate gas inclusions in the ice that was formed during the dissociation of the hydrate. In this case, part of the gas can dissolve in unfrozen water, but this will be limited by the concentration of salts in the liquid phase (Figure 39b).

Phase transitions that occur during the salts migration and accumulation in ice- and hydrate-bearing sediments are accompanied by temperature effects associated with the melting of pore ice and the dissociation of pore gas hydrate. Given that these processes are endothermic, a decrease in the temperature of saline sediment samples is observed. The effect of temperature decrease is most pronounced in frozen hydrate-bearing sediment, since the enthalpy of dissociation of pore gas hydrate is 1.5 times higher than during ice melting.

A decrease in the temperature of the frozen sediment when interacting with a salt solution is associated with a shift of the thermodynamic equilibrium that exists in frozen sediment at a fixed negative temperature. As a result of salt transfer in frozen sediment, the content of salt ions in films of unfrozen water increases. This disturbs the existing phase equilibrium and causes the melting of pore ice, which is accompanied by a decrease in temperature. As the salts continue to accumulate, additional melting of pore ice and a further decrease in temperature occur. This decrease will continue until an equilibrium occurs between the sediment temperature, pore solution concentration and ice. This can be clearly seen in the phase diagram for the NaCl solution (Figure 15), where at a given concentration of the salt solution, there is an equilibrium temperature corresponding to the freezing point of this solution. So, the change of temperature stops, when there is an equilibrium between the solution and ice at a fixed negative temperature. Based on this, the maximum decrease in temperature will correspond to the equilibrium temperature formed during the interaction of the salt solution with the pore ice of the concentration of the salt solution.

Taking into account that in the process of salt migration, the salinization front in the sediment sample shifts, the area of temperature decrease will also shift, which is clearly seen from the data obtained (Figure 33). The maximum decrease in temperature, which is fixed in the experiments, corresponds to the maximum rate of ice melting and hydrate dissociation. Subsequently, when the melting of the pore ice and the dissociation of the gas hydrate dies out, or when the pore ice turns into liquid water and the gas hydrate inclusions completely decompose, a gradual increase in the sample temperature to ambient temperature is observed. In hydrate-containing samples, this process is longer than in frozen ones, which is due to deeper cooling and the formation of additional ice during the dissociation of porous gas hydrate.

The temperature effect in frozen hydrate-containing sediment caused by salt transfer can be schematically represented as follows (Figure 40). Initially, before bringing the sediment into contact with the salt solution, a uniform temperature field (Figure 40a) is observed in the sample at a fixed negative ambient temperature. The temperature of the frozen hydrate-containing sample corresponds to the ambient temperature. After bringing the sample into contact with the salt solution under isothermal conditions, the temperature decreases in the near-contact region of the sample due to the migration and accumulation of salt ions, which causes the melting of ice and the dissociation of the pore gas hydrate (Figure 40b). With the accumulation of salt ions, the process of dissociation of the pore gas hydrate increases, which contributes to the continuation of the process of lowering the temperature (Figure 40c). Subsequently, the process of temperature decreases in this interval decays and shifts into the depth of the sample, where the salinization front moves (Figure 40d). Depending on the intensity of salinity, which is determined by the concentration of the contact solution, partial or complete decomposition of gas hydrates is possible. In the latter case, along with the salinization front, a porous gas hydrate dissociation front appears and even a porous ice thaw front, if there is a significant accumulation of salts.




Figure 40. Scheme of temperature changes during the interaction of a frozen hydrate-containing sample with saline solution.

The displacement of the cooling wave to the end of the sample opposite from the contact during salt transfer is replaced by a gradual increase in the temperature of the sample to the ambient temperature. That is due to the damping of phase transitions along the sample and the establishment of a new thermodynamic equilibrium between the pore ice and the salt solution (Figure 40f).

A more detailed scheme of changes occurring in frozen samples containing a metastable hydrate in the contact with saline solution can be seen in Figure 41.




Figure 41. Scheme of temperature changes in a frozen hydrate-containing sample in the process of salt transfer, phase transformations, and changes in ice and hydrate saturation along the height of the investigated sample.

At some point in time after the start of the interaction with the salt solution, a cooling zone is clearly visible in the sample, caused by the dissociation of the pore gas hydrate and the melting of ice due to the migration and accumulation of salt ions. An analysis of phase transformations in the pore space of sediment at a given point in time shows that in the region far from the contact, to where salt ions have not yet penetrated, the frozen hydrate-saturated sample has an initial temperature equal to the ambient temperature and initial values of ice and hydrate content. Under these conditions, the hydrate continues to exist in a **conserved** state without phase transitions (Figure 41a).

In the middle part of the sample, where the minimum temperature values are fixed, there is a dissociation front of the porous gas hydrate, which causes a local decrease in temperature (Figure 41b). In this region, a sharp decrease in hydrate saturation is observed due to the decomposition of pore hydrate. The water formed during the dissociation of the pore gas hydrate reduces the concentration of salts in the liquid phase. This, occurring under lower temperatures, contributes to its freezing, as a result of which an increased ice content is observed in the near-contact zone compared to the initial value, despite an increase in salt accumulation and an increase in the content of unfrozen water (Figure 41c). As a result, the temperature of the sample in the contact zone rises to the initial values, both due to the release of heat during freezing of water, and due to the temperature equalizing with the ambient temperature. In addition, in the near-contact area, when approaching the contact solution, ice saturation can decrease due to the high accumulation of salts, and, accordingly, the high content of unfrozen water.

Thus, salt migrates in frozen hydrate saturated sediments more actively than in frozen non-hydrate saturated frozen sediment and with greater temperature effect. That is due to a higher amount of potential paths for salt ions migration (mainly crystals of ice and unfrozen water). And the lower the pressure, the more active the hydrate dissociation due to higher unfrozen content.

Salt migration and accumulation has high influence on the process of hydrate decomposition, and decrease in temperature corresponds to the process of hydrate dissociation in the porous space. the temperature reaching its minimum values indicates a decrease in the rate of decomposition of the gas hydrate and the beginning of the attenuation of this process. Then the temperature of the cooled zone of the sample begins to rise gradually and comes into equilibrium with the ambient temperature. This occurs both due to heat exchange with the environment, and due to the heat released during freezing of the moisture formed during the dissociation of the hydrate.

Chapter 7. Influence of the different factors on the process of salt migration in frozen hydrate saturated sediments

During experimental modelling of the interactions of saline solution with methane hydrate-saturated frozen sediments the following factors were investigated: pressure, temperature, contact solution concentration and composition, and dispersity of the sediment.

7.1. Salt and mass transfer in sediment saturated with gas hydrate, depending on pressure

Earlier in Chapter 5.1, an analysis of the influence of pressure on interactions between saline solution and both hydrate-bearing and non-hydrate-bearing frozen sediments was made (Figure 26). And it was shown that the process of salt migration in frozen hydrate-containing samples strongly depends on thermobaric conditions, which determine gas hydrate stability in frozen sediments.

Figure 42 shows the thermobaric conditions for the existence of methane hydrates in **free volume**, obtained by a computer simulation using the HydraFlash program. In this case, points (T1-T6) show the P-T conditions of the aforementioned experiments investigating the interaction of frozen fine sand-2 containing methane hydrates with saline solution. As a part of the investigation, a series of 6 experiments were devised, which were carried at a constant negative temperature of -6°C and at pressures ranging from atmospheric (0.1MPa) to 6.0 MPa. **Two series of experiments were** carried out at a pressure below equilibrium (under self-preservation conditions) (T1 - 0.1 MPa, T2 - 1.8 MPa), and the other three were carried out above equilibrium conditions (T4 - 3.5 MPa, T5 - 4.0 MPa, T6 – 6.0 MPa). In addition, P-T conditions of one series of experiment were close to the hydrate stability limit (T3 – 2.5 MPa at $P_{\text{eq}}=2.4$ MPa).




Figure 42. Modeling P-T conditions of methane hydrate stability in the HydraFlash program and thermobaric conditions for conducting experiments on the migration of salt ions in frozen hydrate-saturated sediments (T1-T6).

A comparison of experimental data on the interaction of frozen hydrate-bearing sediments with salt solutions at a fixed negative temperature (-6°C) and pressures ranging from 0.1 MPa to 4.0 MPa demonstrates that with increasing pressure, the intensity of accumulation of salt ions (Na⁺) decreases. A particularly significant decrease in the accumulation of salt ions was observed at gas pressures above equilibrium. The decrease in the intensity of accumulation of salt ions with increasing pressure is reflected in the increase in the pore hydrate preservation at a fixed time of interaction.

Thus, in a frozen hydrate-containing sample (fine sand-2) 4 hours after interaction with a 0.1 N NaCl solution, Na⁺ ions penetrated 5.2 cm deep into the sample at a pressure of 0.1 MPa, and at a pressure of 4.0 MPa the penetration depth was only 3.4 cm. At the same time, the salt content in the contact zone decreased from 22 mg-EQ/100 g to 15 mg-EQ/100 g (Figure 43).




Figure 43. The effect of pressure on the accumulation of salt ions (Na⁺) (a) and the decrease in the hydrate coefficient (K_h) (b) in samples of frozen hydrate saturated fine sand-2 ($K_h^{in} \sim 0.3$, $W=12\%$) at non-equilibrium and equilibrium pressure 4 hours after the start of interaction with a frozen NaCl solution 0.1 N ($t = -6^\circ\text{C}$).

The amount of salt ions accumulated in the studied samples determines the hydrate content. In the initial state (before the contact with the NaCl solution) for fine sand-2 samples, about 25% of the pore moisture was in the hydrate form. During one-sided salinization of the sample, a regular decrease in hydrate content was observed from the side of the contact zone. The hydrate coefficient decreases

when moving from a pressure above equilibrium (4.0 MPa) to non-equilibrium (0.1 MPa), while the areas where there is a sharp decrease in hydrate content expand. So, 4 hours after the start of the interaction of hydrate-containing samples with a solution of 0.1 NaCl, the area of complete decomposition of the gas hydrate due to salinization changed from 1.8 cm at a pressure of 4.0 MPa to 3.4 cm at a pressure of 0.1 MPa (Figure 43b). Thus, it is possible to trace the presence of a gas hydrate decomposition front in the porous media of a frozen sample, the position of which depends on the pressure at fixed values of temperature and concentration of the contact salt solution.

The calculation of the average migration flow of salt through the frozen hydrate saturated sample of sand for the described experiments shows that its value sharply decreases from $38 \cdot 10^{-10}$ to $0.5 \cdot 10^{-10}$ mol/(cm²·s) with an increase in pressure from 0.1 MPa to 6.0 MPa (Figure 45).




Figure 44. Influence of gas pressure on the flux density of Na⁺ ions through the cross section of a frozen hydrate-containing sample 4 hours after the start of interaction with a frozen salt solution 0.1 N NaCl (t = -6°C).

Analysis of the salt accumulation in the studied samples and the change in hydrate content in the process of salt transfer made it possible to reveal the dependence of the critical concentration of salts in the sample (C_{cr}), which causes the complete decomposition of pore gas hydrates, on gas pressure

at a fixed negative temperature (-6°C) and the concentration of the contact solution (0.1 N) (Figure 46).




Figure 45. Dependence of the critical concentration (C_{cr}) causing decomposition of pore hydrate in a frozen sand sample on gas pressure ($t = -6^{\circ}\text{C}$, $C_{sol} = 0.1 \text{ N}$).

Studies show that the value of the critical concentration (C_{cr}) increases almost linearly ($R=0.9672$) with increasing gas pressure at a fixed negative temperature. When the pressure rises from 0.1 to 6.0 MPa, the value of (C_{cr}) at -6°C increases nearly a factor of 7. Thus, at atmospheric pressure, the critical concentration of salts causing complete gas hydrate dissociation is 0.2% NaCl, and at 6.0 MPa it reaches 1.3% NaCl.

The empirical dependence of the critical salt concentration on pressure (Figure 46) obtained on the basis of experimental studies made it possible to calculate, for the interval values of the salt ion concentration in the sample, the critical pressure values below which complete decomposition of the pore gas hydrate occurs.

The critical pressure values were compared with the gas pressure specified in the experiment. As a result, a salinization interval was identified along the length of the sample, where the calculated critical pressures corresponding to the values of salt accumulation exceed the gas pressures specified

in the experiment. At a critical pressure below the experimental pressure, a residual content of pore hydrate is observed. **In this case**, the lower the concentration of salts, and, accordingly, the lower the critical pressure, the greater the residual content of hydrate. This made it possible to explain the nature of the distribution of salts and pore hydrate along the length of the sample and to find their relationship with the value of the gas pressure specified in the experiment. This pattern can be traced in Figure 46, which shows the distribution of sodium ion concentration and hydrate content along the height of a frozen hydrate-containing sample after interaction with NaCl salt solution at a fixed negative temperature (-6°C) and a given gas pressure (2.5 MPa).




Figure 46. Altitude distribution of a frozen hydrate-containing sample: (A) salt concentrations and hydration coefficient; (B) – critical pressure of dissociation of methane hydrate (concentration of contacting NaCl solution 0.1 N, $t = -6^{\circ}\text{C}$).

So, at the distance of 4 cm from the point of contact with a saline solution, no hydrate was recorded, but at the distances greater than 4cm, a residual hydrate content was observed. Figure 11b shows the critical pressure of pore gas hydrate decomposition calculated from the salt concentration for each interval of a frozen hydrate-containing sample and the given gas pressure of the experiment, as well as the equilibrium pressure characteristic of the initial non-saline sample. At distances up to 4 cm from the contact, the critical pressure of hydrate dissociation was higher than the given gas pressure, so the hydrate in this region completely dissociated. Thus, in few centimeters of the sample (from the contact area), as a result of the accumulation of salts, the critical pressure was 2.68 MPa, which is 0.18 MPa higher than the specified gas pressure in the experiment (2.5 MPa). And at a distance of more than 4 cm, the given gas pressure exceeds the critical pressure and therefore residual hydrate content is observed. Thus, in the range from 4.5 to 5 cm, the critical pressure is 2.45 MPa, which is 0.05 lower than the experimental pressure.

Also, thermodynamic modeling was carried out using software “HydraFlash” software to assess the equilibrium values of salt concentration for methane hydrate for given thermobaric conditions. According to HydraFlash calculations, the equilibrium values of salt concentrations for the considered pressures turned out to be higher than the critical concentrations obtained in the experiments. So, at a temperature of -6°C and a pressure of 4.0 MPa, the equilibrium concentration for methane hydrate is 15%, and at 2.5 MPa it is 0.8% (1% and 0.6% according to the calculations). Such a difference between them is primarily due to the fact that in calculations of the equilibrium concentration of salts, its value is considered in relation to the volume of hydrate. And in the experiment, there was a local non-equilibrium accumulation of salt ions at the boundary with hydrate particles.

So, during the experiments, it was revealed that the intensity of the accumulation of salt ions in frozen sandy sediments containing gas hydrates is higher than in frozen sands that do not contain

gas hydrate formations. In contrast to frozen sediments, in hydrate-containing sediment media, pressure has a huge impact on the processes of salt transfer. It was experimentally found that in frozen hydrate-containing sediment samples, when the pressure decreases from values above equilibrium (>2.4 MPa) to pressures below equilibrium (up to 0.1 MPa), the flow of salts into the sample increases significantly. When the pressure drops from 4.0 MPa to 0.1 MPa at a fixed negative temperature (-6°C), the density of salt flow into a frozen hydrate-containing sample increases by a factor of 40 .

7.2. Temperature influence on the process of hydrate-bearing sediment interaction with saline solutions

As is well known, from the thermobaric point of view an increase in temperature shifts the conditions for the stability of gas hydrates towards higher pressures. The study of the temperature influence on the dissociation of pore gas hydrate was considered on the model fine sand-2, frozen and saturated with methane hydrate. Samples in the initial state had a uniform distribution of methane hydrate ($K_h=0.4$) in porous media, and water content equal to 12% .

The temperature influence on the processes of salt transfer in frozen hydrate-containing sandy sediments was considered in the range from -3°C to -20°C . During the experiments, it was found that with an increase in the ambient temperature, salinization processes are intensified, which accelerates the hydrate decomposition in the porous space of frozen hydrate-containing sediments. At a higher negative temperature (-3°C), during the experiment of 2 hours, NaCl ions, unlike other temperature conditions (-6°C and -20°C), penetrate almost the entire depth (8 cm) of a frozen hydrate-containing sample of fine sand-2 (Figure 47). At the same time, in a sample with a lower temperature (-6°C),

salt ions migrated to a depth of up to 6 cm from contact, and at -20°C only up to 2.2 cm from contact with a salt solution ($C_{\text{sol}}=0.2\text{N NaCl}$).




Figure 47. Effect of temperature on the accumulation of salt ions (Na^+) in frozen hydrate-containing samples (fine sand-2) after 2 hours of interaction with 0.2 N NaCl solution.

As a result of migration and accumulation of salts in frozen hydrate-saturated samples (fine sand-2), interacting for 2 hours with a salt solution (0.2 N NaCl), the pore methane hydrate completely decomposed to a depth of 7 cm from contact at a temperature of -3°C . In addition, the residual hydrate content ($K_h=0.22$ in comparison with the initial value equals $K_h^{\text{in}}=0.4$) was observed only in the top part of the sample. In the experiment at an ambient temperature of -6°C , the porous

gas hydrate completely decomposed in the region of 3.5 cm from the contact. Although at a lower negative temperature (-20°C), partial decomposition of the hydrate was recorded only in the near-contact region (1 cm from the contact), where the hydrate coefficient decreased from the initial 0.37 to 0.1 (Figure 48).




Figure 48. The effect of temperature (-3, -6, -20°C) on the decrease in the hydration coefficient (K_h) in samples of frozen hydrate-containing fine sand-2 ($K_h^{in} \sim 0.4$, $W=12\%$) 2 hours after the start of interaction with frozen 0.2 N NaCl salt solution.

The calculation of the average migration flow of salt for the described experiments shows that its value decreases with decreasing temperature (Figure 49). So, at -2°C the salt flux was $70 \cdot 10^{-10}$ mol/cm²·s, at -6°C the Na⁺ flux decreased more than 3 times to $20 \cdot 10^{-10}$ mol/cm²·s, and at -20°C Na⁺ flux did not exceed $4.2 \cdot 10^{-10}$ mol/cm²·s (Figure 49).




Figure 49. Effect of temperature on the average Na⁺ flux into a frozen hydrate-containing sample (fine sand-2) for 4 hours of interaction with a frozen salt solution of 0.1N NaCl (P=0.1MPa).

When study of the temperature influence on salt migration in hydrate saturated frozen sediment (fine sand-2) at the pressure above equilibrium (4 MPa), the same general dependence was obtained: decrease of temperature lowering the Na⁺ flux. However, the values of average salt flow is lower at the gas pressure 4 MPa, than at the atmospheric pressure. Thus, Na⁺ flux was equal to $20 \cdot 10^{-10}$ mol/cm²·s at gas pressure 4 MPa and temperature -2°C , what is more than 3 times lower, than at the atmospheric pressure. And at the pressure 4 MPa and -6°C Na⁺ flux was equal to $2 \cdot 10^{-10}$ mol/cm²·s – by factor 10 lower, than at the pressure 0.1 MPa (Figure 49).

The analysis of salt accumulation in the studied samples and the change in hydrate content in the process of salt transfer made it possible to reveal the dependence of the critical salt concentration in the sample (C_{cr}), which causes complete decomposition of metastable (self-preserved) porous gas

hydrates (at $P=0.1$ and 4 MPa), on temperature during the interaction of frozen hydrate-containing sediments (fine sand-2) with NaCl solution ($0.1N$) (Figure 50).




Figure 50. Effect of temperature on the critical concentration (C_{cr}) causing decomposition of pore hydrate in a frozen sand sample (fine sand-2, $P=0.1$ and 4 MPa, $C_{sol}=0.1N$).

Experimental data show that the critical concentration (C_{cr}) increases with decreasing ambient temperature (Figure 50). So, at atmospheric pressure and a temperature of $-2^{\circ}C$, the critical concentration (C_{cr}) was 0.6% NaCl for pore moisture, at a temperature of $-10^{\circ}C$ - 2.5% , and at $-16^{\circ}C$ - it reached 3.4% , which is actually 8 times exceeds the critical concentration at $-2^{\circ}C$.

With an increase in pressure and the transition of a hydrate-containing system to an equilibrium state, the value of the critical concentration at a fixed negative temperature increases. But general dependence of the critical concentration increase with temperature decrease was observed as well. Hence, critical concentration increases from 0.8 to 1.8% NaCl in pore moisture at gas pressure 4 MPa with temperature decrease from -2 to $-6^{\circ}C$, while at 0.1 MPa C_{cr} change in this temperature range **0.6 to 1.8%.**

The temperature dependence of the critical concentration of hydrate dissociation in frozen **deposits is linear.** The higher the negative temperature, the lower the concentration of salt ions of the

pore solution, which causes complete decomposition of the gas hydrate in the porous media of frozen deposits. According to the established empirical dependence, it is possible to obtain the critical concentration of the complete decomposition of the pore gas hydrate through temperature value.

Such obtained dependence can be explained the following way. Under conditions of increasing ambient temperature, the process of migration of salts from the contact solution into the frozen hydrate-containing sample increases, which leads to faster dissociation of the porous gas hydrate due to an increase in the accumulation of salt ions and a decrease in the critical concentration required for the complete decomposition of the porous gas hydrate, as in conditions of initially equilibrium pressure, and under conditions of self-preservation.

Thus, the influence of temperature on the intensity of mass transfer in frozen hydrate-containing sediments and the intensity of hydrate dissociation during interaction with salt solutions have been experimentally obtained. It is noted that with a decrease in temperature, the preservation of methane hydrate in porous media of frozen sediment increases, due to the slowing down of the processes of migration of salt ions. The critical concentration of Na^+ ions in frozen hydrate-containing sediment samples increases by a factor of 8 with a decrease in negative temperature from -3 to -16°C . Finally, the average Na^+ flux decreases exponentially (by a factor 10) when temperature decreases from -3 to -20°C .

7.3. Influence of salt solution concentration on the intensity of salt migration and hydrate dissociation in hydrate saturated sediments

Before the start of the experimental modeling on the interaction of frozen hydrate saturated sediments with salt solution of different concentration, computer simulation of thermobaric conditions for the stability of methane hydrate in salt solution system were calculated depending on the concentration of NaCl in “HydraFlash” software (Figure 51).




Figure 51. Thermobaric conditions of hydrate stability in the water-NaCl solution of different concentration simulated in “HydraFlash” software.

The results of thermodynamic modeling showed that the presence of salts shifts the stability conditions of methane hydrate towards higher pressures and lower temperatures from a thermodynamic point of view. This shift depends on the concentration of salts. So, at a fixed pressure of 2.5 MPa, with an increase in NaCl concentration from 1 to 5%, the equilibrium temperature decreases from -5 to -10°C.

It can be assumed that for metastable hydrates in the pore space of frozen sediment, the concentration of salts (NaCl) will also have a significant impact on their preservation at a fixed negative temperature.

As part of the experimental simulation, two series of experiments were carried out. The first series was aimed at studying the influence of the concentration of the contacting NaCl solution on the processes of dissociation of pore gas hydrate in frozen sediment (fine sand-2) under non-equilibrium (metastable) conditions ($P=0.1$ MPa), when the dissociation of pore gas hydrate is determined mainly by the kinetics of salt transfer. The second series of experiments is devoted to

studying the influence of the concentration of the contacting NaCl solution on the processes of hydrate dissociation under stable conditions ($P=4$ MPa), when the dissociation of pore gas hydrate is determined by thermodynamic and kinetic factors.

An increase in the concentration of a salt solution in contact with frozen hydrate-saturated fine sand-2 at a fixed negative temperature (-6°C) and atmospheric pressure (i.e. below equilibrium) leads to more intensive migration and accumulation of salt ions in the sample. Thus, at a contact solution concentration of 0.1N at a fixed interaction time (4 hours), salt ions (Na^+) migrated to a depth of 4.5 cm from the contact into hydrate-saturated frozen sand (Figure 52). With an increase in the concentration of the contact solution to 0.2N, the salinization zone of the frozen hydrated sample increased to 5.2 cm (Figure 52), and at a concentration of 0.4N, salt ions penetrated almost the entire depth of the sample (7 cm) (Figure 52).




Figure 52. Effect of contact solution concentration on the accumulation of salt ions (Na⁺) in frozen hydrate-containing samples (fine sand-2) after 4 hours of interaction with NaCl solution (t=-6°C, P=0.1 MPa).

The amount of salt ions accumulation in the studied sand samples determines the distribution of hydrate over the sample. In the initial state (before contact with the solution), about 40% of the pore moisture in the sand samples (fine sand-2) was in the hydrate form. In the course of one-sided salinization of the sample, a regular decrease in hydrate content was observed from the side of the contact zone. With an increase in the concentration of the NaCl contact solution from 0.1 to 0.4N at a fixed interaction time (4 hours) and a temperature (-6°C), the area of a sharp decrease in the hydrate coefficient expands significantly.

Thus, with an increase in the concentration of the contact solution from 0.1 to 0.4 N, the area of complete decomposition of the gas hydrate due to salinity changed from 2.5 cm to 5.6 cm (Figure 53). At the same time, a decrease in the hydrate coefficient (K_h) was also observed in the interval of the sample (fine sand-2), where residual hydrate was fixed.

Thus, it is possible to trace the shift of the decomposition front of porous gas hydrate in frozen samples depending on the concentration of the contact salt solution.




Figure 53. Influence of NaCl contact solution concentration on the distribution of hydrate coefficient (K_h) along the length of frozen hydrate-containing samples (fine sand-2) ($K_h^{in} \sim 0.4$, $W=12\%$) after 4 hours of interaction ($t=-6^\circ\text{C}$, $P=0.1$ MPa).

In the experiment with a frozen hydrate-containing sample (fine sand-2) interacting with a NaCl solution at a pressure above equilibrium (4.0 MPa), similar dependence was obtained, which indicate a more active accumulation of salt ions and dissociation of pore methane hydrate with an increase in the concentration of the contact NaCl solution. At this pressure, after 4 hours salt ions (Na^+) migrated into hydrate-saturated frozen fine sand-2 to a depth of 3.4 cm from contact with a 0.1N NaCl solution. And with an increase in the concentration of the solution to 0.2N, salt ions were fixed at a depth of 6 cm (Figure 54).




Figure 54. Influence of the contact solution concentration on the accumulation of salt ions (Na^+) in frozen hydrate-containing samples (fine sand-2) interacting for 4 hours under a pressure of 4.0 MPa with a NaCl solution ($t=-6^{\circ}\text{C}$).

An increase in the intensity of salt ions migration and accumulation in frozen hydrate-saturated sediments interacting with a NaCl solution at a pressure above equilibrium (4.0 MPa) with an increase in the concentration of the contact solution was also accompanied by a more active hydrate decomposition. Thus, in the initial state (before contact with saline solution) in frozen hydrate-containing samples, about 40% of pore moisture was in the gas hydrate form. After 4 hours of the contact of the frozen hydrate-containing sample (fine sand-2) with saline solution at -6°C , the pore hydrate completely decomposed at a distance of 1.6 cm from the contact area with 0.1N NaCl solution, and at a distance of 4.4 cm from the contact area with 0.2N NaCl solution (Figure 55).




Figure 55. Effect of contact solution concentration on distribution of hydrate coefficient (K_h) along the height of frozen hydrate-containing sample (fine sand-2, $K_h^{\text{in}} \sim 0.4$, $W=12\%$) after 4 hours of the interaction with frozen NaCl salt solution ($t=-6^\circ\text{C}$, $P=4.0$ MPa).

With an increase in the concentration of the contact solution, the average Na^+ flux into the frozen hydrate-containing sample increases. Thus, when frozen hydrate-saturated fine sand-2 interacted with a 0.1N NaCl solution at atmospheric pressure, the Na^+ flux was $1 \cdot 10^{-10}$ mol/cm²·s, and with an increase in concentration to 0.4 N, the Na^+ flux increased to $8.2 \cdot 10^{-10}$ mol/cm²·sec. A similar dependence was also observed in experiments at a pressure above equilibrium (4.0 MPa): with an increase in the concentration of the NaCl contact solution from 0.1 to 0.4 N, the average Na^+ flux in them increased from 0.4 to $4.6 \cdot 10^{-10}$ mol/cm²·s, however these values were lower than the salt fluxes in frozen hydrate-saturated fine sand-2 under atmospheric pressure conditions (Figure 56).





Figure 56. Influence of contact solution concentration on the average Na⁺ flux in frozen hydrate-saturated sample (fine sand-2) after 4 hours of the interaction ($t = -6^{\circ}\text{C}$, $P = 0.1 \text{ MPa}$ and 4.0 MPa).

An analysis of the experimental modeling data makes it possible to describe phase changes in frozen hydrate-containing sediment in the process of salt migration at equilibrium and non-equilibrium (with self-preservation) gas pressure under the influence of different concentration (Table 9). In the initial state before the start of the experiment, the concentration of salts in the pore solution of the sample is close to zero. At the same time, under conditions of a fixed negative temperature (-6°C) and a pressure above equilibrium (4 MPa), the pore methane hydrate in the initially non-saline frozen sand was in stable conditions (Table 9a).

After contact with a saline solution at a temperature of -6°C in a frozen sample, an increase in the concentration of salts in the pore space occurs as a result of their migration and accumulation. **The migration of salt ions from the contact solution occurs primarily along the films of unfrozen water on the surface of sediment particles,** as well as along the boundaries of ice crystals. This is accompanied by an increase in the thickness of unfrozen water films due to partial dissociation of the hydrate and melting of pore ice (Table 9c).

Table 9. Scheme of phase-transformation in frozen hydrate-saturated sediments during increase of salt solution concentration at stable conditions (experimental pressure $P_{exp} \geq P_{eq}$) and metastable conditions ($P_{exp} < P_{eq}$).



Upon reaching a certain critical salt concentration ($C > C_{cr\ hydr}$) there is a sharp increase in the rate of hydrate dissociation, which leads to its complete decomposition. The moisture formed in this case under conditions of negative temperature and the endothermic effect of dissociation is supercooled and then frozen (Table 9e).

A further increase in salt concentration to values above the critical concentration of pore ice melting ($C > C_{cr \text{ ice}}$), which can be observed when a frozen hydrate-containing sample comes into the contact with a high-concentration salt solution (non-freezing at a given negative temperature), will be accompanied by complete thawing of the sample (Table 9g).

With a decrease in gas pressure below the equilibrium, the pore hydrate in frozen sediment can pass in a metastable state due to the manifestation of the self-preservation effect (Table 9b). Under these conditions, the residual gas hydrate inclusions in frozen sediment will be under the ice shell, which was formed due to the crystallization of water released during the partial dissociation of the pore gas hydrate at a negative temperature.

Under these conditions, the interaction of a salt solution with frozen sediments containing porous gas hydrate under metastable conditions is also accompanied by the migration and accumulation of salt ions. However, this process is more active, and the critical concentration causing the decomposition of pore methane hydrate will be lower than when interacting under stable conditions ($P > P_{eq}$) (Table 9d).

Studies [214] show that the equilibrium content of the liquid phase of water at a pressure above equilibrium ($P > P_{eq}$) is less than in a hydrate-containing sample in a metastable state ($P < P_{eq}$). Taking into account that the migration of salt ions occurs mainly through films of unfrozen water, the process of salt ions migration in frozen sediments containing metastable gas hydrate will occur more intensively than at pressures above equilibrium.

In addition, the role of intergranular (along crystal boundaries) permeability of pore hydrate and ice in the migration of salt ions in frozen hydrate-bearing sediment should be noted. From the literature [315] it is known that the intergranular (ionic) permeability of pore ice contributes significantly to salt transport in frozen sediment. It can be assumed that the intergranular permeability of the hydrate is much lower than the permeability of ice. Thus, in frozen hydrate-containing

sediment under non-equilibrium conditions, which are usually characterized by an increased ice content compared to stable conditions, the critical concentration of pore gas hydrate dissociation is reached faster (Table 9f).

Thus, the experimental data show that an increase in the concentration of the contact solution (NaCl) increases the accumulation of salt ions in frozen hydrate-bearing sediments and activates the dissociation of pore gas hydrate. This is observed both at pressures below equilibrium, when the porous gas hydrate is in the metastable conditions, and at pressures above equilibrium (4.0 MPa). However, in the latter case, the intensity of these processes decreases.

7.4. The process of interaction of hydrate-containing sediments with a salt solution of various chemical composition

Analysis of the results of thermodynamic modeling of the conditions of P-T stability of methane hydrate in the free volume for negative temperatures (from 0 to -20°C) for the most widespread salts in the permafrost (NaCl, CaCl₂, MgCl₂, KCl and Na₂SO₄) in the "HydraFlash" software, carried out before the start of the experiments. It demonstrated a significant shift in the stability conditions of methane hydrate, depending on the chemical composition of the studied salts. It was found that the value of the equilibrium conditions for hydrate stability shifts to lower temperatures and pressures in the following series of salts of 5% concentration: Na₂SO₄ - KCl - CaCl₂ - NaCl - MgCl₂. Thus, for a temperature of -5°C, the pressure shift for Na₂SO₄ was 2.8 MPa, and for MgCl₂ it was 3.8 MPa compared to a non-saline gas-water system (Figure 57).




Figure 57. Influence of a chemical composition of salt solution (Na₂SO₄, KCl, CaCl₂, NaCl, MgCl₂) with a concentration of 5% on the conditions of methane hydrate stability in the range of negative temperatures from 0 to -20°C (calculations were made in the “HydraFlash” software).

The obtained results of thermodynamic modeling generally coincide with the existing data presented by other researchers [173,182,196].

For salts of various chemical compositions used in the calculations, special experiments were carried out on their interaction with samples of frozen hydrate-saturated sand (fine sand-2). The experiments were carried out at a fixed temperature (-6°C), atmospheric pressure and salt concentration (0.1N).

Figure 58 shows the accumulation of salt ions in samples of frozen hydrate-saturated sand interacting with salt solutions of various chemical compositions (NaCl, CaCl₂, MgCl₂, KCl and Na₂SO₄). The obtained experimental data demonstrate that the intensity of migration and accumulation of salts increases in the series Na₂SO₄ – KCl – CaCl₂ – NaCl – MgCl₂ (Figure 58).




Figure 58. Accumulation of Na⁺ ions in samples of frozen hydrate saturated sand (fine sand-2, W=12%) after 4 hours of contact with 0.1N NaCl solution of different composition ($t = -6^{\circ}\text{C}$, $P = 0.1\text{MPa}$).

Thus, 4 hours after the start of the interaction, the maximum penetration of salt ions (up to 5 cm) was recorded in the sample in contact with the MgCl₂ solution. For the NaCl solution, this value did not exceed 4.6 cm, while for the CaCl₂ and KCl solutions it was 2.8 and 2 cm, respectively. The smallest depth of salt accumulations was observed in the sample interacting with the Na₂SO₄ solution (about 0.6 cm).

The distribution of the residual content of hydrates in samples of frozen hydrate-saturated sand after 4 hours of their interaction with salt solution of various chemical composition confirms the revealed pattern in the described series: $\text{Na}_2\text{SO}_4 - \text{KCl} - \text{CaCl}_2 - \text{NaCl} - \text{MgCl}_2$, which is determined by the intensity and magnitude of the salt accumulations and hydrate dissociation. In this series, there is a regular decrease in the residual content of gas hydrate in the pore space of sand (fine sand-2). So, if in the initial (before contact) state, about 40-50% of the pore moisture was in the hydrate form, then after 4 hours of interaction with the MgCl_2 solution, the pore hydrate completely dissociated in the range up to 4.5 cm from the contact. In the case of the NaCl solution hydrate dissociate up to 3.4 cm, and in the case of CaCl_2 and KCl , this interval did not exceed 2 cm and 1 cm, respectively. In the experiment with a solution of Na_2SO_4 , in contrast to the solutions described above, only a partial decrease in the hydrate coefficient in the contact zone was noted. In the range up to 0.5 cm, this decrease was twofold (from the initial 0.5 to 0.25), and the decrease in the hydrate coefficient was generally recorded only at a distance of 3 cm from the contact of the sample with the saline solution (Figure 59).




Figure 59. The influence of the chemical composition of the contact salt solution on the decrease in the hydrate coefficient (K_h) in frozen hydrate-containing samples (fine sand-2, $K_h^{\text{in}} \sim 0.5$, $W=12\%$) after 4 hours of interaction with the frozen salt solution ($t=-6.5^\circ\text{C}$, $C_{\text{sol}}=0.1\text{N}$, $P=0.1\text{MPa}$).

For each contact solution of different chemical composition, the average salt flux into a frozen hydrate-saturated fine sand-2 sample was calculated (Figure 60). The highest value of salt ions flux was obtained upon the contact of frozen hydrate-saturated fine sand-2 with MgCl₂ solution ($5.4 \cdot 10^{-100\text{g}}/\text{cm}^2 \cdot \text{sec}$). At the same time, in the series MgCl₂ - NaCl - CaCl₂ - KCl - Na₂SO₄, the salt flux decreases by almost 5 times.




Figure 60. The influence of the chemical composition of the salt contact solution on the salt flux into frozen hydrate-containing sample (fine sand-2) after 4 hours of interaction with a frozen salt solution ($t=-6^{\circ}\text{C}$, $C_{\text{sol}}=0.1\text{N}$, $P=0.1\text{MPa}$).

The critical concentration of pore hydrate decomposition was calculated for the studied salts during their interaction with frozen hydrate-saturated sand at a fixed negative temperature (-6°C) and atmospheric pressure. This concentration (C_{cr}) varied from 3.23 mg - EQ/100 g (0.08%) for MgCl_2 solution to 3.5 mg - EQ/100 g (0.16%) for Na_2SO_4 . In the described series ($\text{MgCl}_2 - \text{NaCl} - \text{CaCl}_2 - \text{KCl} - \text{Na}_2\text{SO}_4$), the critical concentration of complete hydrate dissociation increased almost 2 times (Table 10).

Table 10. Dependence of the critical concentration (C_{cr}) causing the decomposition of pore hydrate in a frozen fine sand-2 sample on the chemical composition of the contacting solution ($t=-6^{\circ}\text{C}$, $P=0.1\text{MPa}$, $C_{\text{sol}}=0.1\text{N}$).

Salt	C_{cr} , mg - EQ/100 g	C_{cr} , %
MgCl ₂	3.23	0.08
NaCl	4.89	0.11
CaCl ₂	3.29	0.13

KCl	3.90	0.15
Na ₂ SO ₄	3.50	0.16

The influence of the chemical composition of the contacting solution at a fixed concentration is primarily associated with the migration ability of salt ions. Thus, chloride salts have a greater migration capacity in frozen sediments compared to sulfates [46]. The results of experimental studies are consistent with this relationship: chloride salts, compared with sulfates, more actively penetrate into frozen hydrate-saturated sediment and cause more intense hydrate dissociation. In this case, in a series of cations, the mobility of ions is determined by their radius [97,312]. Thus, among the chloride salts used in the experiments, the cation radii increase in the series: $Mg^{2+} < Na^{+} < Ca^{2+} < K^{+}$. In the same series, a decrease in the salt ions migration intensity into frozen hydrate-containing sediments was experimentally obtained, which was reflected in a decrease in the activity of hydrate dissociation in porous media of frozen sediment.

Therefore, in the frame of the experimental modeling influence of different compositions of the contact solution on the process of salt migration in frozen hydrate saturated sediments was tested. It was found, that in the raw $MgCl_2$ - $NaCl$ - $CaCl_2$ - KCl - Na_2SO_4 accumulation of salts in frozen hydrate saturated sand decreases, and as a result, less active hydrate dissociation is observed. Less accumulation of salts can be explained by the decrease in the average density flow by 5 times, and the increase in the critical concentration by 2 times in the described raw.

7.5. Sediment dispersion influences the process of salt migration in hydrate saturated sediments

To reveal the role of the dispersity of the sediment medium and various clay minerals, special studies were carried out on the interaction of frozen hydrate-saturated sediments of various

compositions with a 0.1 N NaCl salt solution at a fixed negative temperature (-6°C) and atmospheric pressure.

Several series of experiments were performed. First of all, an assessment was made of the influence of the content of clay particles and their mineral composition on salt transfer in frozen hydrate-saturated sand samples.

Samples were prepared using quartz sand with a fraction of 0.1–0.2 mm (fine sand-2) to which kaolinite particles (up to 15%) were added. The initial characteristics of frozen hydrate-saturated samples are presented in Table 11.

Table 11. Properties of hydrate-saturated frozen sand with different amount of kaolinite.

Sample	Kaolinite, %	Water content, %	Density, g/cm ³	Porosity, %	Ice saturation, %	Hydrate saturation, %
1	0	12	1.8	39	14	50
2	5	12	1.8	38	14	50
3	7	12	1.82	37	12	49
4	10	12	1.86	35	15	52
5	15	12	1.85	34	18	51

In the course of experiments on the interaction of frozen hydrate-containing samples with saline solution (NaCl), it was found that with an increase in the content of kaolinite particles, the intensity of salt transfer decreases. So, 4 hours after the interaction of a frozen hydrate-containing sample with a NaCl 0.1N solution, the depth of penetration of Na⁺ ions into a frozen hydrate-containing sand sample decreases from 5.3 cm to 2.6 cm with an increase in the number of clay particles from 0 to 10% (Figure 61).




Figure 61. Influence of kaolinite content in fine-grained sand (fine sand-2) on the accumulation of Na⁺ ions in frozen hydrate-saturated samples 4 hours after interaction with 0.1 N NaCl solution at -6 °C.

A decrease in the intensity of salt accumulation with an increase in the amount of kaolinite particles leads to a decrease in the dissociation of pore gas hydrate. Thus, 4 hours after the contact of frozen hydrate-containing samples with 0.1N NaCl solution, the area of complete decomposition of pore hydrate decreased from 4.8 cm in sand without clay particles to 1.6 cm in a sample with 10% kaolinite (Figure 62).




Figure 62. Effect of kaolinite content in sand (fine sand-2) on hydrate dissociation in a frozen hydrate-saturated sample ($K_h^{in} \sim 0.5$, $W=12\%$) 4 hours after the start of interaction with a frozen salt solution of 0.1 N NaCl at -6 °C.

With an increase in the amount of clay particles in frozen hydrate-saturated sand in the process of interaction with a saline solution, the accumulation of moisture increases. Thus, in frozen hydrate-saturated sand without clay additives, moisture accumulation was practically not observed, ΔW in the near-contact area (about 2.5 cm) did not exceed 1.5% (Figure 63). With an increase in the amount of clay particles to 5%, an increase in water content in the near-contact area up to 7% was recorded, and in a sample of frozen hydrate-containing sand with 10% kaolinite, the increase in water content reached almost 8% (Figure 63).




Figure 63. Influence of kaolinite content in sand (fine sand-2) on moisture accumulation in a frozen hydrate-saturated sample ($W=12\%$) after 4 hours of interaction with salt solution of 0.1 N NaCl at $-6\text{ }^{\circ}\text{C}$.

Thus, an increase in the amount of clay particles in sandy hydrate-containing frozen samples led to an increase in the moisture flux from the frozen salt solution. So, with an increase in the number of kaolinite particles from 0 to 15%, the average moisture flux increased by more than 6 times (from 1.5 to $6.8 \cdot 10^{-7}$ g/sec) (Figure 64).




Figure 64. Effect of kaolinite content in sand (fine sand-2) on the average water flux through a frozen hydrate-bearing sample (W=12%) after 4 h of interaction with a frozen salt solution of 0.1 N NaCl at -6 °C.

An experimental study of the influence of clay mineralogy on the process of interaction of frozen hydrate-saturated samples with saline solution was carried out by adding 5% kaolinite or montmorillonite to quartz sand (fine sand-2). The initial characteristics of frozen hydrate-saturated samples are presented in Table 12.

Table 12. Properties of hydrate-saturated frozen sand with different composition of clay.

Sample	Type of clay	Water Content, %	Density, g/cm ³	Porosity, %	Ice Saturation, %	Hydrate Saturation, %
1	kaolinite	12	1.8	38	15	47
2	montmorillonite	12	1.81	36	18	47

An analysis of the distribution of Na⁺ ions along the height of a frozen hydrate-containing sandy sample demonstrates that less accumulation of salt ions is typical for a sediment mixture with montmorillonite than with the addition of kaolinite particles. So, after 4 hours of interaction, Na⁺ ions migrated to a depth of up to 6 cm with the addition of kaolinite particles, and with the use of montmorillonite particles only up to 4 cm (Figure 65).




Figure 65. Influence of the mineralogical composition of clay in quartz sand (fine sand-2 with 5% additives) on the accumulation of Na^+ ions in frozen hydrate-saturated samples after 4 hours of interaction with 0.1 N NaCl solution at $t=-6^\circ\text{C}$: A – kaolinite; B – montmorillonite.

As a result of the migration and accumulation of salt ions in frozen hydrate-saturated sand samples containing 5% clay particles of various mineral composition, the pore methane hydrate completely decomposed to a depth of 5 cm from the contact area with the frozen salt solution in the sample with kaolinite particles and only 2 cm in the sample with 5% montmorillonite (Figure 66).




Figure 66. Influence of the mineralogical composition of clay in quartz sand (fine sand-2) on the decrease in the hydration coefficient (K_h) in frozen hydrate-containing samples ($K_h^{\text{in}} \sim 0.4$, $W=12\%$) 4 hours after the start of interaction with frozen salt solution 0.1 N NaCl at $t=-6^\circ\text{C}$: A – kaolinite; B, montmorillonite.

In frozen hydrate-saturated sand containing montmorillonite particles, more moisture accumulation was observed compared to the sample containing kaolinite particles. Thus, in the contact

area in a sample of frozen hydrate-containing sand with montmorillonite particles, water content increased from the initial 12% to 20%, and in a sample with a kaolinite additive, only up to 16.8% (Figure 67).




Figure 67. Influence of the mineralogical composition of clay in quartz sand (fine sand-1) on the accumulation of moisture in a frozen hydrate-saturated sample ($W=12\%$) after 4 hours of interaction with a frozen salt solution of 0.1 N NaCl at $-6\text{ }^{\circ}\text{C}$: A - kaolinite; B - montmorillonite.

The calculation of the average migration flow of salt for the described experiments shows that its value decreases with an increase in the amount of clay particles (Figure 68). Thus, at a temperature of $-6\text{ }^{\circ}\text{C}$, the average flux density of salt ions in frozen hydrate-saturated fine sand-2 without impurities was $10 \cdot 10^{-10} \text{ mol/cm}^2 \cdot \text{s}$, and with the addition of 15% kaolinite, the flux decreased almost 2 times (to $5.5 \cdot 10^{-10} \text{ mol/cm}^2 \cdot \text{s}$). At the same time, when clay particles of montmorillonite composition were added to quartz sand in an amount of 15%, the value of the average migration flow decreased even more (almost 4 times) and amounted to $2.5 \cdot 10^{-10} \text{ mol/cm}^2 \cdot \text{s}$ (Figure 68).




Figure 68. Influence of the content of clay particles of various compositions (kaolinite and montmorillonite) in quartz sand (fine sand-2) on the average flux density of Na⁺ ions into a frozen hydrate-containing sample over 4 hours of interaction with a frozen salt solution of 0.1N NaCl (p=0.1MPa, t= -6°C): A – kaolinite; B – montmorillonite.

Analysis of the accumulation of salts in the studied samples and the change in hydrate content in the process of salt transfer made it possible to reveal the dependence of the critical concentration of salts in the samples (C_{cr}), which causes the complete decomposition of metastable pore gas hydrates (at P=0.1 MPa), on the content of clay particles of various mineral composition in quartz sand (Figure 69).




Figure 69. Influence of the content of clay particles of different mineral composition in quartz sand (fine sand-2) on the critical concentration (C_{cr}) causing the decomposition of pore hydrate in a frozen sand sample

(t= -6°C, p=0.1MPa, C_{sol} =0.1N): A – kaolinite; B – montmorillonite.

As experimental data show, the critical concentration (C_{cr}) increases with the increase in the content of clay particles in quartz sand (Figure 69). Thus, with an increase in the content of kaolinite to 15%, the critical concentration (C_{cr}) increased 4 times (from 0.5% to 2.1%), and with an increase in the content of montmorillonite particles, the critical concentration increased almost 6 times (from 0.5 to 3.1%).

An experimental study of the influence of the particle size of the studied quartz sand on salt transfer in frozen hydrate-containing sediment showed that with an increase in the size of the predominant sand fraction from 0.1-0.2 (fine sand-2) to 0.2-0.5 mm (medium sand), the intensity of accumulation of salt ions increases, which accompanied by an increase in the dissociation of pore methane hydrate. Thus, at a fixed negative temperature (-6°C), 4 hours after the interaction of frozen hydrate-saturated fine sand-2 with NaCl 0.1N salt solution, Na^+ ions penetrated 5.4 cm into the depth of the sample, and 7.2 cm into medium sand (Figure 70).




Figure 70. Influence of the size of sand particles on the accumulation of salt ions (Na^+) and the decrease in the hydrate coefficient (K_h) in frozen hydrate-containing samples ($K_h^{\text{in}} \sim 0.4$, $W=12\%$, $t = -6^{\circ}\text{C}$) 4 hours after the start of interaction with frozen salt solution 0.1 N NaCl: A - fine sand-2 (0.1-0.2 mm ~ 80%); B – medium sand (0.2-0.5 mm ~ 60%).

An increase in the intensity of migration and accumulation of salt ions in frozen hydrate-saturated sand interacting with a NaCl solution at a fixed negative temperature (-6°C), with an increase in the predominant size of sand particles (from 0.1-0.2 to 0.2-0.5 mm) is accompanied by a more active decomposition of the pore hydrate. Thus, 4 hours after the contact of a frozen hydrate-containing fine sand-2 sample (0.1–0.2 cm) with a 0.1N NaCl solution at -6°C , the pore hydrate completely decomposed at a distance of 5 cm from the contact area. In a frozen hydrate-saturated sample with a predominance of a coarser fraction (medium sand), complete decomposition of the hydrate was recorded to a depth of 6.2 cm. At the same time, K_h , in the area far from the contact, decreased significantly (from 0.4 to 0.2) (Figure 70).

A special series of experiments was devoted to the analysis of the migration of salt ions in frozen hydrate saturated sediment with different content of particles less than 0.05 mm (silt): fine sand-2 (~1% silt), fine sand-3 (~5% silt) and sandy loam (~ 18% silt). The initial characteristics of frozen hydrate-saturated samples are presented in Table 13.

Table 13. Properties of hydrate-saturated frozen sand with different amount of silt.

Sample	Type of sediment	Amount of silt, %	Water content, %	Density, g/cm^3	Porosity, %	Ice saturation, %	Hydrate saturation, %
1	Fine sand-2	1	12	1.8	38	15	44
2	Fine sand-3	5	12	1.83	37	16	37
3	Sandy loam	18	12	1.86	35	18	37

In the course of experimental modeling, it was found that with an increase in the content of silt, the migration and accumulation of salt ions in frozen hydrate-containing samples decreases. So, after 4 hours of interaction with NaCl 0.1 N salt solution at a fixed negative temperature of -6°C , with an increase in the amount of silt from 1% to 18%, the penetration depth of ions in the samples decreased from 5.3 cm to 3.8 cm (Figure 71).




Figure 71. Influence of silt content on the accumulation of salt ions in frozen hydrate-containing samples 4 hours after interaction with 0.1 N NaCl solution at $t = -6^{\circ}\text{C}$: A – fine sand-2 (silt ~ 1%); B – fine sand-3 (silt ~ 5%); C – sandy loam (silt ~ 18%).

A decrease in the migration and accumulation of salts with an increase in the content of silt is accompanied by a decrease in the dissociation of pore gas hydrate. For example, in frozen hydrate-saturated fine sand-2, where the silt content did not exceed 1%, the hydrate completely dissociated in a region of 4.8 cm from the contact. And in the frozen hydrate-containing fine sand-3 sample with a silt content of about 5%, the pore hydrate dissociated to a depth of 4.2 cm. In the sandy loam sample with a silt content of 18%, the pore hydrate decomposition area was only 3.3 cm (Figure 72).




Figure 72. Influence of silt content on the hydrate coefficient (K_h) of frozen hydrate-containing samples ($K_h^{\text{in}} \sim 0.4-0.5$, $W=12\%$) 4 hours after interaction with 0.1 N NaCl solution at $t = -6^{\circ}\text{C}$: A – fine sand-2 (silt ~ 1%); B – fine sand-3 (silt ~ 5%); C – sandy loam (silt ~ 18%).

The different content of silt in frozen hydrate-saturated samples interacting with salt solution is also reflected in moisture accumulation during salt migration. With an increase in the content of silt in frozen hydrate-saturated samples, an increase in water content was recorded in the near-contact area. So, in a sample with a silt content of 5%, the increase in water content was up to 5%, and in a sample with a silt content of 18%, water content increased by more than 2 times (from 12 to 28%) (Figure 73).




Figure 73. Influence of silt content on moisture accumulation in a frozen hydrate-saturated sample ($W=12\%$) after 4 hours of interaction with a frozen salt solution of 0.1 N NaCl at $-6\text{ }^{\circ}\text{C}$: A – fine sand-2 (silt $\sim 1\%$); B – fine sand-3 (silt $\sim 5\%$); C – sandy loam (silt $\sim 18\%$).

An increase in the average water flux in frozen hydrate-saturated samples was also recorded with an increase in the content of silt (from 1 to 18%): from $1.5 \cdot 10^{-7}\text{ g}\cdot\text{s}$ to $22.8 \cdot 10^{-7}\text{ g}\cdot\text{s}$.

For the described samples, the critical concentrations of the decomposition of the pore hydrate during their interaction with the NaCl salt solution were also calculated. This concentration (C_{cr}) increased from 0.05% to 0.19% as silt increased from 1% to 18% (Table 14).

Table 14. Dependence of the critical NaCl concentration on the amount of silt in frozen hydrate saturated sandy sediment at the temperature -6°C .

Sediment	Ccr, mg - EQ/100 g	Ccr, %
Medium sand (silt ~1%)	2.0	0.05
Fine sand-1 (silt ~1%)	3.4	0.13
Fine sand-2 (silt ~5%)	7.5	0.18
Sandy loam (silt~18%)	7.9	0.19

For the studied frozen hydrate-containing samples of **different fineness**, the average salt flux was calculated during their interaction with 0.1N NaCl solution at a temperature of -6°C and $p=0.1\text{MPa}$ (Figure 74). It was found that in sandy-loamy sandy sediments with an increase in the content of silt from 1 to 18%, represented mainly by quartz-feldspar particles, the flux of Na^+ ions increases from $6.2 \cdot 10^{-10} \text{ mol/cm}^2 \cdot \text{s}$ to $16.3 \cdot 10^{-10} \text{ mol/cm}^2 \cdot \text{s}$. **This is due to the increase in the paths for migration with a decrease in particle size without a noticeable increase in their sorption capacity.**




Figure 74. Influence of fineness and amount of clay particles (kaolinite) on the average migration flow of Na⁺ in frozen hydrate-saturated samples interacting with NaCl solution 0.1 N for 4 hours at t= -6°C: A – fine sand-3; B – fine sand-2 with kaolinite (from 0 to 10%); C – medium sand; D – sandy loam.

When finely dispersed clay particles (kaolinite composition) are added to quartz sand (fine sand-2), on the contrary, the density of the ion migration flux decreases, almost 2 times (from 9.9 to 6.5 · 10⁻¹⁰ mol/cm²·s) (Figure 74). This behavior is associated with sorption processes that occur on the surface of clay particles during their interaction with salt ions. This is confirmed by the addition of clay particles of montmorillonite composition (having a high sorption capacity) to the sand, where a 4-fold decrease in the salt flux density is observed (Figure 68).

Thus, with an increase in the content of silt in frozen hydrate-containing sand samples interacting with a salt solution, the intensity of the accumulation of salt ions and the dissociation of pore hydrate increases. When more finely dispersed clay particles (kaolinite and montmorillonite) are added, on the contrary, the accumulation of salt ions and the dissociation of the hydrate decreases.

The experimental results demonstrate the active migration and accumulation of salts in frozen hydrate saturated sediments, what causes active hydrate dissociation in pore space. Salt transfer and

hydrate dissociation during interaction between frozen hydrate saturated sediments and salt solution depends on many factors, such as pressure, temperature, sediments (dispersion, clay content) and contact solution (concentration, chemical composition) characteristics. Experimental data demonstrates, that migration of salt solution may lead to the destruction of intrapermafrost and subpermafrost gas hydrate formations, permafrost thawing, and methane emissions.

Chapter 8. Concept models of hydrate interaction with natural and technical salt solution in hydrate saturated collectors in permafrost

As confirmed by the results of experimental modeling on the interaction between frozen hydrate-saturated sediments and salt solutions, salt ions can migrate from natural and technogenic solutions to frozen horizons containing gas hydrates and can cause intrapermafrost hydrate decomposition in the Arctic region. This is possible when salt ions penetrate from sea water on the Arctic shelf into underwater permafrost containing gas hydrates, when intrapermafrost saline water (cryopegs) flows down the frozen section as a result of the evolution of the permafrost zone, as well as during technogenic impact (for example, when drilling wells in frozen strata containing hydrates). And the intensity of mass transfer processes is defined by internal (sediments characteristics) and external factors (ambient temperature, pressure, salt concentration and composition). These issues have not been sufficiently studied until recently.

The obtained experimental data allow us to present some model-scheme of destabilization of intrapermafrost hydrate during salt migration in frozen hydrate saturated sediments in continental conditions (Figure 73) and on the shelf (Figure 74).

In continental conditions, initially, intrapermafrost gas hydrates exist both in stable and metastable conditions. At the same time, there are cryopegs around which there is a certain salinity

halo (Figure 73a). This salinity halo very slowly spread over time, without significant influence on the surrounding frozen sediments. However, during the evolution of the permafrost zone (for example, with an increase in the sediment temperature) or under technogenic impact (for example, during drilling), the migration of salts from cryopegs proceed faster down the section, and a salinization front is formed (Figure 73b). Salt ions, when reach hydrate saturated frozen horizon, destabilized hydrates and at a certain critical concentration causes their complete dissociation. Since the critical concentration of hydrate dissociation in frozen sediment at a gas pressure below equilibrium is lower than at the pressure above equilibrium, the destabilization of metastable gas hydrate deposits occurs first (Figure 73c), as does the thawing of host frozen sediments. And the diffusion of salts into the frozen horizons, which contain gas hydrates under stable conditions, proceeds much more slowly. When the metastable gas hydrate reservoir had already dissociated, the salt front only partially destabilized gas hydrates at the higher pressure (Figure 73d). The methane released during the dissociation of gas hydrates migrates through permeable areas of frozen horizons (along taliks), and only part of it enters the atmosphere. The main mass of methane accumulates under impermeable frozen horizons, forming a gas deposit and creating zone of increased pressure. Further emission of methane can cause the critical gas pressure to be exceeded, followed by an explosion and the formation of a crater [106,248,249,336].




Figure 75. Proposed model of interaction between saline horizons and frozen deposits containing gas hydrates: a) initial state; b) start of salt migration from cryopegs due to ambient temperature increase; c) dissociation of metastable hydrates occurs; d) dissociation of stable hydrates occurs.

On the Arctic shelf in the conditions of submarine permafrost bottom sediments (in frozen and thawed state), including hydrate containing, are in the contact with sea water. Consequently, salt migration down by the cross-section occur, what can play crucial role in the hydrate destabilization in the submarine permafrost (Figure 74).




Figure 76. The proposed conceptual model on the interaction between frozen hydrate-containing sediments and sea water: a) preliminary state – sea regression; b) after sea transgression – start of interaction between frozen sediments and sea water; c) migration of salt into permafrost sediments (development of salinity and thawing fronts); d) propagation salt front into sediments, saturated with intrapermafrost gas hydrate followed by hydrate decomposition and active methane emission.

Initially (Figure 74A), during sea regression gas hydrates exist in the stable and metastable state without significant change. Then sea transgression occurs (Figure 74B) and frozen hydrate-containing sediments start to interact with salted sea water, sediments temperature rises, and thawing front is formed (Figure 74 – 1) [278,331,337]. Over time, the rate of thawing of underwater permafrost decreases. In this case, the penetration of sea water into the thawing zone is observed (Figure 74b). Since frozen rocks are permeable to salt ions, there is an advancing movement of salt ions from sea water relative to the thawing front. During the migration and accumulation of salt ions in frozen deposits, the temperature of phase transitions decreases and the content of the liquid phase in soils

increases. As a result, the deceleration of the thawing front decreases. Salt ions migrating in frozen deposits, upon reaching horizons containing relict hydrates, will destabilize them and contribute to their active dissociation and methane release (Figure 74c), thus forming the second front of phase transitions - the dissociation front of gas hydrates. It occurs when a certain critical salt concentration is reached in the frozen hydrate-containing rock. This concentration, as shown by experimental studies, depends on the composition of hydrate-containing deposits and thermobaric conditions. During phase transitions ice-water / hydrate-water (ice), gas caused by salt transfer, the permeability of saline permafrost increases [338]. Methane formed during the dissociation of gas hydrates subsequently migrates to the bottom surface and enters the sea water, where it partially dissolves, oxidizes, and also diffuses through the water column and escapes into the atmosphere (Figure 74c).

This scheme makes it possible to explain the possible cause of numerous gas manifestations recorded on the Arctic shelf in areas of submarine permafrost distribution as a result of destabilization of intrapermafrost gas hydrate formations due to heat and mass transfer processes associated with the migration of salt ions from sea water.

Chapter 9. Conclusion

An analysis of the natural conditions of occurrence of frozen hydrate-bearing sediment indicates the possibility of their interaction with salt solutions. This is possible when salt ions penetrate from sea water on the Arctic shelf into underwater permafrost containing gas hydrates, when intrapermafrost saline water (cryopegs) flows down the frozen section as a result of the evolution of the permafrost zone, as well as during technogenic impact (for example, when drilling wells in frozen strata containing hydrates). Salt migration causes thawing of frozen sediment and destabilization of intrapermafrost gas hydrates, which leads to the emission of methane into the atmosphere. Thus, the

transformation of climatic conditions can occur, as well as the emergence of technological complications during the development of Arctic deposits. However, literature analysis has shown that the issue of interaction between frozen hydrate-saturated sediment and salt solutions has not been studied enough so far.

The conducted experimental studies allow us to draw the following conclusions:

1) The new method for determination of water dissolving salt concentration in sandy sediments through water activity was proposed. The method allows to obtain values of salt ions concentration without special probe preparation. It is fast, cheap and has low error ($\pm 0.2\%$), but applicable only to the low concentration of salts (up to 5%);

2) The new method was created for investigating of hydrate saturated frozen sediments interaction with salt solution under the pressure higher than 0.1 MPa in the pressure cell;

3) The method for temperature change investigation during interaction of frozen hydrate saturated sediments interaction with salt solution was developed. The method allows to register temperature change in time with using of special equipment with high precision of temperature sensors ($\pm 0.05^\circ\text{C}$). All external temperature influence in that method is limited, i.e., ambient temperature is fixed at one value. So, only internal temperature changes in the frozen hydrate saturated sample is observed;

4) Mechanism and dynamics of hydrate saturated frozen sediments interaction with salt solution were experimentally investigated:

- during the interaction of frozen hydrate-saturated sediments with salt solutions, not only active processes of salt migration are observed, which are accompanied by the accumulation of salt ions in the pore space, but also phase transitions of pore ice and dissociation of gas hydrates in the pore space, accompanied by anomalous temperature drops;

- the migration of salt ions into a frozen hydrate-containing sample occurs more intensively than into a sample without hydrates (due to the higher content of liquid water films), and with deeper temperature decrease (temperature decrease was 3 times lower in hydrate saturated sand, than in non-hydrate saturated);
- in the process of salts migration and accumulation, two fronts are formed: thawing of frozen sediments and gas hydrate decomposition. It is noted that the front of decomposition of gas hydrate is ahead of the front of thawing, because critical concentration of hydrate dissociation is lower than critical concentration of thawing;
- Na^+ flux is decreasing with time of interaction between frozen hydrate saturated and salt solution. After 25 hours of contact between hydrate saturated sand and NaCl solution salt flux decreased by 10 times.
- based on the experimental data scheme of water phase transition in frozen sand saturated with stable and metastable hydrate is proposed, furthermore according temperature effect during that process is analysed.

5) Experimental investigation of different factors influence on the process of hydrate saturated frozen sediments and salt solution interaction was conducted. The following results was obtained:

- the rate of salt penetration in frozen sediments and the rate of hydrate dissociation is increasing when the experimental conditions are transferred from stable to metastable state (from 4 MPa to 0.1 MPa by 2 times both), as well as Na^+ flux from 1 to $38 \cdot 10^{-10}$ mol/cm²·sec. As it was assumed, such dependence connected with decrease in liquid water amount in hydrate saturated frozen sediments with increasing gas pressure;
- higher temperature leads to more active decomposition of the hydrate in frozen sediments as a result of salt penetration. The dissociation rate is more than 4 times higher when the

- temperature of interaction rises 3 times from -20 to -6.5 °C. Salt critical concentration of hydrate decomposition in porous media of frozen sediment vary from 3.4% to 0.6% when temperature of interaction increase from -16 to -2.5 °C;
- higher concentration of the contact solution leads to more active hydrate decomposition in frozen hydrate saturated sediments interacting with salt solution. The penetration of Na⁺ is 3 times greater with an increase in concentration of the contact solution by 2 times. The rate of hydrate dissociation is 6 times faster in the case of contact with a 0.4N solution than in contact with a 0.2N solution. The higher the concentration of the contacting solution, the more significant the decrease in temperature associated with the more intense dissociation of the hydrate, and the longer the time required for the temperature to reach equilibrium (up to the initial temperature values);
 - hydrate stability in pore media of frozen sediments increases in the series MgCl₂-NaCl-CaCl₂-KCl-Na₂SO₄. The intensity of pore hydrate dissociation is 3 times higher upon contact with a NaCl solution than upon contact with Na₂SO₄ solution, and 2 times higher than upon contact with CaCl₂ solution. Data confirmed by computer simulations in HydraFlash;
 - rise of clay content in sand decreases intensity of salt accumulation and hydrate dissociation: critical concentration increases by 4 times and the average salt flux decreases by 2 times when clay content change from 0 to 15%;
 - more active salt accumulation and hydrate dissociation happen in the sand with kaolinite particles, than in the sand with montmorillonite;
 - salt accumulation is more active and hydrate decomposes more rapidly when amount of silt with low sorption ability increases. Thus, more active salt migration and hydrate dissociation was observed in sand than in sandy loam.

6) On the basis of experimental data, a conceptual scheme for the destabilization of intrapermafrost formations in the areas of permafrost distribution in continental conditions and on the Arctic shelf is proposed. This scheme makes it possible to explain the possible cause of numerous gas manifestations recorded in the Arctic area of permafrost distribution as a result of destabilization of intrapermafrost gas hydrate formations due to heat and mass transfer processes associated with the migration of salt ions.

The conducted experimental studies allow us to take a newly look at the mechanism and patterns of migration of salt ions in frozen hydrate-saturated sediment during their contact. Experimental data demonstrate the potential danger of destabilization of gas hydrate formations in the Arctic as a result of exposure to natural and artificial solutions, especially for the oil and gas industry.

Bibliography

1. Kudryashov, B.; Yakovlev, A. *Well Drilling in Permafrost (Published in Russian)*; Nedra: Moscow, Russia, 1983; p. 283.
2. Lu, S. M. A Global Survey of Gas Hydrate Development and Reserves: Specifically in the Marine Field. *Renew. Sustain. Energy Rev.* **2015**, *41*, 884–900.
<https://doi.org/10.1016/j.rser.2014.08.063>.
3. Timofeev, N. G.; Skryabin, R. M.; Atlasov, R. A. Wells Drilling Temperature Regime under the Cryolithozone Conditions (Published in Russian). *Sci. Educ.* **2017**, No. 3, 51–56.
4. Ma, Y.; Yang, J.; Feng, P.; Zhang, C. The Challenges and Key Technology of Drilling Safety in the Area of the Arctic. In *Springer Series in Geomechanics and Geoengineering*; Springer, 2020; pp 522–532. https://doi.org/10.1007/978-981-13-7127-1_48.
5. Yakushev, V. S.; Collett, T. S. Gas Hydrates in Arctic Regions: Risk to Drilling and Production. In *Proceedings of the Second International Offshore and Polar Engineering Conference*; San Francisco, 1992; Vol. 1, pp 669–673.
<https://doi.org/10.1016/j.jaad.2006.01.013>.
6. Medvedsky, R. I. *Construction and Operation of Oil and Gas Wells in Permafrost (Published in Russian)*; Nedra: Moscow, Russia, 1987; p. 127.
7. Kontorovich, A. E.; Epov, M. I.; Burshtein, L. M.; Kaminskii, V. D.; Kurchikov, A. R.; Malyshev, N. A.; Prischepa, O. M.; Safronov, A. F.; Stupakova, A. V.; Suprunenko, O. I. Geology and Hydrocarbon Resources of the Continental Shelf in Russian Arctic Seas and the Prospects of Their Development. *Russ. Geol. Geophys.* **2010**, *51*, *1*, 3–11.
<https://doi.org/10.1016/j.rgg.2009.12.003>.

8. Yakushev, V. S. *Gas and Gas Condensate Fields Development at Complicated Geocryological Conditions (Published in Russian)*; Gubkin Russian State University: Moscow, Russia, 2014; p. 188, ISBN 978-5-91961-114-1.
9. Wegner, S. A.; Campbell, K. J. Drilling Hazard Assessment for Hydrate Bearing Sediments Including Drilling through the Bottom-Simulating Reflectors. *Mar. Pet. Geol.* **2014**, *58*, PA, 382–405. <https://doi.org/10.1016/j.marpetgeo.2014.08.007>.
10. Nimblett, J. N.; Shipp, R. C.; Strijbos, F. Gas Hydrate as a Drilling Hazard: Examples from Global Deepwater Settings. In *Proceedings of the Annual Offshore Technology Conference*; 2005; Vol. 2005-May, pp 1429–1435. <https://doi.org/10.4043/17476-ms>.
11. McConnell, D. R.; Zhang, Z.; Boswell, R. Review of Progress in Evaluating Gas Hydrate Drilling Hazards. *Mar. Pet. Geol.* **2012**. <https://doi.org/10.1016/j.marpetgeo.2012.02.010>.
12. Max, M. *Natural Gas Hydrate In Oceanic and Permafrost Environments*; Kluwer Academic Publishers: Washington, 2000; p. 419, ISBN 978-1-4020-1362-1. <https://doi.org/10.1007/978-94-011-4387-5>.
13. Collett, T. S.; Dallimore, S. R.; Max, M. D. Natural Gas Hydrate in Oceanic and Permafrost Environments. In *v. 5, chapter 5*; Springer Science & Business Media, 2003; Vol. 5.
14. Sloan Jr., E. D.; Koh, C. A. *Clathrate Hydrates of Natural Gases*; CRC Press: Boca Raton, FL, USA, 2007; p. 752, ISBN 9780429129148. <https://doi.org/10.1201/9781420008494>.
15. Istomin, V. A.; Yakushev, V. S. *Gas Hydrates in Nature (Published in Russian)*; Nedra: Moscow, 1992; p. 235.
16. Boswell, R.; Hancock, S.; Yamamoto, K.; Collett, T.; Pratap, M.; Lee, S.-R. Natural Gas Hydrates. In *Future Energy*; Elsevier, 2020; pp 111–131. <https://doi.org/10.1016/B978-0-08->

102886-5.00006-2.

17. Chuvilin, E.; Davletshina, D. Formation and Accumulation of Pore Methane Hydrates in Permafrost: Experimental Modeling. *Geosciences* **2018**, *8*, *12*, 467.
<https://doi.org/10.3390/geosciences8120467>.
18. Collett, T.; Bahk, J.-J.; Baker, R.; Boswell, R.; Divins, D.; Frye, M.; Goldberg, D.; Husebø, J.; Koh, C.; Malone, M.; et al. Methane Hydrates in Nature—Current Knowledge and Challenges. *J. Chem. Eng. Data* **2015**, *60*, *2*, 319–329. <https://doi.org/10.1021/je500604h>.
19. Ruppel, C. Permafrost-Associated Gas Hydrate: Is It Really Approximately 1 % of the Global System? *J. Chem. Eng. Data* **2015**, *60*, *2*, 429–436. <https://doi.org/10.1021/je500770m>.
20. Waite, W. F.; Ruppel, C. D.; Boze, L.-G.; Lorenson, T. D.; Buczkowski, B. J.; McMullen, K. Y.; Kvenvolden, K. A. Preliminary Global Database of Known and Inferred Gas Hydrate Locations. *U.S. Geol. Surv. data release* **2020**. <https://doi.org/10.5066/P9LLFVJM>.
21. Ruppel, C. D. Gas Hydrate in Nature. *U.S. Geol. Surv.* **2018**, *Fact Sheet*, *4*.
<https://doi.org/https://doi.org/10.3133/fs20173080>.
22. Ershov, E. D.; Lebedenko, Y. P.; Chuvilin, E. M.; Istomin, V. A.; Yakushev, V. S. Features of the Existence of Gas Hydrates in the Cryolithozone (Published in Russian). *Rep. Acad. Sci. USSR* **1991**, *321*, 788–791.
23. Chuvilin, E.; Bukhanov, B.; Davletshina, D.; Grebenkin, S.; Istomin, V. Dissociation and Self-Preservation of Gas Hydrates in Permafrost. *Geosciences* **2018**, *8*, *12*, 431:1-431:12.
<https://doi.org/10.3390/geosciences8120431>.
24. Hachikubo, A.; Takeya, S.; Chuvilin, E.; Istomin, V. Preservation Phenomena of Methane Hydrate in Pore Spaces. *Phys. Chem. Chem. Phys.* **2011**, *13*, *39*, 17449–17452.

<https://doi.org/10.1039/c1cp22353d>.

25. Takeya, S.; Ebinuma, T.; Uchida, T.; Nagao, J.; Narita, H. Self-Preservation Effect and Dissociation Rates of CH₄ Hydrate. *J. Cryst. Growth* **2002**, *237*, 379–382.
[https://doi.org/10.1016/S0022-0248\(01\)01946-7](https://doi.org/10.1016/S0022-0248(01)01946-7).
26. Stern, L. A.; Circone, S.; Kirby, S. H.; Durham, W. B. Anomalous Preservation of Pure Methane Hydrate at 1 Atm. *J. Phys. Chem. B* **2001**, *105*, 9, 1756–1762.
<https://doi.org/10.1021/jp003061s>.
27. Kuhs, W. F.; Genov, G.; Staykova, D. K.; Hansen, T. Ice Perfection and Onset of Anomalous Preservation of Gas Hydrates. *Phys. Chem. Chem. Phys.* **2004**, *6*, 21, 4917–4920.
<https://doi.org/10.1039/B412866D>.
28. Istomin, V. A.; Yakushev, V. S. Makhonina, N. A.; Kwon, V. G.; Chuvilin, E. M. Self-Preservation Phenomenon of Gas Hydrate. *Gas Ind. Russ.* **2006**, *4*, 16–27.
29. Istomin, V. A.; Yakushev, V. S. Gas-Hydrates Self-Preservation Effect. *Phys. Chem. Ice* **1992**, 136–140.
30. Chuvilin, E. M.; Tumskoy, V. E.; Tipenko, G. S.; Gavrilov, A. V.; Bukhanov, B. A.; Tkacheva, E. V.; Audibert-Hayet, A.; Cauquil, E. Relic Gas Hydrate and Possibility of Their Existence in Permafrost Within the South-Tambey Gas Field. *SPE Arct. Extrem. Environ. Tech. Conf. Exhib. Soc. Pet. Eng.* **2013**, 166925:1-166925:9.
31. Chuvilin, E.; Ekimova, V.; Bukhanov, B.; Grebenkin, S.; Shakhova, N.; Semiletov, I. Role of Salt Migration in Destabilization of Intra Permafrost Hydrates in the Arctic Shelf: Experimental Modeling. *Geosciences* **2019**, *9*, 4, 188:1-188:18.
<https://doi.org/10.3390/geosciences9040188>.

32. Chuvilin, E.; Davletshina, D.; Ekimova, V.; Bukhanov, B.; Shakhova, N.; Semiletov, I. Role of Warming in Destabilization of Intrapermafrost Gas Hydrates in the Arctic Shelf: Experimental Modeling. *Geosciences* **2019**, *9*, 407:1-407:12.
<https://doi.org/10.3390/geosciences9100407>.
33. Yakushev, V. S. *Natural Gas and Gas Hydrates in the Permafrost (Published in Russian)*; Gazprom VNIIGAZ: Moscow, 2009; p. 192.
34. Yamamoto, K.; Boswell, R.; Collett, T. S.; Dallimore, S. R.; Lu, H. Review of Past Gas Production Attempts from Subsurface Gas Hydrate Deposits and Necessity of Long-Term Production Testing. *Energy & Fuels* **2022**, *36*, *10*, 5047–5062.
<https://doi.org/10.1021/acs.energyfuels.1c04119>.
35. Zhao, R.; Chang, H. J.; Chen, K. L. Environmental Risk of Gas Hydrates Exploitation in Tibetan Plateau Permafrost. *Adv. Mater. Res.* **2014**, *955–959*, 2114–2117.
<https://doi.org/10.4028/www.scientific.net/AMR.955-959.2114>.
36. Yakushev, V. Natural Gas Liberations around Production Wells at Russian Arctic Gas Fields. *Geosciences* **2020**, *10*, *5*, 184:1-184:6. <https://doi.org/10.3390/geosciences10050184>.
37. Max, M. D.; Johnson, A. H.; Dillon, W. P. *Natural Gas Hydrate - Arctic Ocean Deepwater Resource Potential*; Springer, Cham: Dordrecht, Netherlands, 2013; p. 113 p., ISBN 9783319025070. <https://doi.org/https://doi.org/10.1007/978-3-319-02508-7>.
38. Chuvilin, E.; Tipenko, G.; Bukhanov, B.; Istomin, V.; Pissarenko, D. Simulating Thermal Interaction of Gas Production Wells with Relict Gas Hydrate-Bearing Permafrost. *Geosciences* **2022**, *12*, *3*, 115. <https://doi.org/10.3390/geosciences12030115>.
39. Chuvilin, E.; Ekimova, V.; Davletshina, D.; Krivokhat, E.; Shilenkov, V.; Bukhanov, B.

- Pressure Influence on Salt Migration in Frozen Hydrate-Saturated Sediments: Experimental Modeling. *Energy & Fuels* **2022**, *12*, 7, 261. <https://doi.org/10.1021/acs.energyfuels.2c01282>.
40. Chuvilin, E.; Ekimova, V.; Davletshina, D.; Bukhanov, B.; Krivokhat, E.; Shilenkov, V. Migration of Salt Ions in Frozen Hydrate-Saturated Sediments: Temperature and Chemistry Constraints. *Geosciences* **2022**, *12*, 7, 276. <https://doi.org/10.3390/geosciences12070276>.
41. Chuvilin, E.; Bukhanov, B.; Istomin, V. Cryopegs as Destabilization Factor of Intra-Permafrost Gas Hydrates. In *EGU General Assembly Conference*; 2016; Vol. 18, pp EPSC2016-3509.
42. Yakushev, V. S.; Istomin, V. A. Causes of Gas Emissions in Frozen Rocks of the Yamburg Gas Condensate Field (Published in Russian). In *Features of gas well development in difficult geocryological conditions*; VNIIGAZ: Moscow, Russia, 1987; pp 119–127.
43. Bing, H.; He, P.; Zhang, Y. Cyclic Freeze–Thaw as a Mechanism for Water and Salt Migration in Soil. *Environ. Earth Sci.* **2015**, *74*, 1, 675–681. <https://doi.org/10.1007/s12665-015-4072-9>.
44. Maksimov, A. M.; Tsypkin, G. G. Formation of a Two-Phase Zone in the Interaction of Wet Rocks with Cooled Salt Solution. *J. Eng. Phys.* **1988**, *55*, 3, 1014–1019. <https://doi.org/10.1007/BF00870486>.
45. Ershov, E. D.; She, Z. S.; Lebedenko, Y.; Chuvilin, E. M.; Kryuchkov, K. Y. Mass Transfer and Deformation Processes in Frozen Rocks Interacting with Aqueous Salt Solutions (Published in Russian). In *Engineering-geological study and evaluation of frozen, freezing and thawing soils (IGK-92). Materials of the III Scientific and Technical Workshop*; Russian State Scientific Research Institute of Hydraulic Engineering: St Petersburg, 1993; pp 67–77.

46. Lebedenko, Y. P. Cryogenic Migration of Ions and Bound Moisture in Ice-Saturated Frozen Rocks (Published in Russian). *Eng. Geol.* **1989**, No. 4, 21–30.
47. Ershov, E. D.; Lebedenko, Y. P.; Chuvilin, E. M.; Ganova, S. D. Features of Moisture Migration in Frozen Fine Soils under the Action of Osmotic Forces (Published in Russian). In *Geocryological research*; MSU: Moscow, Russia, 1987; pp 150–157.
48. Chuvilin, E. M.; Ershov, E. D.; Smirnova, O. G. Ionic Migration in Frozen Soils and Ice. In *Proceedings of the 7th International Permafrost Conference*; Yellowknife, Canada, 1998; pp 167–171.
49. Ershov, E. D.; Komarov, I. A.; Chuvilin, E. M. Forecast of Processes of Interaction of Liquid Technogenic Brines Buried in the Frozen Sediment Massif (Published in Russian). *Geoecology. Eng. Geol. Hydrogeol. Geocryol.* **1997**, No. 2, 19–29.
50. Vasiliev, V. I.; Maksimov, A. M.; Petrov, E. E.; Tsytkin, G. G. *Heat and Mass Transfer in Freezing and Thawing Soils (Published in Russian)*; Nauka: Moscow, Russia, 1996; p. 224.
51. Permyakov, P. P. Mathematical Modeling of Technogenic Pollution in Frozen Soils (Published in Russian). *Bull. Tomsk Polytech. Univ. Georesource Eng.* **2004**, 307, 5, 63–68.
52. Permyakov, P. P.; Ammosov, A. P.; Popov, G. G.; Khokholova, S. T. Mathematical Models in Predicting Man-Caused Pollution of the Permafrost Zone (Published in Russian). In *VIII Russian conference “Modern methods of mathematical modeling of natural and man-made disasters” October 26 - 28*; Kemerovo, 2005; pp 1–11.
53. Feng, J.-C.; Wang, Y.; Li, X.-S. Hydrate Dissociation Induced by Depressurization in Conjunction with Warm Brine Stimulation in Cubic Hydrate Simulator with Silica Sand. *Appl. Energy* **2016**, 174, 181–191. <https://doi.org/10.1016/j.apenergy.2016.04.090>.

54. Li, S.; Wang, Z.; Xu, X.; Zheng, R.; Hou, J. Experimental Study on Dissociation of Hydrate Reservoirs with Different Saturations by Hot Brine Injection. *J. Nat. Gas Sci. Eng.* **2017**, *46*, 555–562. <https://doi.org/10.1016/j.jngse.2017.07.032>.
55. Kamath, V. A.; Mutalik, P. N.; Sira, J. H.; Patil, S. L. Experimental Study of Brine Injection Depressurization of Gas Hydrates Dissociation of Gas Hydrates. *SPE Form. Eval.* **1991**, *6*, 04, 477–484. <https://doi.org/10.2118/19810-PA>.
56. Kamath, V. A.; Godbole, S. P. Evaluation of Hot-Brine Stimulation Technique for Gas Production From Natural Gas Hydrates. *J. Pet. Technol.* **1987**, *39*, *11*, 1379–1388. <https://doi.org/10.2118/13596-PA>.
57. Malakhova, V. V.; Eliseev, A. V. Salt Diffusion Effect on the Submarine Permafrost State and Distribution as Well as on the Stability Zone of Methane Hydrates on the Laptev Sea Shelf (Published in Russian). *Ice Snow* **2020**, *60*, *4*, 533–546.
58. Polozkov, K.A., Basniev, K.S., Gaftunyak, P. I. Complications Arising from the Construction and Operation of Wells in Permafrost Zones, and Measures to Prevent Them (Published in Russian). *Constr. oil gas wells L. sea* **2010**, *1*, 6–10.
59. Fjellanger, E.; Kontorovich, A. E.; Barboza, S. A.; Burshtein, L. M.; Hardy, M. J.; Livshits, V. R. Charging the Giant Gas Fields of the NW Siberia Basin. In *Petroleum Geology Conference Proceedings*; Geological Society of London, 2010; Vol. 7, pp 659–668. <https://doi.org/10.1144/0070659>.
60. Ershov, E. D. *General Geocryology (Published in Russian)*; Moscow University Press: Moscow, Russia, 2002; p. 682.
61. Harris, S. A.; Brouchkov, A.; Guodong, C. *Geocryology: Characteristics and Use of Frozen*

Ground and Permafrost Landforms; CRC Press, 2017; p. 1–765, ISBN 9781351681629.

<https://doi.org/10.4324/978135166988>.

62. Kudryavtsev, V. A.; Dostovalov, B. N.; Romanovsky, N. N.; Kondratieva, K. A.; Melamed, V. G. *General Permafrost (Geocryology) (Published in Russian)*; MSU: Moscow, Russia, 1978; p. 464.
63. Chuvilin, E. M.; Perlova, E. V.; Yakushev, V. S. Classification of the Intrapermafrost Sediment Gas Component (Published in Russian). *Cryosph. Earth* **2005**, *9*, 3, 73–76.
64. Istomin, V.; Chuvilin, E.; Bukhanov, B.; Uchida, T. Pore Water Content in Equilibrium with Ice or Gas Hydrate in Sediments. *Cold Reg. Sci. Technol.* **2017**, *137*, 60–67.
<https://doi.org/10.1016/j.coldregions.2017.02.005>.
65. Akagawa, S.; Iwahana, G.; Watanabe, K.; Chuvilin, E. M.; Istomin, V. A.; Hinkel, K. M. Improvement of Pulse NMR Technology for Determination of Unfrozen Water Content in Frozen Soils. In *the Tenth International Conference on Permafrost, June*; Salekhard, Russia, 2012; pp 21–26.
66. Miyata, Y.; Akagawa, S. An Experimental Study of Dynamic Solid-Liquid Phase Equilibrium in a Porous Medium. *JSME Int. Journal, Ser. B Fluids Therm. Eng.* **1998**, *41*, 3, 590–600.
<https://doi.org/10.1299/jsmeb.41.590>.
67. Kruse, A. M.; Darrow, M. M.; Akagawa, S. Improvements in Measuring Unfrozen Water in Frozen Soils Using the Pulsed Nuclear Magnetic Resonance Method. *J. Cold Reg. Eng.* **2018**, *32*, 1. [https://doi.org/10.1061/\(asce\)cr.1943-5495.0000141](https://doi.org/10.1061/(asce)cr.1943-5495.0000141).
68. Chuvilin, E. M.; Yakushev, V. S.; Perlova, E. V. Gas and Possible Gas Hydrates in the Permafrost of Bovanenkovo Gas Field, Yamal Peninsula, West Siberia. *Polarforschung* **2000**,

68, 1–3, 215–219.

69. Obu, J.; Westermann, S.; Bartsch, A.; Berdnikov, N.; Christiansen, H. H.; Dashtseren, A.; Delaloye, R.; Elberling, B.; Etzelmüller, B.; Kholodov, A.; et al. Northern Hemisphere Permafrost Map Based on TTOP Modelling for 2000–2016 at 1 Km² Scale. *Earth-Science Rev.* **2019**, *193*, 299–316. <https://doi.org/10.1016/j.earscirev.2019.04.023>.
70. Vasiliev, A. A.; Gravis, A. G.; Gubarkov, A. A.; Drozdov, D. S.; Korostelev, Y. V.; Malkova, G. V.; Oblogov, G. E.; Ponomareva, O. E.; Sadurtdinov, M.R. Streletskaya, I.D. Streletsky, D. A.; Ustinova, E. V.; et al. Permafrost Degradation: Results of the Long-Term Geocryological Monitoring in the Western Sector of Russian Arctic (Published in Russian). *Kriosf. Zemli* **2020**, *XXIV*, 2, 15–30. [https://doi.org/10.21782/KZ1560-7496-2020-2\(15-30\)](https://doi.org/10.21782/KZ1560-7496-2020-2(15-30)).
71. HydroSpetsGeologiya. *Information Note with an Assessment of the Current State of the Subsoil in Significant and Man-Made Disturbed Conditions of the Territory of the Arctic Zone of the Russian Federation, Incl. at Geocryological Polygons Marre-Sale and Vorkuta in 2020 (Publis; 2021; p. 149.*
72. Misyurkeeva, N.; Buddo, I.; Kraev, G.; Smirnov, A.; Nezhdanov, A.; Shelokhov, I.; Kurchatova, A.; Belonosov, A. Periglacial Landforms and Fluid Dynamics in the Permafrost Domain: A Case from the Taz Peninsula, West Siberia. *Energies* **2022**, *15*, 8, 2794. <https://doi.org/10.3390/en15082794>.
73. Shakhova, N.; Semiletov, I.; Leifer, I.; Salyuk, A.; Rekant, P.; Kosmach, D. Geochemical and Geophysical Evidence of Methane Release over the East Siberian Arctic Shelf. *J. Geophys. Res. Ocean.* **2010**, *115*, 8, C08007:1–C08007:14. <https://doi.org/10.1029/2009JC005602>.
74. Harrison, W. D.; Osterkamp, T. E. Heat and Mass Transport Processes in Subsea Permafrost 1. An Analysis of Molecular Diffusion and Its Consequences. *J. Geophys. Res.* **1978**, *83*, C9, 185

4707. <https://doi.org/10.1029/JC083iC09p04707>.
75. Angelopoulos, M.; Overduin, P. P.; Miesner, F.; Grigoriev, M. N.; Vasiliev, A. A. Recent Advances in the Study of Arctic Submarine Permafrost. *Permafr. Periglac. Process.* **2020**, *31*, 3, 442–453. <https://doi.org/10.1002/PPP.2061>.
76. Matveeva, T.; Savvichev, A. S.; Semenova, A.; Logvina, E.; Kolesnik, A. N.; Bosin, A. A. Source, Origin, and Spatial Distribution of Shallow Sediment Methane in the Chukchi Sea. *Oceanography* **2015**, *28*, 3, 202–217. <https://doi.org/10.5670/oceanog.2015.66>.
77. Lobkovskiy, L. I.; Nikiforov, S. L.; Dmitrevskiy, N. N.; Libina, N. V.; Semiletov, I. P.; Ananiev, R. A.; Meluzov, A. A.; Roslyakov, A. G. Gas Extraction and Degradation of the Submarine Permafrost Rocks on the Laptev Sea Shelf. *Oceanology* **2015**, *55*, 2, 283–290. <https://doi.org/10.1134/s0001437015010129>.
78. Nicolsky, D.; Shakhova, N. Modeling Sub-Sea Permafrost in the East Siberian Arctic Shelf: The Dmitry Laptev Strait. *Environ. Res. Lett.* **2010**, *5*, 1, 015006:1-015006:9. <https://doi.org/10.1088/1748-9326/5/1/015006>.
79. Portnov, A.; Mienert, J.; Serov, P. Modeling the Evolution of Climate-Sensitive Arctic Subsea Permafrost in Regions of Extensive Gas Expulsion at the West Yamal Shelf. *J. Geophys. Res. G Biogeosciences* **2014**, *119*, 11, 2082–2094. <https://doi.org/10.1002/2014JG002685>.
80. Overduin, P. P.; Schneider von Deimling, T.; Miesner, F.; Grigoriev, M. N.; Ruppel, C.; Vasiliev, A.; Lantuit, H.; Juhls, B.; Westermann, S. Submarine Permafrost Map in the Arctic Modeled Using 1-D Transient Heat Flux (SuPerMAP). *J. Geophys. Res. Ocean.* **2019**, *124*, 6, 3490–3507. <https://doi.org/10.1029/2018JC014675>.

81. Romanovskii, N. N.; Hubberten, H.-W.; Gavrilov, A. V.; Eliseeva, A. A.; Tipenko, G. S. Offshore Permafrost and Gas Hydrate Stability Zone on the Shelf of East Siberian Seas. *Geo-Marine Lett.* **2005**, *25*, 2–3, 167–182. <https://doi.org/10.1007/s00367-004-0198-6>.
82. Shpolyanskaya, N. A.; Streletskaya, I. D.; Surkov, A. Cryolithogenesis within the Arctic Shelf (Modern and Ancient) (Published in Russian). *Earth's Cryospher* **2006**, *10*, 3, 49–60.
83. Fartyshev, A. I. *Features of the Coastal-Shelf Cryolithozone of the Laptev Sea (Published in Russian)*; Nauka: Novosibirsk, 1993; p. 135.
84. Romanovskii, N. N.; Tumskoi, V. E. Retrospective Approach to the Estimation of the Contemporary Extension and Structure of the Shelf Cryolithozone in East Arctic (Published in Russian). *Earth's Cryosph.* **2011**, *15*, 1, 3–14.
85. Perlshtein, G. Z.; Sergeev, D. O.; Tipenko, G. S.; Tumskoy, V. E.; Khimenkov, A. N.; Vlasov, A. N.; Merzlyakov, V. P.; Stanilovskaya, Y. V. Hydrocarbon Gases and Permafrost Zone of the Arctic Shelf (Published in Russian). *Arct. Ecol. Econ.* **2015**, *2*, 18, 35–44.
86. Melnikov, V. P.; Spesivtsev, V. I. *Engineering-Geological and Geocryological Conditions of the Shelf of the Barents and Kara Seas (Published in Russian)*; Nauka: Novosibirsk, 1995; p. 195.
87. Zhigarev, L. A. *Oceanic Permafrost (Published in Russian)*; MSU: Moscow, Russia, 1997; p. 318.
88. Anisimov, O. A.; Streletsky, D. A. Geocryological Hazards of Thawing Permafrost (Published in Russian). *Arct. XXI century. Nat. Sci.* **2015**, *2*, 3, 60–74.
89. Li, L.; Yang, J.; Zou, X.; Yan, L.; Zhang, C.; Yang, P.; Ling, T.; Li, S. Monitoring Technology of Gas Hydrate in Polar Drilling. In *The 31st International Ocean and Polar*

Engineering Conference; Rhodes, Greece, 2021.

90. Sergeev, D. Permafrost-Related Geohazards in Cold Russian Regions. *Oxford Res. Encycl. Nat. Hazard Sci.* **2018**. <https://doi.org/10.1093/ACREFORE/9780199389407.013.291>.
91. Kamenskikh, S. V. Logachev, Y. L.; Nor, A. V.; Ulyasheva, N. M.; Fomin, A. S. *Complications and Accidents during the Construction of Oil and Gas Wells (Published in Russian)*; USTU: Ukhta, Russia, 2014; p. 231.
92. Kamel, A. H. A Novel Mud Formulation for Drilling Operations in the Permafrost. In *SPE Western Regional Meeting Proceedings*; 2013; pp 252–261. <https://doi.org/10.2118/165335-ms>.
93. Avetisyan, N. G. *The Choice of the Type of Drilling Fluid for Drilling in Unstable Rocks. Overview (Published in Russian)*; VNIIOENG: Moscow, Russia, 1983; p. 30.
94. Dong, L.; Wan, Y.; Li, Y.; Liao, H.; Wu, N.; Leonenko, Y. 3d Numerical Simulation on Drilling Fluid Invasion into Natural Gas Hydrate Reservoirs. *SSRN Electron. J.* **2021**. <https://doi.org/10.2139/ssrn.3898481>.
95. Aksenova, N. A.; Rozhkova, O. V. *Drilling Fluids and Well Flushing (Published in Russian)*; Tyumen Industrial University: Tyumen, 2016; p. 168.
96. RosTechNadzor. *On Approval of Federal Norms and Rules in the Field of Industrial Safety "Safety Rules in the Oil and Gas Industry (Published in Russian)*; 2020; Vol. Order №34.
97. Smirnova, O. G. Migration of Ions of Chemical Elements in Frozen Rocks and Ice (Published in Russian), Moscow State University, 1997.
98. Maramzin, A. V.; Ryazanov, A. A. *Drilling Exploratory Wells in Areas with Permafrost Distribution (Published in Russian)*; Nedra: Moscow, Russia, 1971; p. 145–163.

99. Avetov, N. R.; Krasnova, E. A.; Yakushev, V. S. About Possible Causes and Nature of Gas Showing around Gas and Gas-Condensate Wells at the Territory of Yamburg Oil-and-Gas-Condensate Field (Published in Russian). *News gas Sci.* **2018**, *1*, 33, 33–40.
100. Streletskii, D. A.; Streletskaya, I. D. Redistribution of Ions within the Active Layer and Upper Permafrost, Yamal, Russia. In *8th International Conference on Permafrost, 21-25 July*; Balkema: Zurich, Switzerland, 2003; pp 1117–1122.
101. Streletskaya, I. D. Cryopeg Responses To Periodic Climate Fluctuations. In *Permafrost - Seventh International Conference, June*; Yellowknife, Canada, 1998; pp 1021–1025.
102. Zhang, Q.; Wang, Z.; Fan, X.; Wei, N.; Zhao, J.; Lu, X.; Yao, B. A Novel Mathematical Method for Evaluating the Wellbore Deformation of a Diagenetic Natural Gas Hydrate Reservoir Considering the Effect of Natural Gas Hydrate Decomposition. *Nat. Gas Ind. B* **2021**. <https://doi.org/10.1016/j.ngib.2021.04.009>.
103. Ulyantsev, A.; Polyakova, N.; Trukhin, I.; Parotkina, Y.; Dudarev, O.; Semiletov, I. Peculiarities of Pore Water Ionic Composition in the Bottom Sediments and Subsea Permafrost: A Case Study in the Buor-Khaya Bay. *Geosciences* **2022**, *12*, 2, 1–49. <https://doi.org/10.3390/geosciences12020049>.
104. Chuvilin, E. M.; Ershov, E. D.; Naletova, N. S. Mass Transfer and Structure Formation in Freezing Saline Soils. In *7th International Conference on Permafrost*; Yellowknife, Canada, 1998; pp 173–179.
105. Iakovlev, A. A.; Turitsyna, M. V.; Mogil'nikov, E. V. Analysis and Justification of Selecting Cleaning Agents and Technology of Their Application in Permafrost Well Drilling (Published in Russian). *Bull. PNIPU. Geol. Oil gas Min.* **2014**, *12*, 22–32. <https://doi.org/10.15593/2224-9923/2014.12.3>.

106. Khimenkov, A. N.; Sergeev, D. O.; Vlasov, A. N.; Volkov-Bogorodsky, D. B.; Tipenko, G. S.; Merzlyakov, V. P.; Stanilovskaya, Y. V. Explosive Processes in Permafrost Areas – New Type of Geocryological Hazard. In *Innovation and Discovery in Russian Science and Engineering*; Springer, Cham, 2021; pp 83–99. https://doi.org/10.1007/978-3-030-63571-8_5.
107. Merey, Ş. Drilling of Gas Hydrate Reservoirs. *Journal of Natural Gas Science and Engineering*. 2016, pp 1167–1179. <https://doi.org/10.1016/j.jngse.2016.09.058>.
108. Ning, F.; Zhang, K.; Wu, N.; Zhang, L.; Li, G.; Jiang, G.; Yu, Y.; Liu, L.; Qin, Y. Invasion of Drilling Mud into Gas-Hydrate-Bearing Sediments. Part i: Effect of Drilling Mud Properties. *Geophys. J. Int.* **2013**, *193*, 3, 1370–1384. <https://doi.org/10.1093/gji/ggt015>.
109. Yakutseni, V. P. Gas Hydrates - Unconventional Gas Raw Materials, Their Formation, Properties, Distribution and Geological Resources (Published in Russian). *Oil gas Geol. Theory Pract.* **2013**, *8*, 4.
110. Bogoyavlensky, V. I. Natural and Technogenic Threats in Fossil Fuels Production in the Earth Cryolithosphere (Published in Russian). *Min. Ind. J. (Gornay Promishlennost)* **2020**, *1*, 97–118. <https://doi.org/10.30686/1609-9192-2020-1-97-118>.
111. Are, F. E. Problem of Deep Gas Emission into the Atmosphere (Published in Russian). *Kriosf. Zemli* **1998**, *2*, 4, 42–50.
112. Rivkin, F. M. Methane in Frozen Rocks and the Forecast of Its Release during Climate Warming and Technogenic Surface Disturbances (Published in Russian). *Proc. Russ. Acad. Sci. Geogr. Ser. Sci. J.* **1998**, *2*, 64–75.
113. Sergienko, V. I.; Lobkovskii, L. I.; Semiletov, I. P.; Dudarev, O. V.; Dmitrievskii, N. N.; Shakhova, N. E.; Romanovskii, N. N.; Kosmach, D. A.; Nikol'skii, D. N.; Nikiforov, S. L.; et

al. The Degradation of Submarine Permafrost and the Destruction of Hydrates on the Shelf of East Arctic Seas as a Potential Cause of the Methane Catastrophe: Some Results of Integrated Studies in 2011. *Dokl. Earth Sci.* **2012**, *446*, *1*, 1132–1137.

<https://doi.org/10.1134/S1028334X12080144>.

114. Anisimov, O. A.; Zaboikina, Y. G.; Kokorev, V. A.; Yurganov, L. N. Possible Causes of Methane Release from the East Arctic Seas Shelf (Published in Russian). *Ice Snow* **2014**, *2*, *126*, 69–81.
115. Dyadin, Y. A.; Gushchin, A. L. Gas Hydrates (Published in Russian). *Soros Educ. J.* **1998**, *3*, 55–64.
116. Anisimov, O. A.; Lavrov, S. A.; Reneva, S. A. Methane Emission from Permafrost Swamps in Russia under Climate Change (Published in Russian). In *Problems of Ecological Modeling and Monitoring of Ecosystems*; Israel, S. A., Ed.; Gidrometeoizdat: Saint Petersburg, Russia, 2005; pp 124–142.
117. Bazhin, N. M. Emission of Methane from Sedimentary Layers (Published in Russian). In *Reducing Methane Emissions*; 2000; pp 239–244.
118. Sergienko, V. I.; Semiletov, I. P.; Shakhova, N. E. Emission of Methane and Carbon Dioxide on the East Siberian Shelf Is a Factor of Global Climate Change (Published in Russian). In *Proceedings of the meeting of the Scientific Council of the Russian Academy of Sciences for the Study of the Arctic and Antarctic, 31 March - 2 April*; Arkhangelsk, Russia, 2010; pp 117–136.
119. Perlova, E. V.; Miklyaeva, E. S.; Leonov, S. A.; Tkacheva, E. V.; Ukhova, Y. A. Gas Hydrates within the Yamal Peninsular and Adjoining Kara Sea Continental Shelf as a Negative Factor of the Yamal Region Exploration (Published in Russian). *News gas Sci.*

2017, 3, 31, 255–262.

120. Collett, T. S.; Johnson, A. H.; Knapp, C. C.; Boswell, R. Natural Gas Hydrates: A Review. *AAPG Mem.* **2009**, 89, 146–219. <https://doi.org/10.1306/13201101M891602>.
121. Kvenvolden, K. A. Gas Hydrates-Geological Perspective and Global Change. *Rev. Geophys.* **1993**, 31, 2, 173–187. <https://doi.org/10.1029/93RG00268>.
122. Moridis, G.; Collett, T. Strategies for Gas Production from Hydrate Accumulations under Various Geologic Conditions. *Proc. TOUGH Symp. Lawrence Berkeley Natl. Lab. Berkeley, California, May 12–14* **2003**.
123. Makogon, Y. F.; Holditch, S. A.; Makogon, T. Y. Natural Gas-Hydrates - A Potential Energy Source for the 21st Century. *J. Pet. Sci. Eng.* **2007**, 56, 1–3, 14–31. <https://doi.org/10.1016/j.petrol.2005.10.009>.
124. Liang, Y.; Tan, Y.; Luo, Y.; Zhang, Y.; Li, B. Progress and Challenges on Gas Production from Natural Gas Hydrate-Bearing Sediment. *J. Clean. Prod.* **2020**, 261, 261:1-261:23. <https://doi.org/10.1016/j.jclepro.2020.121061>.
125. Nair, V. C.; Gupta, P.; Sangwai, J. S. Gas Hydrates as a Potential Energy Resource for Energy Sustainability; 2018; pp 265–287. https://doi.org/10.1007/978-981-10-7188-1_12.
126. Koh, D. Y.; Kang, H.; Lee, J. W.; Park, Y.; Kim, S. J.; Lee, J.; Lee, J. Y.; Lee, H. Energy-Efficient Natural Gas Hydrate Production Using Gas Exchange. *Appl. Energy* **2016**, 162, 114–130. <https://doi.org/10.1016/j.apenergy.2015.10.082>.
127. Makogon, Y. F.; Omelchenko, R. Y. Commercial Gas Production from Messoyakha Deposit in Hydrate Conditions. *J. Nat. Gas Sci. Eng.* **2013**, 11, 1–6. <https://doi.org/10.1016/j.jngse.2012.08.002>.

128. Choi, W.; Mok, J.; Lee, Y.; Lee, J.; Seo, Y. Optimal Driving Force for the Dissociation of CH₄ Hydrates in Hydrate-Bearing Sediments Using Depressurization. *Energy* **2021**, *223*,. <https://doi.org/10.1016/J.ENERGY.2021.120047>.
129. Fereidounpour, A.; Vatani, A. An Investigation of Interaction of Drilling Fluids with Gas Hydrates Indrilling Hydrate Bearing Sediments. *J. Nat. Gas Sci. Eng.* **2014**, *20*, 422–427. <https://doi.org/10.1016/j.jngse.2014.07.006>.
130. Huang, T.-J.; Zhang, Y.; Wang, Y.; Li, X.-S.; Chen, Z.-Y. An Experimental Drilling Apparatus Used for Evaluating Drilling Risks Related to Natural Gas Hydrate. *MethodsX* **2020**, *7*, 101019. <https://doi.org/10.1016/j.mex.2020.101019>.
131. Barker, J. W.; Gomez, R. K. *Formation of Hydrates during Deepwater Drilling Operations*; 1996; p. 105–109. <https://doi.org/10.2118/16130-pa>.
132. Priest, J. A.; Grozic, J. L. H. Stability of Fine-Grained Sediments Subject to Gas Hydrate Dissociation in the Arctic Continental Margin. In *Advances in Natural and Technological Hazards Research*; Springer Netherlands, 2016; Vol. 41, pp 427–436. https://doi.org/10.1007/978-3-319-20979-1_43.
133. Stackelberg, M. V.; Müller, H. R. On the Structure of Gas Hydrates. *The Journal of Chemical Physics*. 1951, pp 1319–1320. <https://doi.org/10.1063/1.1748038>.
134. Carroll, J. *Natural Gas Hydrates - A Guide for Engineers (3rd Edition)*; 2014; ISBN 978-0-12-800074-8.
135. Istomin, V.; Yakushev, V.; Makhonina, N. Self-Preservation Phenomenon of Gas Hydrate (Published in Russian). *Gas Ind.* **2006**, *4*, 36–46.
136. Koh, C. A.; Sloan, E. D. Natural Gas Hydrates: Recent Advances and Challenges in Energy

- and Environmental Applications. *AIChE J.* **2007**, *53*, 7, 1636–1643.
<https://doi.org/10.1002/aic.11219>.
137. Koh, C. A.; Sum, A. K.; Sloan, E. D. State of the Art: Natural Gas Hydrates as a Natural Resource. *J. Nat. Gas Sci. Eng.* **2012**, *8*, 132–138.
<https://doi.org/10.1016/j.jngse.2012.01.005>.
138. Makogon, Y. F. Natural Gas Hydrates - A Promising Source of Energy. *Journal of Natural Gas Science and Engineering*. 2010, pp 49–59. <https://doi.org/10.1016/j.jngse.2009.12.004>.
139. Supak, K. R.; George, D. L. Best Practices for Mitigating Flow Measurement Errors Associated with Hydrate Formation. *Pipeline&Gas J.* **2015**, *242*, 7.
140. Makogon, Y. F. *Hydrates of Natural Gas (Published in Russian)*; Nedra: Moscow, Russia, 1974; p. 208, ISBN 978-364-214-233-8.
141. Galloway, T. J.; Ruska, W.; Chappellear, P. S.; Kobayashi, R. Experimental Measurement of Hydrate Numbers for Methane and Ethane and Comparison with Theoretical Values. *Ind. Eng. Chem. Fundam.* **1970**, *9*, 2, 237–243. <https://doi.org/10.1021/i160034a008>.
142. Sum, A. K.; Burruss, R. C.; Sloan, E. D. Measurement of Clathrate Hydrates via Raman Spectroscopy. *J. Phys. Chem. B* **1997**, *101*, 38, 7371–7377.
<https://doi.org/10.1021/jp970768e>.
143. Groysman, A. G. *Thermophysical Properties of Gas Hydrates*; Nauka: Novosibirsk, 1985; p. 94.
144. Chong, Z. R.; Yang, S. H. B.; Babu, P.; Linga, P.; Li, X. Sen. Review of Natural Gas Hydrates as an Energy Resource: Prospects and Challenges. *Appl. Energy* **2016**, *162*, 1633–1652. <https://doi.org/10.1016/j.apenergy.2014.12.061>.

145. Hassanpouryouzband, A.; Joonaki, E.; Vasheghani Farahani, M.; Takeya, S.; Ruppel, C.; Yang, J.; English, N. J.; Schicks, J. M.; Edlmann, K.; Mehrabian, H.; et al. Gas Hydrates in Sustainable Chemistry. *Chem. Soc. Rev.* **2020**, *49*, *15*, 5225–5309.
<https://doi.org/10.1039/C8CS00989A>.
146. Collett, T. S.; Ginsburg, G. D. Gas Hydrates in the Messoyakha Gas Field of the West Siberian Basin - A Re-Examination of the Geologic Evidence. *Int. J. Offshore Polar Eng.* **1998**, *8*, *1*, 22–29.
147. Wang, X. H.; Sun, C. Y.; Chen, G. J.; He, Y. N.; Sun, Y. F.; Wang, Y. F.; Li, N.; Zhang, X. X.; Liu, B.; Yang, L. Y. Influence of Gas Sweep on Methane Recovery from Hydrate-Bearing Sediments. *Chem. Eng. Sci.* **2015**, *134*, 727–736. <https://doi.org/10.1016/j.ces.2015.05.043>.
148. Boswell, R.; Bünz, S.; Collett, T. S.; Frye, M.; Fujii, T.; McConnell, D.; Mienert, J.; Pecher, I.; Reichel, T.; Ryu, B.-J.; et al. Introduction to Special Section: Exploration and Characterization of Gas Hydrates. *Interpretation* **2016**, *4*, *1*, 2. <https://doi.org/10.1190/int-2016-0103-spseintro.1>.
149. Vasiliev, V. G.; Makogon, Y. F.; Trebin, F. A.; Trofimuk, A. A.; Chersky, N. V. The Property of Natural Gases to Be in a Solid State in the Earth's Crust (Published in Russian), 1970.
150. Johnson, A.; Max, M. Technology Options to Enable Low Environmental Risk Production of Natural Gas Hydrate. *9th Int. Conf. Gas Hydrates* **2017**.
151. Stern, L. A.; Kirby, S. H.; Circone, S.; Durham, W. B. Scanning Electron Microscopy Investigations of Laboratory-Grown Gas Clathrate Hydrates Formed from Melting Ice, and Comparison to Natural Hydrates Stern et Al. *Am. Mineral.* **2004**, *89*, 8–9, 1162–1175.
<https://doi.org/10.2138/am-2004-8-902>.

152. Kawamura, T.; Ohtake, M.; Yamamoto, Y.; Haneda, H.; Sakamoto, Y.; Komai, T.; Higuchi, S. Investigation For High Concentration Inhibitor Injection Into Artificial Methane Hydrate Bearing Sediment. **2009**, 119–123.
153. Zhao, X.; Qiu, Z.; Zhao, C.; Xu, J.; Zhang, Y. Inhibitory Effect of Water-Based Drilling Fluid on Methane Hydrate Dissociation. *Chem. Eng. Sci.* **2019**.
<https://doi.org/10.1016/j.ces.2018.12.057>.
154. Khan, M. S.; Lal, B.; Bustam, M. A. Gas Hydrate Inhibitors. In *Green Energy and Technology*; 2020; pp 27–46. https://doi.org/10.1007/978-3-030-30750-9_2.
155. Sira, J.; Patil, S.; Kamath, V. A. Study of Hydrate Dissociation by Methanol and Glycol Injection. In *Proceedings - SPE Annual Technical Conference and Exhibition*; 1990; Vol. Pi.
<https://doi.org/10.2118/20770-MS>.
156. Halliday, W.; Clapper, D. K.; Smalling, M. New Gas Hydrate Inhibitors for Deepwater Drilling Fluids. In *Proceedings of the IADC/SPE Asia Pacific Drilling Technology Conference, APDT*; 1998; pp 201–211. <https://doi.org/10.2118/1198-0052-jpt>.
157. Chong, Z. R.; Yin, Z.; Tan, J. H. C.; Linga, P. Experimental Investigations on Energy Recovery from Water-Saturated Hydrate Bearing Sediments via Depressurization Approach. *Appl. Energy* **2017**, *204*, 1513–1525. <https://doi.org/10.1016/j.apenergy.2017.04.031>.
158. Zhao, J.; Zhu, Z.; Song, Y.; Liu, W.; Zhang, Y.; Wang, D. Analyzing the Process of Gas Production for Natural Gas Hydrate Using Depressurization. *Appl. Energy* **2015**, *142*, 125–134. <https://doi.org/10.1016/j.apenergy.2014.12.071>.
159. Yang, X.; Sun, C. Y.; Su, K. H.; Yuan, Q.; Li, Q. P.; Chen, G. J. A Three-Dimensional Study on the Formation and Dissociation of Methane Hydrate in Porous Sediment by

- Depressurization. *Energy Convers. Manag.* **2012**, *56*, 1–7.
<https://doi.org/10.1016/j.enconman.2011.11.006>.
160. Heeschen, K. U.; Abendroth, S.; Priegnitz, M.; Spangenberg, E.; Thaler, J.; Schicks, J. M. Gas Production from Methane Hydrate: A Laboratory Simulation of the Multistage Depressurization Test in Mallik, Northwest Territories, Canada. *Energy and Fuels* **2016**, *30*, 8, 6210–6219. <https://doi.org/10.1021/acs.energyfuels.6b00297>.
161. Duan, Z.; Li, D.; Chen, Y.; Sun, R. The Influence of Temperature, Pressure, Salinity and Capillary Force on the Formation of Methane Hydrate. *Geosci. Front.* **2011**, *2*, 2, 125–135.
<https://doi.org/10.1016/j.gsf.2011.03.009>.
162. Dallimore, S. R.; Collett, T. S. Scientific Results from the Mallik 2002 Gas Hydrate Production Research Well Program Mackenzie Delta, Northwest Territories, Canada. **2002**, 1–36.
163. Yang, J.; Okwananke, A.; Tohidi, B.; Chuvilin, E.; Maerle, K.; Istomin, V.; Bukhanov, B.; Cheremisin, A. Flue Gas Injection into Gas Hydrate Reservoirs for Methane Recovery and Carbon Dioxide Sequestration. *Energy Convers. Manag.* **2017**, *136*, 431–438.
<https://doi.org/10.1016/j.enconman.2017.01.043>.
164. Saw, V. K.; Das, B. B.; Ahmad, I.; Mandal, A.; Laik, S. Influence of Electrolytes on Methane Hydrate Formation and Dissociation. *Energy Sources, Part A Recover. Util. Environ. Eff.* **2014**, *36*, 15, 1659–1669. <https://doi.org/10.1080/15567036.2011.557695>.
165. Cherskiy, N. V.; Groysman, A. G.; Nikitina, L. M.; Tserev, V. P. First Experimental Determination of Heats of Decomposition of Natural-Gas Hydrates. *Dokl. Acad. Sci. USSR, Earth Sci. Sect.* **1984**, *265*, 163–167.

166. Handa, P. Y. A Calorimetric Study of Naturally Occurring Gas Hydrates. *Ind. Eng. Chem. Res.* **1988**, *27*, *5*, 872–874. <https://doi.org/10.1021/ie00077a026>.
167. Misyura, S. Y. The Influence of Porosity and Structural Parameters on Different Kinds of Gas Hydrate Dissociation. *Sci. Rep.* **2016**, *6*, *1*, 30324. <https://doi.org/10.1038/srep30324>.
168. Rehder, G.; Eckl, R.; Elfgen, M.; Falenty, A.; Hamann, R.; Kähler, N.; Kuhs, W. F.; Osterkamp, H.; Windmeier, C. Methane Hydrate Pellet Transport Using the Self-Preservation Effect: A Techno-Economic Analysis. *Energies 2012, Vol. 5, Pages 2499-2523* **2012**, *5*, *7*, 2499–2523. <https://doi.org/10.3390/EN5072499>.
169. Feng, J.-C.; Li, G.; Li, X.-S.; Li, B.; Chen, Z.-Y. Evolution of Hydrate Dissociation by Warm Brine Stimulation Combined Depressurization in the South China Sea. *Energies* **2013**, *6*, *10*, 5402–5425. <https://doi.org/10.3390/en6105402>.
170. Grigg, R. B.; Lynes, G. L. Oil-Based Drilling Mud as a Gas-Hydrates Inhibitor. *SPE Drill. Eng.* **1992**, *7*, *1*, 32–38. <https://doi.org/10.2118/19560-PA>.
171. Qi, Y.; Wu, W.; Liu, Y.; Xie, Y.; Chen, X. The Influence of NaCl Ions on Hydrate Structure and Thermodynamic Equilibrium Conditions of Gas Hydrates. *Fluid Phase Equilib.* **2012**, *325*, 6–10. <https://doi.org/10.1016/J.FLUID.2012.04.009>.
172. Sun, S. C.; Kong, Y. Y.; Zhang, Y.; Liu, C. L. Phase Equilibrium of Methane Hydrate in Silica Sand Containing Chloride Salt Solution. *J. Chem. Thermodyn.* **2015**, *90*, 116–121. <https://doi.org/10.1016/j.jct.2015.06.030>.
173. Masoudi, R.; Tohidi, B. On Modelling Gas Hydrate Inhibition by Salts and Organic Inhibitors. *J. Pet. Sci. Eng.* **2010**, *74*, 3–4, 132–137. <https://doi.org/10.1016/J.PETROL.2010.08.010>.

174. Zatsepina, O. Y.; Buffett, B. A. Thermodynamic Conditions for the Stability of Gas Hydrate in the Seafloor. *J. Geophys. Res. Solid Earth* **1998**, *103*, *10*, 24127–24139. <https://doi.org/10.1029/98jb02137>.
175. Najibi, H.; Mohammadi, A. H.; Tohidi, B. Estimating the Hydrate Safety Margin in the Presence of Salt and/or Organic Inhibitor Using Freezing Point Depression Data of Aqueous Solutions. *Ind. Eng. Chem. Res.* **2006**, *45*, *12*, 4441–4446. <https://doi.org/10.1021/ie051265v>.
176. Cha, M.; Hu, Y.; Sum, A. K. Methane Hydrate Phase Equilibria for Systems Containing NaCl, KCl, and NH₄Cl. *Fluid Phase Equilib.* **2016**, *413*, 2–9. <https://doi.org/10.1016/j.fluid.2015.08.010>.
177. Kharrat, M.; Dalmazzone, D. Experimental Determination of Stability Conditions of Methane Hydrate in Aqueous Calcium Chloride Solutions Using High Pressure Differential Scanning Calorimetry. *J. Chem. Thermodyn.* **2003**, *35*, *9*, 1489–1505. [https://doi.org/10.1016/S0021-9614\(03\)00121-6](https://doi.org/10.1016/S0021-9614(03)00121-6).
178. De Roo, J. L.; Peters, C. J.; Lichtenthaler, R. N.; Diepen, G. A. M. Occurrence of Methane Hydrate in Saturated and Unsaturated Solutions of Sodium Chloride and Water in Dependence of Temperature and Pressure. *AIChE J.* **1983**, *29*, *4*, 651–657. <https://doi.org/10.1002/aic.690290420>.
179. Hu, Y.; Lee, B. R.; Sum, A. K. Universal Correlation for Gas Hydrates Suppression Temperature of Inhibited Systems: I. Single Salts. *AIChE J.* **2017**, *63*, *11*, 5111–5124. <https://doi.org/10.1002/aic.15846>.
180. Zeng, H.; Zhang, Y.; Li, X.; Chen, C.; Zhang, L.; Chen, Z. Experimental Study on the Influence of Brine Concentration on the Dissociation Characteristics of Methane Hydrate. *J. Nat. Gas Sci. Eng.* **2022**, *100*, 104492. <https://doi.org/10.1016/J.JNGSE.2022.104492>.

181. Jager, M. D.; Sloan, E. D. The Effect of Pressure on Methane Hydration in Pure Water and Sodium Chloride Solutions. *Fluid Phase Equilib.* **2001**, *185*, 1–2, 89–99.
[https://doi.org/10.1016/S0378-3812\(01\)00459-9](https://doi.org/10.1016/S0378-3812(01)00459-9).
182. Zheng, R.; Li, X.; Negahban, S. Phase Boundary of Gas Hydrates in Single and Mixed Electrolyte Solutions: Using a Novel Unified Equation of State. *J. Mol. Liq.* **2022**, *345*, 117825:1-117825:13. <https://doi.org/10.1016/j.molliq.2021.117825>.
183. Seo, Y.; Lee, H. Hydrate Phase Equilibria of the Ternary CH₄ + NaCl + Water, CO₂ + NaCl + Water and CH₄ + CO₂ + Water Mixtures in Silica Gel Pores. *J. Phys. Chem. B* **2003**, *107*, 3, 889–894. <https://doi.org/10.1021/jp026776z>.
184. Semenov, A. P.; Stoporev, A. S.; Mendgaziev, R. I.; Gushchin, P. A.; Khlebnikov, V. N.; Yakushev, V. S.; Istomin, V. A.; Sergeeva, D. V.; Vinokurov, V. A. Synergistic Effect of Salts and Methanol in Thermodynamic Inhibition of SII Gas Hydrates. *J. Chem. Thermodyn.* **2019**, *137*, 119–130. <https://doi.org/10.1016/j.jct.2019.05.013>.
185. Clennell, M. Ben; Hovland, M.; Booth, J. S.; Henry, P.; Winters, W. J. Formation of Natural Gas Hydrates in Marine Sediments: 1. Conceptual Model of Gas Hydrate Growth Conditioned by Host Sediment Properties. *J. Geophys. Res. Solid Earth* **1999**, *104*, B10, 22985–23003. <https://doi.org/10.1029/1999JB900175>.
186. Kumar, N.; Chowdhury, N. B.; Beltran, J. G. A 3-In-1 Approach to Evaluate Gas Hydrate Inhibitors. *Energies* **2019**, *12*, 15, 2921. <https://doi.org/10.3390/en12152921>.
187. Englezos, P.; Bishnoi, P. R. Prediction of Gas Hydrate Formation Conditions in Aqueous Electrolyte Solutions. *AIChE J.* **1988**, *34*, 10, 1718–1721.
<https://doi.org/10.1002/aic.690341017>.

188. Maekawa, T.; Itoh, S.; Sakata, S.; Igari, S. I.; Imai, N. Pressure and Temperature Conditions for Methane Hydrate Dissociation in Sodium Chloride Solutions. *Geochem. J.* **1995**, *29*, 5, 325–329.
189. Li, S.; Wang, J.; Lv, X.; Ge, K.; Jiang, Z.; Li, Y. Experimental Measurement and Thermodynamic Modeling of Methane Hydrate Phase Equilibria in the Presence of Chloride Salts. *Chem. Eng. J.* **2020**, *395*, 125126. <https://doi.org/10.1016/j.cej.2020.125126>.
190. Dickens, G. R.; Quinby-Hunt, M. S. Methane Hydrate Stability in Seawater. *Geophys. Res. Lett.* **1994**, *21*, 19, 2115–2118. <https://doi.org/10.1029/94GL01858>.
191. Dholabhai, P. D.; Englezos, P.; Kalogerakis, N.; Bishnoi, P. R. Equilibrium Conditions for Methane Hydrate Formation in Aqueous Mixed Electrolyte Solutions. *Can. J. Chem. Eng.* **1991**, *69*, 3, 800–805. <https://doi.org/10.1002/CJCE.5450690324>.
192. Choi, W.; Lee, J.; Kim, Y.-G.; Kim, H.; Rhee, T. S.; Jin, Y. K.; Kim, J.-H.; Seo, Y. The Impact of the Abnormal Salinity Enrichment in Pore Water on the Thermodynamic Stability of Marine Natural Gas Hydrates in the Arctic Region. *Sci. Total Environ.* **2021**, *799*, 149357:1-149357:9. <https://doi.org/10.1016/j.scitotenv.2021.149357>.
193. Li, S.; Zheng, R.; Xu, X.; Chen, Y. Dissociation of Methane Hydrate by Hot Brine. *Pet. Sci. Technol.* **2015**, *33*, 6, 671–677. <https://doi.org/10.1080/10916466.2015.1005845>.
194. Ahn, T.; Park, C.; Lee, J.; Kang, J. M.; Nguyen, H. T. Experimental Characterization of Production Behaviour Accompanying the Hydrate Reformation in Methane-Hydrate-Bearing Sediments. *J. Can. Pet. Technol.* **2012**, *51*, 01, 14–19. <https://doi.org/10.2118/136737-PA>.
195. Jin, Y.; Li, S.; Yang, D.; Jiang, X. Determination of Dissociation Front and Operational Optimization for Hydrate Development by Combining Depressurization and Hot Brine

- Stimulation. *J. Nat. Gas Sci. Eng.* **2018**, *50*, 215–230.
<https://doi.org/10.1016/j.jngse.2017.12.009>.
196. Xu, J.; Chen, Z.; Liu, J.; Sun, Z.; Wang, X.; Zhang, J. A Molecular Dynamic Study on the Dissociation Mechanism of SI Methane Hydrate in Inorganic Salt Aqueous Solutions. *J. Mol. Graph. Model.* **2017**, *75*, 403–412. <https://doi.org/10.1016/j.jmgm.2017.03.022>.
197. Ding, T.; Wang, R.; Xu, J.; Camara, M.; Zhou, W.; Zhang, J. Dissociation Mechanism of Methane Hydrate by CaCl₂: An Experimental and Molecular Dynamics Study. *J. Mol. Model.* **2022**, *28*, 4, 109. <https://doi.org/10.1007/s00894-022-05070-6>.
198. Sung, W.; Kang, H. Experimental Investigation of Production Behaviors of Methane Hydrate Saturated in Porous Rock. *Energy Sources* **2003**, *25*, 8, 845–856.
<https://doi.org/10.1080/00908310390207873>.
199. Li, X.-S.; Wan, L.-H.; Li, G.; Li, Q.-P.; Chen, Z.-Y.; Yan, K.-F. Experimental Investigation into the Production Behavior of Methane Hydrate in Porous Sediment with Hot Brine Stimulation. *Ind. Eng. Chem. Res.* **2008**, *47*, 23, 9696–9702.
<https://doi.org/10.1021/ie8009582>.
200. Xu, J.; Gu, T.; Sun, Z.; Li, X.; Wang, X. Molecular Dynamics Study on the Dissociation of Methane Hydrate via Inorganic Salts. *Mol. Phys.* **2016**, *114*, 1, 34–43.
<https://doi.org/10.1080/00268976.2015.1081708>.
201. Chen, J.; Liu, C.; Zhang, Z.; Wu, N.; Liu, C.; Ning, F.; Fang, B.; Wan, Y.; Bu, Q.; Hu, G. Molecular Study on the Behavior of Methane Hydrate Decomposition Induced by Ions Electrophoresis. *Fuel* **2022**, *307*, 121866:1-121866:7.
<https://doi.org/10.1016/j.fuel.2021.121866>.

202. Mohammadi, A. H.; Afzal, W.; Richon, D. Gas Hydrates of Methane, Ethane, Propane, and Carbon Dioxide in the Presence of Single NaCl, KCl, and CaCl₂ Aqueous Solutions: Experimental Measurements and Predictions of Dissociation Conditions. *J. Chem. Thermodyn.* **2008**, *40*, 12, 1693–1697. <https://doi.org/10.1016/j.jct.2008.06.015>.
203. Aregbe, A.; Sun, B.; Engineering, L. C.-J. of C. & 2019, undefined. Methane Hydrate Dissociation Conditions in High-Concentration NaCl/KCl/CaCl₂ Aqueous Solution: Experiment and Correlation. *ACS Publ.* **2019**, *64*, 7, 2929–2939. <https://doi.org/10.1021/acs.jced.8b01173>.
204. Atik, Z.; Windmeier, C.; Oellrich, L. R. Experimental Gas Hydrate Dissociation Pressures for Pure Methane in Aqueous Solutions of MgCl₂ and CaCl₂ and for a (Methane + Ethane) Gas Mixture in an Aqueous Solution of (NaCl + MgCl₂). *J. Chem. Eng. Data* **2006**, *51*, 5, 1862–1867. <https://doi.org/10.1021/je060225a>.
205. Chong, Z. R.; Chan, A. H. M.; Babu, P.; Yang, M.; Linga, P. Effect of NaCl on Methane Hydrate Formation and Dissociation in Porous Media. *J. Nat. Gas Sci. Eng.* **2015**, *27*, 178–189. <https://doi.org/10.1016/J.JNGSE.2015.08.055>.
206. Stoporev, A. S.; Semenov, A. P.; Medvedev, V. I.; Kidyarov, B. I.; Manakov, A. Y.; Vinokurov, V. A. Nucleation of Gas Hydrates in Multiphase Systems with Several Types of Interfaces. *J. Therm. Anal. Calorim.* **2018**, *134*, 1, 783–795. <https://doi.org/10.1007/s10973-018-7352-2>.
207. Li, H.; Wang, L. Hydrophobized Particles Can Accelerate Nucleation of Clathrate Hydrates. *Fuel* **2015**, *140*, 440–445. <https://doi.org/10.1016/j.fuel.2014.10.005>.
208. Prasad, P. S. R.; Kiran, B. S. Self-Preservation and Stability of Methane Hydrates in the Presence of NaCl. *Sci. Rep.* **2019**, *9*, 1, 5860. <https://doi.org/10.1038/s41598-019-42336-1>.

209. Henry, P.; Thomas, M.; Clennell, M. Ben. Formation of Natural Gas Hydrates in Marine Sediments: 2. Thermodynamic Calculations of Stability Conditions in Porous Sediments. *J. Geophys. Res. Solid Earth* **1999**, *104*, B10, 23005–23022.
<https://doi.org/10.1029/1999JB900167>.
210. Zaripova, Y.; Yarkovoi, V.; Varfolomeev, M.; Kadyrov, R.; Stoporev, A. Influence of Water Saturation, Grain Size of Quartz Sand and Hydrate-Former on the Gas Hydrate Formation. *Energies* **2021**, *14*, 5, 1272. <https://doi.org/10.3390/en14051272>.
211. Daigle, H.; Dugan, B. Capillary Controls on Methane Hydrate Distribution and Fracturing in Advective Systems. *Geochemistry, Geophys. Geosystems* **2011**, *12*, 1, Q01003:1-Q01003:18.
<https://doi.org/10.1029/2010GC003392>.
212. Smith, D. H.; Wilder, J. W.; Seshadri, K. Methane Hydrate Equilibria in Silica Gels with Broad Pore-Size Distributions. *AIChE J.* **2002**, *48*, 2, 393–400.
<https://doi.org/10.1002/aic.690480222>.
213. Anderson, R.; Llamedo, M.; Tohidi, B.; Burgass, R. W. Characteristics of Clathrate Hydrate Equilibria in Mesopores and Interpretation of Experimental Data. *J. Phys. Chem. B* **2003**, *107*, 15, 3500–3506. <https://doi.org/10.1021/jp0263368>.
214. Sergeeva, D.; Istomin, V.; Chuvilin, E.; Bukhanov, B.; Sokolova, N. Influence of Hydrate-Forming Gas Pressure on Equilibrium Pore Water Content in Soils. *Energies* **2021**, *Vol. 14*, Page 1841 **2021**, *14*, 7, 1841. <https://doi.org/10.3390/EN14071841>.
215. Wu, G.; Ji, H.; Tian, L.; Chen, D. Effects of Salt Ions on the Methane Hydrate Formation and Dissociation in the Clay Pore Water and Bulk Water. *Energy & Fuels* **2018**, *32*, 12, 12486–12494. <https://doi.org/10.1021/acs.energyfuels.8b03486>.

216. Bello-Palacios, A.; Almenningen, S.; Fotland, P.; Ersland, G. Experimental and Numerical Analysis of the Effects of Clay Content on CH₄ Hydrate Formation in Sand. *Energy & Fuels* **2021**, acs.energyfuels.1c00549. <https://doi.org/10.1021/acs.energyfuels.1c00549>.
217. Qin, Y.; Pan, Z.; Liu, Z.; Shang, L.; Zhou, L. Influence of the Particle Size of Porous Media on the Formation of Natural Gas Hydrate: A Review. *Energy & Fuels* **2021**, *35*, *15*, 11640–11664. <https://doi.org/10.1021/acs.energyfuels.1c00936>.
218. Kang, S.; Ryu, H.; Seo, Y. Phase Behavior of CO₂ and CH₄ Hydrate in Porous Media. *World Acad. Sci. Eng. Technol.* **2007**, *33*, 183–188.
219. Istomin, V.; Chuvilin, E.; Makhonina, N.; Kvon, V.; Safonov, S. Description of Gas Hydrates Equilibria in Sediments Using Experimental Data of Soil Water Potential. In *ICGH-2008, July 6-10; Vancouver, Canada, 2008*; pp 1–12.
220. Johnson, A. H. Global Resource Potential of Gas Hydrate – a New Calculation. *Nat. Gas Oil* **2011**, *304*, 285–4541.
221. Solov'yev, V. A.; Ginsburg, G. D. Formation of Submarine Gas Hydrates. *Bull. Geol. Soc. Denmark* **1994**, *41*, 86–94.
222. Okwananke, A.; Yang, J.; Tohidi, B.; Chuvilin, E.; Istomin, V.; Cheremisin, A. Compressed Air Injection for Methane Recovery from Gas Hydrate-Bearing Sediments. *Ninth Int. Conf. Gas Hydrates* **2017**.
223. Davletshina, D. A. Regularities of Formation and Dissociation of Gas Hydrates in Frozen Sediments (Published in Russian), Institute of permafrost, Yakutsk, 2021.
224. Romanovskii, N. N.; Hubberten, H. W.; Gavrilov, A. V.; Eliseeva, A. A.; Tipenko, G. S.; Kholodov, A. L.; Romanovsky, V. E. Permafrost and Gas Hydrate Stability Zone Evolution

on the Eastern Part of the Eurasia Arctic Sea Shelf in the Middle Pleistocene-Holocene
(Published in Russian). *Earth's Cryosph.* **2003**, 7, 4, 51–64.

225. Malakhova, V.; Golubeva, E. Model Study of the Effects of Climate Change on the Methane Emissions on the Arctic Shelves. *Atmosphere (Basel)*. **2022**, 13, 2, 274:1-274:21.
<https://doi.org/10.3390/atmos13020274>.
226. Chuvilin, E.; Davletshina, D. A.; Lupachik, M. Hydrate Formation in Frozen and Thawing Methane Saturated Sediments (Published in Russian). *Earth's Cryosph.* **2019**, 23, 2, 50–61.
227. Kvenvolden, K. A. Natural Gas Hydrate Occurrence and Issues. *Ann. N. Y. Acad. Sci.* **1994**, 715, 1, 232–246. <https://doi.org/10.1111/j.1749-6632.1994.tb38838.x>.
228. Sun, Y.; Goldberg, D.; Collett, T.; Hunter, R. High-Resolution Well-Log Derived Dielectric Properties of Gas-Hydrate-Bearing Sediments, Mount Elbert Gas Hydrate Stratigraphic Test Well, Alaska North Slope. *Mar. Pet. Geol.* **2011**, 28, 2, 450–459.
<https://doi.org/10.1016/j.marpetgeo.2010.03.001>.
229. Zhu, Y.; Zhang, Y.; Wen, H.; Lu, Z.; Jia, Z.; Li, Y.; Li, Q.; Liu, C.; Wang, P.; Guo, X. Gas Hydrates in the Qilian Mountain Permafrost, Qinghai, Northwest China. *Acta Geol. Sin. - English Ed.* **2010**, 84, 1, 1–10. <https://doi.org/10.1111/j.1755-6724.2010.00164.x>.
230. Liu, C.; Meng, Q.; He, X.; Li, C.; Ye, Y.; Lu, Z.; Zhu, Y.; Li, Y.; Liang, J. Comparison of the Characteristics for Natural Gas Hydrate Recovered from Marine and Terrestrial Areas in China. *J. Geochemical Explor.* **2015**, 152, 67–74.
<https://doi.org/10.1016/j.gexplo.2015.02.002>.
231. Lu, Z.; Zhu, Y.; Zhang, Y.; Wen, H.; Li, Y.; Liu, C. Gas Hydrate Occurrences in the Qilian Mountain Permafrost, Qinghai Province, China. *Cold Reg. Sci. Technol.* **2011**, 66, 2–3, 93–

104. <https://doi.org/10.1016/j.coldregions.2011.01.008>.
232. Lu, Z.; Zhu, Y.; Liu, H.; Zhang, Y.; Jin, C.; Huang, X.; Wang, P. Gas Source for Gas Hydrate and Its Significance in the Qilian Mountain Permafrost, Qinghai. *Mar. Pet. Geol.* **2013**, *43*, 341–348. <https://doi.org/10.1016/j.marpetgeo.2013.01.003>.
233. Golmohammadi, S. M.; Nakhaee, A. A Cylindrical Model for Hydrate Dissociation near Wellbore during Drilling Operations. *J. Nat. Gas Sci. Eng.* **2015**, *27*, 1641–1648. <https://doi.org/10.1016/j.jngse.2015.10.032>.
234. Agalakov, S. E.; Kurchikov, A. R.; Baburin, A. N. Geological and Geophysical Background the Existence of Gas Hydrates in the Turonian Deposits of the East Messoyakha Deposits (Published in Russian). *Geol. Geophys.* **2001**, *42*, 11–12, 1785–1791.
235. Andreassen, K.; Hubbard, A.; Winsborrow, M.; Patton, H.; Vadakkepuliambatta, S.; Plaza-faverola, A.; Deryabin, A.; Mattingsdal, R.; Mienert, J. Gas Hydrate Regulate Methane Emissions from Arctic Petroleum Basins. In *Proceeding of the 9th International Conference on Gas Hydrates (ICGH9)*; 2017; p 2.
236. Chuvilin, E.; Ekimova, V.; Davletshina, D.; Sokolova, N.; Bukhanov, B. Evidence of Gas Emissions from Permafrost in the Russian Arctic. *Geosciences* **2020**, *10*, 10, 383:1-383:23. <https://doi.org/10.3390/geosciences10100383>.
237. Anisimov, O. A.; Kokorev, V. A. Comparative Analysis of Land, Marine, and Satellite Observations of Methane in the Lower Atmosphere in the Russian Arctic under Conditions of Climate Change. *Izv. Atmos. Ocean. Phys.* **2015**, *51*, 9, 979–991. <https://doi.org/10.1134/s0001433815090030>.
238. Bogoyavlensky, V. I. Prospects and Problems of the Arctic Shelf Oil and Gas Fields

- Development (Published in Russian). *Drill. Oil* **2012**, *11*, 4–9.
239. Bogoyavlensky, V. I. Gas-Hydrodynamics in the Arctic Craters of Gas Blowout (Published in Russia). *Arct. Ecol. Econ.* **2018**, No. 1(29), 48–55. <https://doi.org/10.25283/2223-4594-2018-1-48-55>.
240. Bogoyavlensky, V. I.; Sizov, O. S.; Bogoyavlensky, I. V.; Nikonov, R. A. Remote Detection of Areas of Surface Gas Manifestations and Gas Emissions in the Arctic: The Yamal Peninsula (Published in Russian). *Arct. Ecol. Econ.* **2016**, *3*, 23, 4–15.
241. Sizov, O. S. Remote Analysis of the Consequences of Surface Gas Show in the North of Western Siberia (Published in Russian). *Geomatics* **2015**, *21*, *1*, 53–68.
242. Buldovicz, S. N.; Khilimonyuk, V. Z.; Bychkov, A. Y.; Ospennikov, E. N.; Vorobyev, S. A.; Gunar, A. Y.; Gorshkov, E. I.; Chuvilin, E. M.; Cherbunina, M. Y.; Kotov, P. I.; et al. Cryovolcanism on the Earth: Origin of a Spectacular Crater in the Yamal Peninsula (Russia). *Sci. Rep.* **2018**, *8*, *1*, 13534. <https://doi.org/10.1038/s41598-018-31858-9>.
243. Bogoyavlensky, V. I. Oil and Gas Emissions on Land and Offshore Areas of the Arctic and World Ocean (Published in Russian). *Drill. oil* **2015**, No. 6, 4–10.
244. Leibman, M. O.; Plekhanov, A. V. Yamal Gas Emission Crater: Results of Preliminary Survey (Published in Russian). *KholodOk* **2014**, *2*, 9–15.
245. Leibman, M. O.; Kizyakov, A. I.; Plekhanov, A. V.; Streletskaya, I. D. New Permafrost Feature – Deep Crater in Central Yamal (West Siberia, Russia) as a Response to Local Climate Fluctuations. *Geogr. Environ. Sustain.* **2014**, *7*, *4*, 68–79. <https://doi.org/10.24057/2071-9388-2014-7-4-68-79>.
246. Olenchenko, V. V.; Sinitsky, A. I.; Antonov, E. Y.; Eltsov, I. N.; Kushnarenko, O. N.;

- Plotnikov, A. E.; Potapov, V. V.; Epov, M. I. Results of Geophysical Researches of the Area of New Geological Formation “Yamal Crater” (Published in Russian). *Kriosf. Zemli* **2015**, *19*, *4*, 94–106.
247. Kizyakov, A. I.; Sonyushkin, A. V.; Leibman, M. O.; Zimin, M. V. Geomorphological Conditions of the Gas-Emission Crater and Its Dynamics in Central Yamal (In Russian). *Kriosf. Zemli* **2015**, *19*, *2*, 13–23.
248. Kizyakov, A.; Zimin, M.; Sonyushkin, A.; Dvornikov, Y.; Khomutov, A.; Leibman, M. Comparison of Gas Emission Crater Geomorphodynamics on Yamal and Gydan Peninsulas (Russia), Based on Repeat Very-High-Resolution Stereopairs. *Remote Sens.* **2017**, *9*, *10*, 1023. <https://doi.org/10.3390/rs9101023>.
249. Chuvilin, E.; Stanilovskaya, J.; Titovsky, A.; Sinitsky, A.; Sokolova, N.; Bukhanov, B.; Spasennykh, M.; Cheremisin, A.; Grebenkin, S.; Davletshina, D.; et al. A Gas-Emission Crater in the Erkuta River Valley, Yamal Peninsula: Characteristics and Potential Formation Model. *Geosciences* **2020**, *10*, *5*, 170. <https://doi.org/10.3390/geosciences10050170>.
250. Bogoyavlensky, V. I.; Bogoyavlensky, I. V.; Sizov, O. S.; Nikonov, R. A.; Kargina, T. N. Earth Degassing in the Arctic: Remote and Field Studies of the Thermokarst Lakes Gas Eruption (Published in Russian). *Arct. Ecol. Econ.* **2019**, No. 2(34), 31–47. <https://doi.org/10.25283/2223-4594-2019-2-31-47>.
251. Bogoyavlensky, V. I.; Erokhin, G. N.; Nikonov, R. A.; Bogoyavlensky, I. V.; Bryksin, V. M. Study of Catastrophic Gas Blowout Zones in the Arctic Based on Passive Microseismic Monitoring (on the Example of Lake Otkrytiye) (Published in Russian). *Arct. Ecol. Econ.* **2020**, *1*, *37*, 93–104. <https://doi.org/doi:10.25283/2223-4594-2020-1-93-104>.
252. Kalinko, M. K. *History of Geological Development and Oil and Gas Potential of the*

- Khataigekoy Depression (Published in Russian)*; Gostoptekhizdat: Leningrad, 1959; p. 358.
253. Mamzelev, A. P.; Are, F. E. Geological-Engineering Conditions of Yamal Peninsula along Designing Railroad. In *6th International Conference on Permafrost 5-9 July*; South China University of Technology Press: Beijing, China, 1993; pp 436–442.
254. Melnikov, P. I.; Melnikov, V. P.; Tsarev, V. P. On Generation of Hydrocarbons in the Thicknesses of the Permafrost Rocks (Published in Russian). *Izv. AN SSSR. Seriya Geol.* **1989**, 2, 118–128.
255. Badu, Y. B. The Cryolithic Structure of the Ground. In *The cryosphere of oil and gas condensate fields, Yamal Peninsula*; Nedra: Saint Petersburg, Russia, 2006; pp 85–111.
256. Chuvilin, E. M.; Perlova, E. V.; Baranov, Y. B.; Kondakov, V. V.; Osokin, A. B.; Yakushev, V. S. *Structure and Properties of Rocks in Cryolithozone of the Southern Part of Bovanenkovo Gas-Condensate Deposit (Published in Russian)*; GEOS: Moscow, Russia, 2007; p. 137.
257. Kondakov, V. V.; Kusova, O. F.; Kondakov, M. V. Geocryological Conditions of the North-Eastern Part of the Yamal Peninsula (Published in Russian). In *The fourth conference of geocryologists of Russia*; University book: Moscow, Russia, 2011; pp 89–94.
258. Bolshakov, Yu.Ya. Kultikov, A. M. *Analysis of Gas Showings during Drilling on the Territory of Bovanenkovo and Kharasavey Fields (Published in Russian): Report of the Institute of North Development Problems of the Siberian Branch of the USSR Academy of Sciences. N2 1*; Tyumen, 1989; p. 74.
259. Kondakov, V. V.; Galyavich, A. S. Comprehensive Studies of Permafrost Sediments with an Assessment of Their Water and Gas Saturation (Published in Russian). In *Problems of*

cryology of the Earth: abstracts. report conferences, 20-24 April; Pushchino, Russia, 1998; p 105.

260. Ivanov, M. S. Modern Permafrost Coastal Deltaic Sediments of the Yansky Seaside (Published in Russian). *Quest. Geogr. Yakutia* **1969**, 5, 138–147.
261. Masurenkov, Y. P.; Slezin, Y. B.; Sobisevich, A. L. Gas Plumes off Bennett’s Island (Published in Russian). *Bull. Russ. Acad. Sci. Geogr. Ser.* **2015**, 3, 86–95.
262. Rekant, P. V.; Tumskoy, V. E.; Gusev, E. A.; Schwenk, T.; Spiess, F.; Cherkashev, G. A.; Kassens, H. Distribution and Features of Occurrence of the Subaqueous Cryolithozone in the Area of the Semenovskaya and Vasilievskaya Banks (the Laptev Sea) According to Seismoacoustic Profiling (Published in Russian). In *System of the Laptev Sea and Adjacent Seas of the Arctic: Current State and History of development*; Moscow State University: Moscow, Russia, 2009; pp 292–308.
263. Shakhova, N. E.; Semiletov, I. P.; Salyuk, A. N.; Belcheva, N. N.; Kosmach, D. A. Anomalies of Methane in the Water Layer of the Atmosphere on the Shelf of the East Siberian Arctic (Published in Russian). *Reports Acad. Sci.* **2007**, 414, 6, 819–823.
264. Shakhova, N.; Semiletov, I.; Leifer, I.; Sergienko, V.; Salyuk, A.; Kosmach, D.; Chernykh, D.; Stubbs, C.; Nicolsky, D.; Tumskoy, V.; et al. Ebullition and Storm-Induced Methane Release from the East Siberian Arctic Shelf. *Nat. Geosci.* **2014**, 7, 1, 64–70.
<https://doi.org/10.1038/ngeo2007>.
265. Shakhova, N.; Semiletov, I.; Sergienko, V.; Lobkovsky, L.; Yusupov, V.; Salyuk, A.; Salomatin, A.; Chernykh, D.; Kosmach, D.; Panteleev, G.; et al. The East Siberian Arctic Shelf: Towards Further Assessment of Permafrost-Related Methane Fluxes and Role of Sea Ice. *Philos. Trans. R. Soc. A Math. Phys. Eng. Sci.* **2015**, 373, 2052:1-2052:13.

<https://doi.org/10.1098/rsta.2014.0451>.

266. Shakhova, N. E.; Sergienko, V. I.; Semiletov, I. P. The Contribution of the East Siberian Shelf to the Modern Methane Cycle. *Her. Russ. Acad. Sci.* **2009**, *79*, 3, 237–246.
<https://doi.org/10.1134/S101933160903006X>.
267. Baranov, B.; Galkin, S.; Vedenin, A.; Dozorova, K.; Gebruk, A.; Flint, M. Methane Seeps on the Outer Shelf of the Laptev Sea: Characteristic Features, Structural Control, and Benthic Fauna. *Geo-Marine Lett.* **2020**. <https://doi.org/10.1007/s00367-020-00655-7>.
268. Samarkin, V.; Semiletov, I. P.; Finke, N.; Shakhova, N. E.; Joye, S. B. Methane Carbon Stable Isotope Signatures in Waters and Sediments of the Laptev Sea Shelf. In *American Geophysical Union, 3-7 December*; San Francisco, USA, 2012; pp B21D-0411.
269. Weidner, E.; Weber, T. C.; Mayer, L.; Jakobsson, M.; Chernykh, D.; Semiletov, I. A Wideband Acoustic Method for Direct Assessment of Bubble-Mediated Methane Flux. *Cont. Shelf Res.* **2019**, *173*, 104–115. <https://doi.org/10.1016/j.csr.2018.12.005>.
270. Firsov, Y. G.; Ivanov, M. V.; Koloskov, E. N. A New Stage in Bathymetric Studies of the Northern Waters of Russia on the Example of the Kara Sea (Published in Russian). *Bull. State Univ. Mar. River Fleet Admiral Makarov* **2014**, *6*, 28, 115–124.
271. Koloskov, E. N.; Firsov, Y. G. The Use of Modern Hydrographic Technologies for the Study of Topography and Bottom Gas Occurrence in the Northern Seas of Russia (Published in Russian). *Bull. State Univ. Marit. River Fleet Admiral SB Makarov* **2015**, *3*, 31, 54–62.
272. Bondarev, V. N.; Rokos, S. I.; Kostin, D. A.; Dlugach, A. G.; Polyakova, N. A. Underpermafrost Accumulations of Gas in the Upper Part of the Sedimentary Cover of the Pechora Sea (Published in Russian). *Geol. Geophys.* **2002**, *43*, 7, 587–598.

273. Rokos, S. I. Engineering and Geological Features of Near-Surface Zones of Anomalously High Reservoir Pressure on the Shelf of the Pechora and Southern Kara Seas (Published in Russian). *Eng. Geol.* **2008**, *4*, 22–28.
274. Grigoriev, M. N. Studies of the Degradation of Permafrost Rocks in the Seas of Eastern Siberia (Based on the Results of Expeditions in 2014–2016) (Published in Russian). *Arct. Antarct. Probl.* **2018**, *1*, 89–96.
275. Safronov, A. F.; Shits, E. Y.; Grigor'ev, M. N.; Semenov, M. E. Formation of Gas Hydrate Deposits in the Siberian Arctic Shelf (Published in Russian). *Russ. Geol. Geophys.* **2010**, *51*, *1*, 83–87. <https://doi.org/10.1016/j.rgg.2009.12.006>.
276. Delisle, G. Temporal Variability of Subsea Permafrost and Gas Hydrate Occurrences as Function of Climate Change in the Laptev Sea, Siberia. *Polarforschung* **2000**, *68*, *1–3*, 221–225.
277. Shakhova, N.; Semiletov, I.; Salyuk, A.; Yusupov, V.; Kosmach, D.; Gustafsson, Ö. Extensive Methane Venting to the Atmosphere from Sediments of the East Siberian Arctic Shelf. *Science (80-.)*. **2010**, *327*, 5970, 1246–1250. <https://doi.org/10.1126/science.1182221>.
278. Nicolsky, D. J.; Romanovsky, V. E.; Romanovskii, N. N.; Kholodov, A. L.; Shakhova, N. E.; Semiletov, I. P. Modeling Sub-Sea Permafrost in the East Siberian Arctic Shelf: The Laptev Sea Region. *J. Geophys. Res. Earth Surf.* **2012**, *117*, *3*, F03028:1-F03028:22. <https://doi.org/10.1029/2012JF002358>.
279. Bondur, V. G.; Kuznetsova, T. V.; Vorobyev, V. E.; Zamshin, V. Detection of Gas Shows (Gas Seeps) on the Russian Shelf Using Satellite Data (Published in Russian). *Georesources. Geoenergy. Geopolit.* **2014**, *1*, *9*, 1–23.

280. Bondur, V. G.; Kuznetsova, T. V. Detecting Gas Seeps in Arctic Water Areas Using Remote Sensing Data. *Izv. Atmos. Ocean. Phys.* **2015**, *51*, 9, 1060–1072.
<https://doi.org/10.1134/S0001433815090066>.
281. Ananiev, R. A.; Dmitrevsky, N. N.; Roslyakov, A. G.; Chernykh, D. V.; Moroz, E. A.; Zarayskaya, Y. A.; Semiletov, I. P. Acoustic Monitoring of Gas Emission Processes in the Arctic Shelf Seas. *Oceanol.* **2022**, *62*, 1, 127–132.
<https://doi.org/10.1134/S0001437022010015>.
282. Bogoyavlensky, V.; Kishankov, A.; Kazanin, A.; Kazanin, G. Distribution of Permafrost and Gas Hydrates in Relation to Intensive Gas Emission in the Central Part of the Laptev Sea (Russian Arctic). *Mar. Pet. Geol.* **2022**, *138*, 105527.
<https://doi.org/10.1016/J.MARPETGEO.2022.105527>.
283. Anisimov, O. A.; Borzenkova, I. I.; Lavrov, S. A.; Strelchenko, Y. G. Modern Dynamics of Underwater Permafrost and Methane Emission on the Shelf of the Seas of the Eastern Arctic (Published in Russian). *Ice snow* **2012**, *2*, 97–105.
284. Ruppel, C. D.; Kessler, J. D. The Interaction of Climate Change and Methane Hydrates. *Rev. Geophys.* **2017**, *55*, 1, 126–168. <https://doi.org/10.1002/2016RG000534>.
285. Fedoseev, S. M. Gas Hydrates of Permafrost (Published in Russian). *Nat. Resour. Arct. Subarct.* **2006**, *1*, 22–27.
286. Ershov, E. D.; Lebedenko, Y. P.; Chuvilin, E. M.; Benediktova, N. A. Migration of Chemical Elements and Related Environmental Problems (Published in Russian). In *the III Republican scientific and technical seminar “Modern methods of exploration and development of minerals in the Far North”*; Syktyvkar, Russia, 1989; pp 26–30.

287. Hao, S. A Study to Optimize Drilling Fluids to Improve Borehole Stability in Natural Gas Hydrate Frozen Ground. *J. Pet. Sci. Eng.* **2011**, *76*, 3–4, 109–115.
<https://doi.org/10.1016/j.petrol.2011.01.014>.
288. Srungavarapu, M.; Patidar, K. K.; Pathak, A. K.; Mandal, A. Performance Studies of Water-Based Drilling Fluid for Drilling through Hydrate Bearing Sediments. *Appl. Clay Sci.* **2018**.
<https://doi.org/10.1016/j.clay.2017.11.014>.
289. Li, Q.; Liu, L.; Yu, B.; Guo, L.; Shi, S.; Miao, L. Borehole Enlargement Rate as a Measure of Borehole Instability in Hydrate Reservoir and Its Relationship with Drilling Mud Density. *J. Pet. Explor. Prod.* **2021**, *11*, 3, 1185–1198. <https://doi.org/10.1007/s13202-021-01097-2>.
290. Sun, S.; Ye, Y.; Liu, C.; Xiang, F.; Ma, Y. P-T Stability Conditions of Methane Hydrate in Sediment from South China Sea. *J. Nat. Gas Chem.* **2011**, *20*, 5, 531–536.
[https://doi.org/10.1016/S1003-9953\(10\)60224-1](https://doi.org/10.1016/S1003-9953(10)60224-1).
291. Lee, J. Experimental Study on the Dissociation Behavior and Productivity of Gas Hydrate by Brine Injection Scheme in Porous Rock. *Energy & Fuels* **2010**, *24*, 1, 456–463.
<https://doi.org/10.1021/ef900791r>.
292. Chong, Z. R.; Koh, J. W.; Linga, P. Effect of KCl and MgCl₂ on the Kinetics of Methane Hydrate Formation and Dissociation in Sandy Sediments. *Energy* **2017**, *137*, 518–529.
<https://doi.org/10.1016/j.energy.2017.01.154>.
293. Zheng, M.; Liu, T.; Jiang, G.; Wei, M.; Huo, Y.; Liu, L. Large-Scale and High-Similarity Experimental Study of the Effect of Drilling Fluid Penetration on Physical Properties of Gas Hydrate-Bearing Sediments in the Gulf of Mexico. *J. Pet. Sci. Eng.* **2020**, *187*,.
<https://doi.org/10.1016/j.petrol.2019.106832>.

294. Huang, T.; Zhang, Y.; Li, G.; Li, X.; Chen, Z. Numerical Modeling for Drilling Fluid Invasion into Hydrate-Bearing Sediments and Effects of Permeability. *J. Nat. Gas Sci. Eng.* **2020**, *77*,. <https://doi.org/10.1016/j.jngse.2020.103239>.
295. Zuev, V. M.; Porokhnyak, A. M.; Drozdov, A. V.; Sergievsky, V. V. Zaborovsky, V. V. Method of Storing Toxic Brines in Permafrost Environment (Published in Russian), 1988.
296. Porokhniak, A. M. *Gas Hydrates of the Permafrost Zone in Western Yakutia (Published in Russian)*; Publishing house of TsNIITsvetmet: Moscow, Russia, 1988; p. 30.
297. Porokhnyak, A. M.; Rassudov, A. V. *Disposal of Liquid Waste in Permafrost (Published in Russian)*; Nedra: Moscow, Russia, 1993; p. 111.
298. The process of melting ice with chemical reagents (published in Russian) https://bstudy.net/645082/tehnika/protsess_plavleniya_lda_himicheskimi_reagentami (accessed Mar 22, 2022).
299. Gaidaenko, E. P. Solubility of Ice in Frozen Soils under the Influence of Saline Solutions (Published in Russian). In *Problems of engineering glaciology*; Nauka: Novosibirsk, Russia, 1986; pp 32–36.
300. Pekhovich, A. I.; Shatalina, I. N. Experimental Studies of Ice Melting in Aqueous Solution (Published in Russian). In *Heat and mass transfer (volume 2)*; Minsk, 1968; pp 98–104.
301. Fedorov, A. M. Laboratory Studies of the Destruction of Ice and Ice-Rock Monoliths by Brines (Published in Russian). In *Conditions and processes of cryogenic migration of matter*; Yakutsk, Russia, 1989; pp 83–96.
302. Alekseev, S. *Cryogenesis of Groundwater and Rocks (on the Example of the Daldino-Alakitsky Region of Western Yakutia) (Published in Russian)*; SRC OIGGM SO: Novosibirsk,

- Russia, 2000; p. 119.
303. Borisov, V.; Alekseev, S. Factors of Interaction of Brines with Ice (Frozen Rock) at a Negative Temperature (Published in Russian). In *Fundamental problems of water and water resources at the turn of the III millennium*; NTL: Tomsk, Russia, 2000; pp 584–589.
304. Volkov, N. G.; Komarov, I. A.; Mironenko, M. V.; Fotiev, S. M. Methods for Assessing the Formation Temperature of the Ion-Salt Composition of Cryopegs (Published in Russian). *Kriosf. Zemli* **2005**, *9*, *4*, 54–61.
305. Farnam, Y.; Bentz, D.; Sakulich, A.; Flynn, D.; Weiss, J. Measuring Freeze and Thaw Damage in Mortars Containing Deicing Salt Using a Low-Temperature Longitudinal Guarded Comparative Calorimeter and Acoustic Emission. *Adv. Civ. Eng. Mater.* **2014**, *3*, *1*, 20130095:1-20130095:27. <https://doi.org/10.1520/acem20130095>.
306. Tyutyunov, I. A. *Introduction to the Theory of Permafrost Formation (Published in Russian)*; Academy of Science USSR: Moscow, Russia, 1960; p. 144.
307. Mantovani, S.; Valeri, S.; Loria, A.; Del Pennino, U. Viscosity of the Ice Surface Layer. *J. Chem. Phys.* **1980**, *72*, *2*, 1077–1083. <https://doi.org/10.1063/1.439248>.
308. Minlikaeva, O. V. Liquid-like Film on the Surface of Ice as a Medium for the Migration of Soluble Chemicals (Published in Russian). *Nat. Resour. Arct. Subarct.* **2007**, *4*, 103–105.
309. Chuvilin, E. M.; Bukhanov, B. A.; Mukhametdinova, A. Z.; Grechishcheva, E. S.; Sokolova, N. S.; Alekseev, A. G.; Istomin, V. A. Freezing Point and Unfrozen Water Contents of Permafrost Soils: Estimation by the Water Potential Method. *Cold Reg. Sci. Technol.* **2022**, *196*, 103488. <https://doi.org/10.1016/j.coldregions.2022.103488>.
310. Nechaev, E. A.; Kahn, E. V. Migration of Salts in Water-Saturated Sands at Negative

- Temperatures (Published in Russian). *Geochemistry* **1980**, 7,.
311. Romanov, V. P. Investigation of the Connection between Electrical Surface Properties and Ion Transport in Frozen Rocks (Published in Russian). In *Migration of chemical elements in permafrost*; Nauka: Novosibirsk, Russia, 1985.
312. Benediktova, N. A. Physical and Chemical Processes in Frozen Rocks during Their Interaction with Saline Solutions (Published in Russian), Moscow State University, 1992.
313. Chuvilin, E. M.; Smirnova, O. G. Migration of Chemical Elements in Frozen Rocks (Published in Russian). In *Proceedings of the First Conference of Geocryologists of Russia*; Moscow State University Publishing House: Moscow, Russia, 1996; pp 116–129.
314. Andersland, O. B.; Biggar, K. W. Site Investigations of Fuel Spill Migration into Permafrost. *J. Cold Reg. Eng.* **2002**, 13, 3, 165–166. [https://doi.org/10.1061/\(asce\)0887-381x\(1999\)13:3\(165\)](https://doi.org/10.1061/(asce)0887-381x(1999)13:3(165)).
315. Ershov, E. D.; Chuvilin, E. N.; Smirnova, O. G. Mobility of Ions of Chemical Elements in Ice and Frozen Rocks (Published in Russian). *Dokl. Akad. Nauk* **1999**, 367, 6, 796–798.
316. Tsyppkin, G. G. *Flows with Phase Transitions in Porous Media (Published in Russian)*; Fizmatlit: Moscow, Russia, 2009; p. 232, ISBN 978-5-9221-1102-7.
317. Lekhov, V. A.; Sokolov, V. N. Experimental Determination of the Hydraulic Conductivity and Diffusion Coefficient in Low Permeable Deposits (Published in Russian). *Geoecology. Eng. Geol. Hydrogeol. Geocryol.* **2017**, 3, 67–75.
318. Belousova, A. P. Preliminary Predictive Assessment of the Conditions of Pollution of the Unsaturated Zone by Oilfield Brines (Published in Russian). *Geoecology. Eng. Geol. Hydrogeol. Geocryol.* **1998**, 1, 75–89.

319. Malakhova, V. V. Mathematical Modeling of the Long-Term Dynamics of the Submarine Permafrost of the Arctic Shelf (Published in Russian). *Interexpo Geo-Siberia* **2014**, *4*, *1*, 1:1-1:5.
320. Tsyarkin, G. G. Effect of Liquid Phase Mobility on Gas Hydrate Dissociation in Reservoirs. *Fluid Dyn.* **1992**, *26*, *4*, 564–572. <https://doi.org/10.1007/BF01050319>.
321. Vasiliev, V. I.; Danilov, Y. G.; Ereemeev, I. S.; Popov, V. V.; Tsyarkin, G. G.; Yuzhui, S.; Yandong, C. Comparison of Mathematical Models of Heat and Mass Transfer in Soils (Published in Russian). *Bull. North-Eastern Fed. Univ.* **2013**, *10*, *4*, 5–10.
322. Aksenov, V. I.; Bubnov, N. G.; Klinova, G. I.; Iospa, A. V.; Gevorkyan, S. G. Phase Transformations of Water in Frozen Soils under the Cryopeg Influence (Published in Russian). *Geoecology. Eng. Geol. Hydrogeol. Geocryol.* **2010**, *1*, 40–51.
323. Galushkin, Y. I.; Sitar, K. A.; Frolov, S. V. Permafrost Formation and Degradation in the Urengoy and Kuyumbinskaya Areas of the Siberia. Part 1. Application of Basin Modeling System (Published in Russian). *Kriosf. Zemli* **2012**, *16*, *1*, 3–11.
324. Razumov, S. O.; Spektor, V. B.; Grigoriev, M. N. Model of the Post-Cenozoic Evolution of the Cryolithozone of the Shelf of the Western Part of the Laptev Sea (Published in Russian). *Oceanology* **2014**, *54*, *5*, 637–649.
325. Wei, J.; Cheng, Y.; Yan, C.; Li, Q.; Han, S.; Ansari, U. Decomposition Prevention through Thermal Sensitivity of Hydrate Formations around Wellbore. *Appl. Therm. Eng.* **2019**, *159*, 113921. <https://doi.org/10.1016/j.applthermaleng.2019.113921>.
326. Chuvilin, E. M.; Kozlova, E. V. Experimental Estimation of Hydrate-Containing Sediments Stability. In *5th International Conference on Gas Hydrates*; 2005.

327. Chuvilin, E. M.; Guryeva, O. M. Experimental Study of Self-Preservation Effect of Gas Hydrates in Frozen Sediments. In *Proceedings of the 9th International Conference on Permafrost, Fairbanks, AK, USA; 2008; Vol. 28.*
328. Trofimov, V. T.; Korolev, V. A. *Workshop on Soil Science (Published in Russian);* Moscow State University: Moscow, Russia, 1993; p. 390.
329. *GOST 12536-2014. Soils. Laboratory Determination of the Grain (Grain-Size) and Microaggregate Composition (Published in Russian);* Standartinform: Moscow, Russia, 2014.
330. 5180-84, G. *Soils. Methods for Laboratory Determination of Physical Characteristics / GOST of October 24, No. 5180-84; 1984.*
331. Shakhova, N.; Semiletov, I.; Gustafsson, O.; Sergienko, V.; Lobkovsky, L.; Dudarev, O.; Tumskey, V.; Grigoriev, M.; Chernykh, D.; Koshurnikov, A.; et al. Current Rates and Mechanisms of Subsea Permafrost Degradation in the East Siberian Arctic Shelf. *Nat. Commun.* **2017**, *8*, 15872:1-15872:13. <https://doi.org/10.1038/ncomms15872>.
332. Chuvilin, E.; Bukhanov, B.; Grebenkin, S.; Tysmskoy, V.; Shakhova, N.; Dudarev, O.; Semiletov, I. Thermal Conductivity of Bottom Sediments in the East Siberian Arctic Seas: A Case Study in the Buor-Khaya Bay. In *Proceedings of the 7th Canadian Permafrost Conference; Québec City, Québec, Canada; 2015; p ABS557:1-ABS557:6.*
333. Chuvilin, E.; Kozlova, E.; Makhonina, N. Experimental Investigation of Gas Hydrate and Ice Formation in Methane-Saturated Sediments. In *Proceedings of 8th International Conference on Permafrost; 2003; pp 145–150.*
334. Chuvilin, E.; Istomin, V.; Ekimova, V.; Bukhanov, B.; Hassanpouryouzband, A.; Yang, J.; Tohidi, B. Experimental Modeling of Methane Recovery by Flue Gas Injection Into Frozen

Hydrate-Bearing Reservoir. *Geomodel* **2018**.

335. Chuvilin, E. M.; Kozlova, E. V.; Skolotneva, T. S. Experimental Simulation of Frozen Hydrate-Containing Sediments Formation. In *Proceedings of the Fifth International Conference on Gas Hydrates*; Trondheim, Norway, 2005; pp 1561–1567.
336. Andreassen, K.; Hubbard, A.; Winsborrow, M.; Patton, H.; Vadakkepuliambatta, S.; Plaza-Faverola, A.; Gudlaugsson, E.; Serov, P.; Deryabin, A.; Mattingsdal, R.; et al. Massive Blow-out Craters Formed by Hydrate-Controlled Methane Expulsion from the Arctic Seafloor. *Science (80-.)*. **2017**, *356*, 6341, 948–953. <https://doi.org/10.1126/science.aal4500>.
337. Overduin, P.; Liebner, S.; Knoblauch, C.; Kneier, F. Subsea Permafrost Degradation and Inferred Methane Release in Shallow Coastal Water of the Central Laptev Sea. **2013**.
338. Chuvilin, E. M.; Grebenkin, S. I.; Sacleux, M. Influence of Moisture Content on Permeability of Frozen and Unfrozen Soils. *Kriosf. Zemli* **2016**, *20*, 3, 66–72.